

Floods of North-Central Indiana, July 2003

The storm front that moved across large parts of Indiana during the afternoon and evening of July 4, 2003, was more than a precursor to that night's Independence Day fireworks. The showers at first were seen by Hoosiers as a welcome source of moisture for lawns, gardens, and cropland. This storm, however, turned out to be the opening salvo of a 2-week onslaught of torrential rains that resulted in record-breaking floods in many basins. The heaviest rains were concentrated in north-central Indiana, July 5–10.

During the 2 weeks of downpours, water filled floodplains, overtopped roadways, and poured into homes. While flood waters ravaged Indiana communities, field crews from the U.S. Geological Survey (USGS) raced to gather vital hydrologic data. They measured the river flow and water levels to provide the information needed by river forecasters, engineers, and emergency managers who rely on data from the USGS network of about 170 Indiana streamflow-gaging stations (fig. 1). The network is funded in a cooperative program of federal, State, and local agencies, and is part of a national network maintained by the USGS. Many of these stations are equipped to transmit real-time streamflow data. In times of emergency, real-time streamflow information is a key element in the protection of lives and property. Real-time data are available to the public on the World Wide Web at

<http://in.waterdata.usgs.gov/nwis/rt>

New Peaks of Record

During the July 2003 floods, new records were set at 12 USGS streamflow-gaging stations in Indiana. The new peaks of record gage heights and new peaks of record discharges, along with other significant peaks, are listed in table 1.



Figure 1. Map of U.S. Geological Survey streamflow-gaging stations in Indiana, 2003. Stations in red set new record-peak discharge or gage height in July 2003 flood.

Significant Items:

- New peaks of record at 12 USGS streamflow-gaging stations in Indiana.
- 100-Year-Flood discharges on Deer Creek, Iroquois River, Kokomo Creek, St. Marys River, and the Wabash River.
- At White River at Noblesville, the water level exceeds the 1964 and 1991 record water levels by one-half foot, even though the July 2003 flood had 24 percent less discharge.
- Potential overall impact of flood is lessened as large volumes of water are dispersed to different major river basins, rather than concentrated toward a single major stream.

Discussion of the magnitude of the peaks follows.

Kokomo Creek and Wildcat Creek

Local news media reported nearly 10 inches of rain falling on the Kokomo area, July 4–5. The deluge flooded streets and backed up storm sewers. The runoff overwhelmed Kokomo Creek, a tributary to Wildcat Creek. The July 5 peak dis-

charge at the gaging station Kokomo Creek near Kokomo (03333600) was 2,940 ft³/s (cubic feet per second). This discharge was more than double the designated 100-year-flood discharge of 1,400 ft³/s. (A 100-year flood is that discharge for which there is a 1-percent chance of occurrence in any given year.) Coordinated flood frequencies for selected reaches of Indiana streams are available in the

Table 1. Peaks for period of record and for July 2003 at selected U.S. Geological Survey streamflow-gaging stations in Indiana[ft, foot; ft³/s, cubic foot per second; --, no data; >, greater than; <, less than; S.O., stage-only gaging station, beginning October 1, 2001; **red**, sites where new peak recorded July 2003; elevation of water surface can be obtained by adding the station datum to the gage height]

Station number	Station name	Station datum (ft)	Period of record, in water years ^a	Discharge peak of record (ft ³ /s) through 2002 water year	Gage-height peak of record (ft) through 2002 water year	Gage-height (ft) extreme outside period of record	Date and time of peak in July 2003	Discharge peak (ft ³ /s) in July 2003	Gage-height peak (ft) in July 2003	Recurrence interval
03322900	Wabash River at Linn Grove, IN	808.00	1965–2003	9,560 March 17, 1978	13.87 March 17, 1978	--	July 8 0500	14,500	14.76	>100 year
03322985	Wabash River near Bluffton, IN	795.42	2002–2003	4,640 April 1, 2002	12.88 April 1, 2002	--	July 9	15,300	18.43	60 year
03323500	Wabash River at Huntington, IN	700.04	1951–2003	S.O.	23.20 February 10, 1959	22.7 March 1913	July 11 0600	Unknown	19.37	--
03325000	Wabash River at Wabash, IN	642.66	1923–2003	49,600 May 18, 1943	24.44 February 11, 1959	28.7 March 26, 1913	July 7 1100	14,700	16.56	< 10 year
03329000	Wabash River at Logansport, IN	573.28	1923–2003 ^b	89,800 May 18, 1943	21.32 May 18, 1943	25.3 March 26, 1913	July 7 1200	37,100	13.60	< 10 year
03329700	Deer Creek near Delphi, IN	553.81	1944–2003	14,400 June 10, 1958	18.26 June 10, 1958	19.8 May 1943	July 5 2130	16,200	18.64	100 year
03331753	Tippecanoe River at Winamac, IN	674.19	2001–2003	4,600 May 16, 2002	11.48 May 16, 2002	--	July 11 0900	3,110	9.53	< 10 year
03333050	Tippecanoe River near Delphi, IN	535.00	1939–2003 ^b	20,600 April 12, 1994	13.72 April 12, 1994	--	July 10 0800	18,500	12.80	10 year
03333450	Wildcat Creek near Jerome, IN	820.04	1961–2993	7,120 July 14, 1992	13.71 December 30, 1990	18 March 1913	July 6 0015	7,180	14.35	30 year
03333600	Kokomo Creek near Kokomo, IN	807.68	1959–2003	1,040 April 20, 1964	9.88 April 20, 1964	--	July 5 0630	2,940	11.84	>100 year
03333700	Wildcat Creek at Kokomo, IN	775.62	1956–2003	8,100 February 10, 1959	16.95 December 30, 1990	--	July 5 1115	8,710	17.75	30 year
03335000	Wildcat Creek near Lafayette, IN	527.66	1954–2003	25,000 June 10, 1958	21.52 June 10, 1958	25.4 March 1913	July 7 0000	23,500	23.64	30 year
03335500	Wabash River at Lafayette, IN	504.14	1924–2003 ^b	131,000 May 19, 1943	28.47 May 19, 1943	32.9 March 26, 1913	July 10 1800	80,000	25.05	30 year
03336000	Wabash River at Covington, IN	473.97	1940–2003	147,000 May 20, 1943	32.44 May 20, 1943	35.1 March 1913	July 11 1000	89,000	28.90	30 year
03340500	Wabash River at Montezuma, IN	457.75	1928–2003	184,000 May 20, 1943	32.83 May 20, 1943	34.0 March 27, 1913	July 12 0900	108,000	30.35	15 year
03341500	Wabash River at Terre Haute, IN	445.78	1928–2003 ^b	189,000 May 20, 1943	30.50 May 20, 1943	31.2 March 27, 1913	July 13 1900	96,900	25.09	10 year
03347000	White River at Muncie, IN	917.10	1931–2003	14,300 April 21, 1964	21.07 January 15, 1937	22.6 March 1913	July 10 1000	6,950	10.94	< 10 year
03349000	White River at Noblesville, IN	738.16	1947–2003	27,000 December 31, 1990	21.31 April 22, 1964	--	July 10 1550	20,400	21.86	10 year
03351000	White River near Nora, IN	710.94	1930–2003	32,400 May 19, 1943	19.19 January 1, 1991	22.4 March 26, 1913	July 11 0100	24,900	17.78	10 year
03353000	White River at Indianapolis, IN	662.26	1930–2003 ^b	38,000 December 31, 1990	21.57 January 16, 1937	30.0 March 26, 1913	July 11 0300	27,900	17.19	< 10 year
03354000	White River near Centeron, IN	595.44	1947–2003 ^b	50,500 April 22, 1964	18.38 December 31, 1990	22.8 March 1913	July 12 0600	36,600	16.85	< 10 year
03360500	White River at Newberry, IN	465.59	1928–2003	105,000 November 18, 1993	25.87 November 18, 1993	27.5 March 27, 1913	July 15 2000	34,800	20.35	< 10 year
03374000	White River at Petersburg, IN	400.00	1928–2003	183,000 January 22, 1937	28.30 January 22, 1937	29.5 March 1913	July 18 0400	48,800	20.97	< 10 year
03377500	Wabash River at Mount Carmel, IL	369.46	1928–2003 ^b	305,000 May 25, 1943	32.35 May 17, 2002	33.0 March 30, 1913	July 20 0900	117,000	24.85	< 10 year
04181500	St. Marys River at Decatur, IN	760.44	1947–2003	11,300 February 10, 1959	24.40 March 14, 1982	--	July 9 0800	15,500	26.92	>100 year
04182000	St. Marys River near Fort Wayne, IN	748.97	1931–2003	13,600 February 11, 1959	19.66 March 14, 1982	--	July 9 2300	15,000	21.20	100 year
04183000	Maumee River at New Haven, IN	724.51	1957–2003 ^b	26,600 March 17, 1982	25.49 March 17, 1982	--	July 10 0900	19,400	21.58	< 10 year
05521000	Iroquois River at Rosebud, IN	661.47	1948–2003	656 December 30, 1990	8.86 February 10, 1959	--	July 21 1615	475	6.30	100 year
05522500	Iroquois River at Rensselaer, IN	642.29	1948–2003	2,550 June 10, 1958	16.54 June 10, 1958	--	July 10 1200	2,610	16.57	100 year
05524500	Iroquois River near Foresman, IN	624.00	1949–2003	5,930 June 14, 1958	24.42 June 14, 1958	--	July 8 1915	>5,210 (July 8)	24.87 (July 11)	Backwater ^c

^aWater year is the 12-month period October 1 through September 30. The 2003 water year, for example, includes October 1, 2002, through September 30, 2003.^bFragmented periods of data exist for other years.^cBackwater is the condition when water is backed up or retarded in its course, compared with its normal condition of flow.

Indiana Department of Natural Resources online publication, Coordinated Discharges of Selected Streams in Indiana at

http://www.state.in.us/dnr/water/surface_water/coordinated_discharges/index.html

The peak gage height for the station was 11.84 ft (feet), nearly 2 ft above the previous record set in 1964 (table 1). Most reaches of Wildcat Creek registered discharges in the range of a 30-year flood, as indicated by data collected at the gaging stations located near Jerome (03333450), at Kokomo (03333700), and near Lafayette (03335000).

St. Marys River

The St. Marys River at Decatur gaging station (04181500) recorded the peak gage height on July 9 at more than 2.5 ft above the previous peak of record in March 1982. The discharge was greater than a 100-year flood. Even though the St. Marys River peaked at the

St. Marys near Fort Wayne gaging station (04182000) at more than 1.5 ft above the 1982 flood level and the discharge was in the range of the 100-year flood, the city of Fort Wayne was spared a repeat of the 1982 devastation. In July 2003, the St. Joseph River, which flows from the north to join the St. Marys River to form the Maumee River, did not receive excessive amounts of precipitation. Consequently, the July 10 peak discharge at the Maumee River at New Haven gaging station (04183000) was less than a 10-year flood.

Wabash River

The July 8 crest at the Wabash River at Linn Grove gaging station (03322900) set new records. The peak gage height of 14.76 ft was almost 1 ft above the previous peak of record in March 1978. The peak discharge of 14,500 ft³/s was greater than a 100-year discharge.

The heavy rains generated a surge of water flowing down the upper reaches

of the Wabash River, as indicated by the flows at Linn Grove. Much of this surge was captured in J. Edward Roush Lake in Huntington. The peak discharges on July 7 were less than a 10-year flood at Wabash River at Wabash (03325000) and Wabash River at Logansport (03329000), about 22 and 55 miles, respectively, downstream from the lake. Farther downstream, the Wabash River at Lafayette gaging station (03335500) peaked on July 10 with discharges in the 30-year-flood range. The Wabash River at Covington gaging station (03336000) peaked on July 11 with discharges also in the 30-year-flood range.

The peaks attenuated as the flows continued downstream. At the Wabash River at Montezuma gaging station (03340500), where local police assisted USGS field crews in launching their boat, the river peaked on July 12 with discharges in the 15-year-flood range. The Wabash River at Terre Haute gaging station (03341500) peaked on July 13 with discharges in the 10-year-flood range.

Although the initial surge of water from the upper reaches of the Wabash River Basin was captured by J. Edward Roush Lake, continued large flows into the lake required lake managers to release water when the lake's holding capacity had been exceeded. The stage-only gaging station, Wabash River at Huntington (03323500), just below the outflow structure of the lake, recorded a peak gage height of 19.37 ft on July 11. This was the highest water level since the peak of record, 23.20 ft, February 10, 1959, which was prior to construction of the reservoir.

Deer Creek

The media reported loss of life in the Delphi area when high waters flooded homes and neighborhoods. At the Deer Creek near Delphi gaging station (03329700), the July 5 peak gage height of 18.64 ft exceeded the previous record peak of 18.26 ft, set in June 1958. The peak discharge of 16,200 ft³/s was in the range of a 100-year flood.



The Wabash River floods major roadways near the town of Bluffton, Indiana.



The Wabash River rises near the town of Montezuma, Indiana.



U.S. Geological Survey streamgager is shin deep in the Iroquois River as she inspects the gage near Foresman, Indiana.



Kokomo Creek flows over the bridge in the town of Kokomo, Indiana, as a U.S. Geological Survey streamgager prepares to make a discharge measurement.

Iroquois River

In the Iroquois River Basin, on the western side of the State, peaks gained in magnitude as flows moved downstream. The July 9 discharge at the Iroquois River at Rosebud gaging station (05521000) was in the range of a 40-year flood. By the time the Iroquois River peaked at the Rensselaer gaging station (05522500) on July 10, discharges were in the 100-year-flood range, eclipsing the June 1958 discharge. At the Iroquois River near Foresman gaging station (05524500), roads were submerged as water pooled up in the area faster than it could be drained, causing significant backwater conditions. Local citizens assisted USGS field crews by transporting them to the gaging station, using large trucks. The July 11 peak gage height at Foresman reached 24.87 ft, nearly one-half foot above the previous peak of record set in June 1958.

Intense rainfall several days later in the Wheatfield area caused the Iroquois River at Rosebud gaging station to peak on July 21 in the range of a 100-year-flood discharge. The effects of that later storm, however, seemed to lessen as the flow moved to the downstream stations.

White River

In parts of Marion and Hamilton Counties, large tracts of land along with homes and businesses were inundated

by White River flood waters. At the White River at Muncie gaging station (03347000), the July 10 peak discharge was less than a 10-year flood. Downstream, peak discharges at White River at Noblesville (03349000) and White River near Nora (03351000) gaging stations were in the 10-year-flood range. Farther downstream, the peak discharges diminished to less than a 10-year flood.

At the White River at Noblesville gaging station, a new record gage height was reached, even though the discharge was in the range of a 10-year flood. The previous peak-of-record gage height was on April 22, 1964, when the gage height reached 21.31 ft, with a discharge of 26,800 ft³/s. The peak-of-record discharge, 27,000 ft³/s, was on December 31, 1990, when the gage height reached 21.29 ft. On July 10, 2003, the stream peaked at a discharge of 20,400 ft³/s, with a new record gage height of 21.86 ft. Even though the peak discharge for the most recent flood was about 24 percent less than the two previous record peaks, the gage height crested about one-half foot higher.

Distribution of Flood Waters

The July 2003 flood was notable not only in the magnitude of the peaks but also in the number of separate basins affected. By the end of July, the Federal Emergency Management Agency (<http://www.fema.gov>) listed 40 Indiana coun-

ties as declared disaster areas. Although the destruction caused by the flooding was significant, the geographic patterns of the precipitation resulted in the dispersal of large volumes of water, thus averting a concentration of the runoff toward a single receiving stream. The flood waters of the Iroquois River drained west toward the Illinois River, and the flood waters on the St. Marys River drained east into Ohio by way of the Maumee River. The upper Wabash River and upper White River drained south, with the peaks attenuating as they moved downstream into areas where there had been less rainfall.

By Donald V. Arvin, Hieu T. Nguyen, and L. Mark Huff

For more information, please contact:

District Chief
U.S. Geological Survey
5957 Lakeside Boulevard
Indianapolis, IN 46278
317-290-3333
Fax: (317) 290-3313
E-mail: dc_in@usgs.gov

Additional information can be found by accessing the Indiana District Home Page on the World Wide Web at

<http://in.water.usgs.gov>