

ATTITUDES TOWARD MENTAL ILLNESS

Results From the Behavioral Risk Factor
Surveillance System



Suggested Citation

Centers for Disease Control and Prevention, Substance Abuse and Mental Health Services Administration, National Association of County Behavioral Health & Developmental Disability Directors, National Institute of Mental Health, The Carter Center Mental Health Program. *Attitudes Toward Mental Illness: Results from the Behavioral Risk Factor Surveillance System*. Atlanta (GA); Centers for Disease Control and Prevention; 2012.

Contributing Authors

Rosemarie Kobau, MPH, MAPP¹
Public Health Advisor
Centers for Disease Control and Prevention

Matthew M. Zack, MD, MPH¹
Medical Epidemiologist
Centers for Disease Control and Prevention

Cecily Luncheon, MD, DrPH, MPH¹
ORISE Fellow
Centers for Disease Control and Prevention

John P. Barile, PhD¹
ORISE Fellow
Centers for Disease Control and Prevention

Chris Marshall⁴
Special Assistant to the Director
Center for Mental Health Services
Substance Abuse and Mental Health Services
Administration

Thomas Bornemann, PhD⁵
Director, The Carter Center Mental Health Program

Emeline Otey, PhD⁷
Office of Research Training and Career Development
Division of Services and Intervention Research
National Institute of Mental Health

Robin K. Davis, PhD²
ICF Macro

Lucas Godoy Garraza, MS²
ICF Macro

Christine Walrath, PhD²
ICF Macro

Ron Manderscheid, PhD³
Executive Director, National Association of
County Behavioral Health & Developmental
Disability Directors

Rebecca G. Palpant, MS⁵
Assistant Director, The Rosalynn Carter
Fellowships for Mental Health Journalism
The Carter Center Mental Health Program

Diana S. Morales, MPH⁶
Public Health Analyst
Office of Constituency Relations and Public
Liaison
National Institute of Mental Health

Acknowledgments

Kurt Greenlund, PhD¹
Acting Director, Prevention Research Centers Program
Centers for Disease Control and Prevention

Susan L. Jernick, DPM, MPH¹
Contractor, DB Consulting Group
Centers for Disease Control and Prevention

Deborah Leiter⁸
Vice President, Campaign Director
The Advertising Council

Adrienne Ziluca⁸
Campaign Manager
The Advertising Council

¹ Division of Population Health, 4770 Buford Highway NE, Atlanta, GA 30341

² ICF Macro, 3 Corporate Square, NE, Suite 370, Atlanta, GA 30329

³ National Association of County Behavioral Health & Developmental Disability Directors,
25 Massachusetts Ave, NW, Suite 500, Washington, DC 20001

⁴ Center for Mental Health Services, Substance Abuse and Mental Health Services Administration,
1 Choke Cherry Road, 6-1059, Rockville, MD 20857

⁵ One Copenhill, 453 Freedom Parkway, Atlanta, GA 30307

⁶ 6001 Executive Boulevard, Bethesda, MD 20892-9629

⁷ 6001 Executive Boulevard, Bethesda, MD 20892-9669

⁸ The Advertising Council, 1707 L Street, NW, Suite 600, Washington, DC 20036

CONTENTS

- Executive Summary 1
 - Introduction to Using This Report 2
- Background..... 3
 - Attitudes Toward Mental Illness and Stigma 3
 - What Are the Consequences of Negative Attitudes Toward Mental Illness and Stigma? 4
 - Why Is It Important to Track Attitudes Toward Mental Illness? 5
- Methodology 6
 - BRFSS 6
 - Measures 6
 - Individual Level Predictors..... 7
 - State-level Predictors 7
 - What a Difference a Friend Makes Campaign Overview 8
 - Analysis 8
- Key Findings in Surveyed States 10
 - Overall Study Results 10
 - Treatment Effectiveness 10
 - People Are Generally Caring and Sympathetic to People With Mental Illness 10
 - Discussion 53
 - Strategies for Combating Stigma 54
 - Limitations 54
 - Acknowledgment 55
- Resources for States to Address Mental Illness Stigma 55
 - Federal Resources 55
 - Nonfederal Resources 55
- References 56
- Appendix A: Mental Health-related and BRFSS Indicators 59
- Appendix B: State-specific Data Tables 64



EXECUTIVE SUMMARY

In 2005, the Centers for Disease Control and Prevention (CDC) and the Substance Abuse and Mental Health Services Administration (SAMHSA) collaborated to track state-level attitudes toward mental illness. Following recommendations from the 2002 President's New Freedom Commission on Mental Health, these agencies sought to examine public perceptions regarding treatment effectiveness and perceptions regarding people being caring and sympathetic to people with mental illness. This collaboration paralleled the release of a SAMHSA national anti-stigma campaign, *What a Difference a Friend Makes*, designed to help young adults support friends with a mental health problem. Two questions that asked about attitudes toward mental illness, along with other questions about mental illness symptoms, were included on the CDC-supported Behavioral Risk Factor Surveillance System (BRFSS). These questions comprised the BRFSS Mental Illness and Stigma Module. In 2007, 35 states, the District of Columbia, and Puerto Rico received SAMHSA support to collect data using the BRFSS Mental Illness and Stigma Module, and, in 2009, 16 states received support to do so. While a 2010 CDC study published in the *Morbidity and Mortality Weekly Report* (CDC, 2010) described some limited findings, the goals of this study were to expand on the previous analysis to (1) provide state-level estimates of attitudes toward mental illness by select socio-demographic factors, mental illness symptoms, and mental health treatment; (2) examine individual (e.g., age, race/ethnicity) and state-level predictors (e.g., per capita expenditures on mental health services; the average annual unemployment rate; and exposure to the *What a Difference a Friend Makes* campaign) on attitudes toward mental illness; (3) highlight population subgroups who strongly disagree with the statements (i.e., a vulnerable group) for each state; and (4) provide resources and strategies for targeting these groups, and combating stigma in general.

As more individuals seek help and share their stories with friends and relatives, compassion will be the response, not ridicule (President's New Freedom Commission on Mental Health, p. 7, 2003).

This study found that most adults (>80%) in the states surveyed agreed that mental illness treatment is effective, but substantially fewer adults (35%–67%) agreed that people are caring and sympathetic to people with mental illness. Some population subgroups (e.g., black, non-Hispanic adults, Hispanic adults, those with less than a high school education) were more likely to strongly disagree that treatment is effective. Women, adults with chronic disease (e.g., arthritis, heart disease), and adults who were unemployed or unable to work were more likely to strongly disagree that people are caring and sympathetic to people with mental illness. In general, adults with mental illness symptoms, including those receiving treatment for a mental health problem were less likely to agree that people are caring and sympathetic to people with mental illness.

Adults who lived in states with higher per capita expenditures on mental health services were more likely to agree that treatment is effective, and were more likely to report receiving treatment. Adults who lived in areas with more mental health professionals were more likely to agree that other people are caring and sympathetic to people with mental illness. Young adults (ages 18–24) who lived in states with greater donated media time for SAMHSA's *What a Difference a Friend Makes* campaign were more likely to agree that people are caring and sympathetic to people with mental illness.

These BRFSS data demonstrate the feasibility and usefulness of tracking attitudes toward mental illness at the state level as well as for assessment, program development, and evaluation of public health anti-stigma activities.



Introduction to Using This Report

This report examines individual and state-level predictors of attitudes toward mental illness to highlight subgroups that disagreed with the attitudinal statements and, therefore, might benefit from educational programs and services. The report's eight sections include the following:

- 1) **Background:** This section provides information about stigma as it interferes with public health prevention efforts and the rationale for the selection of study questions.
- 2) **Study Methodology:** This section describes the source of questions used for this study, participating states, data collection methods, individual-level predictors, and state-level predictors (Appendix A) used in the analysis as well as analytical strategies used.
- 3) **Key Findings:** This section briefly summarizes the main study findings.
- 4) **State-Specific Findings (Graphs):** This section provides a one-page graphical overview of key findings on total disagreement for each attitudinal question by the state overall and by age, sex, race/ethnicity, serious psychological distress (SPD), frequent mental distress (FMD), and receipt of mental health treatment for each state participating in this study. This section also includes a *Call to Action* subsection that highlights groups found to significantly differ in their responses from the state average. Additionally, this section provides resources targeting these groups.
- 5) **Discussion:** This section compares and contrasts these findings in relation to other research, reviews the implications of these findings, and describes study limitations.
- 6) **Resources:** This section provides links to resources for those interested in strategies to combat mental illness stigma.
- 7) **References:** This section cites other published studies and resources for those who wish to delve further.
- 8) **State-Specific Findings (Tables [Appendix B]):** This section tabulates detailed results by state for attitudes toward mental illness by age, sex, race/ethnicity, educational level, household income, employment status, veteran status, diabetes, cardiovascular disease, asthma, arthritis, FMD, SPD, and receipt of mental health treatment (adjusted for sex, age, racial/ethnic group, education, and household income).

Understanding attitudes toward mental illness at the state level can help identify and inform priorities that support the efforts of mental health state agencies, providers, policymakers, educators, and others to reduce stigma. The collection and the analysis of state-level data can provide valuable insight into the unique needs of various subgroups, such as racial and ethnic minorities as well as people living with chronic disease and co-occurring mental illness. It is our hope that this information can help shape initiatives effective in reducing stigma and removing barriers for those seeking or receiving treatment for mental illness.



BACKGROUND

Attitudes Toward Mental Illness and Stigma

People’s beliefs and attitudes toward mental illness set the stage for how they interact with, provide opportunities for, and help support a person with mental illness. People’s beliefs and attitudes toward mental illness also frame how they experience and express their own emotional problems and psychological distress and whether they disclose these symptoms and seek care. About one in four U.S. adults (26.2%) age 18 and older, in any given year, has a mental disorder (e.g., mood disorder, anxiety disorder, impulse control disorder, or substance abuse disorder) (Kessler, Chiu, Demler, & Walters, 2005), meaning that mental disorders are common and can affect anyone. Many adults with common chronic conditions such as arthritis, cancer, diabetes, heart disease, and epilepsy experience concurrent depression and anxiety—further complicating self-management of these disorders and adversely affecting quality of life (Chapman et al., 2005; El-Gabalawy et al., 2010; IOM, 2012).

Attitudes and beliefs about mental illness are shaped by personal knowledge about mental illness, knowing and interacting with someone living with mental illness, cultural stereotypes about mental illness, media stories, and familiarity with institutional practices and past restrictions (e.g., health insurance restrictions, employment restrictions; adoption restrictions) (Corrigan et al., 2004; Wahl, 2003). When such attitudes and beliefs are expressed positively, they can result in supportive and inclusive behaviors (e.g., willingness to date a person with mental illness or to hire a person with mental illness). When such attitudes and beliefs are expressed negatively, they may result in avoidance, exclusion from daily activities, and, in the worst case, exploitation and discrimination.

Stigma has been described as “a cluster of negative attitudes and beliefs that motivate the general public to fear, reject, avoid, and discriminate against people with mental illnesses” (President’s New Freedom Commission on Mental Health, p. 4, 2003). When stigma leads to social exclusion or discrimination (“experienced” stigma), it results in unequal access to resources that all people need to function well: educational opportunities, employment, a supportive community, including friends and family, and access to quality health care (Link & Phelan, 2001; Corrigan et al., 2004). These types of disparities in education, employment, and access to care can have cumulative long-term negative consequences.

The prevailing view of health-related stigma is that it refers to perceived, enacted, or anticipated avoidance or social exclusion, and not to an individual blemish or mark
(Weiss, Ramakrishna & Somma 2006; Goffman 1963).

For example, a young adult with untreated mental illness who is unable to graduate from high school is less likely to find a good paying job that can support his or her basic needs, including access to health care. These disadvantages can cause a person to experience more negative outcomes. Being unemployed, living at or below the poverty line, being socially isolated, and living with other social disadvantages can further deflate self-esteem, compounding mental illness symptoms, and add to the burden of stigma (Sartorius, 2005). Sometimes stigma is simply “felt” in the absence of being discriminated against and results from internalizing perceived negative attitudes associated with a characteristic (e.g., age), a disorder (e.g., HIV-AIDS), a behavior (e.g., smoking), or other factor (e.g., place of birth).



Whether stigma is experienced as social exclusion or discrimination or felt as a pervasive and underlying sense of being different from others, it can be debilitating for people and poses a challenge for public health prevention efforts. Different opinions exist regarding the implications of different labels associated with describing mental illness (e.g., brain disease) and felt or experienced stigma (Sayce, 1998; Corrigan & Watson 2004; Corrigan, Markowitz, & Watson, 2004; Pescosolido et al., 2010). However, the prevailing view of health-related stigma is that it refers to perceived, enacted, or anticipated avoidance or social exclusion, and not to an individual blemish or mark (Weiss, Ramakrishna, & Somma, 2006; Goffman, 1963). Different methods exist for measuring health-related stigma, and challenges and limitations associated with distinguishing between felt versus experienced stigma in attitudinal research have been described (Link et al., 2004; Green, 1995).

What Are the Consequences of Negative Attitudes Toward Mental Illness and Stigma?

Only about 20% of adults with a diagnosable mental disorder (Wang et al., 2005) or with a self-reported mental health condition (Hennessy et al., 2012) saw a mental health provider in the previous year. Embarrassment associated with accessing mental health services is one of the many barriers that cause people to hide their symptoms and to prevent them from getting necessary treatment for their mental illness symptoms (U.S. Department of Health and Human Services, 1999; 2001; Greene-Shortridge et al., 2007; Nadeem et al., 2007). Stigma poses a barrier for public health primary prevention efforts designed to minimize the onset of mental illness, as well as with secondary prevention efforts aimed at promoting early treatment to prevent worsening of symptoms over time (Weiss, Ramakrishna, & Somma, 2006).

Stigma can also interfere with self-management of mental disorders (tertiary prevention) (Sirey et al., 2001). Untreated symptoms can have grave consequences for people living with mental illness and negatively impact families affected by these disorders. For example, most people with serious and persistent mental illness (mental disorders that interfere with some area of social functioning) are unemployed and live below the poverty line, and many face major barriers to obtaining decent, affordable housing (U.S. Department of Health and Human Services, 1999). These individuals may need a number of additional social supports (e.g., job training, peer-support networks) to live successfully in the community, but such supports may not be available. Other individuals with depression and anxiety might avoid disclosing their symptoms and instead adopt unhealthy behaviors to help them cope with their distress (e.g., smoking, excessive alcohol use, binge-eating). These behaviors can increase their risk for developing chronic diseases, worsening their overall health over time. Recent studies have found an increased risk of death at younger ages for people with mental illness (Colton & Manderscheid, 2006).

- **Stigma about mental illness can lead people to fear disclosing that they have mental health problems, which may prevent treatment and recovery.**
- **Stigma can result in limited life opportunities.**
- **Stigma poses barriers for public health prevention efforts designed to minimize onset of mental illness and the prevention or worsening of symptoms over time.**
- **Stigma can result in lower prioritization for public resources allocated to mental health services and poorer quality of care delivered to people with mental illness.**



Attitudes toward mental illness can also influence how policymakers allocate public resources to mental health services, pose challenges for staff retention in mental health settings, result in poorer quality of medical care administered to people with mental illness, and create fundraising challenges for organizations who serve people with mental illness and their families (Kadri & Sartorius, 2005; Pescosolido et al., 2010; Stuart, 2005). State-level factors such as unemployment levels and access to mental health services, and the presence or absence of other state resources may affect public attitudes and merit study.

Why Is It Important to Track Attitudes Toward Mental Illness?

People's attitudes and beliefs predict their behavior (Ajzen & Fishbein, 1980). People's beliefs and attitudes about mental illness might predict whether they disclose their symptoms and seek treatment and support. Knowledge and beliefs that can aid in the recognition, management, or prevention of mental health disorders are defined as mental health literacy (Jorm et al., 1997). Tracking attitudes toward mental illness can serve as an indicator of the public's mental health literacy. For example, in a 1996 study, 54% of the U.S. public attributed major depression to neurobiological causes, and this increased to 67% in 2006 (Pescosolido et al., 2010). Similarly, a larger percentage of people endorsed the benefits of treatment by a physician for people with major depression in 2006 (91%) than in 1996 (78%) (Pescosolido et al., 2010). However, improvements in neurobiological understanding of mental illness were unrelated to negative attitudes and, in some cases, increased the odds of negative attitudes (e.g., need for social distance, perceived dangerousness) (Pescosolido et al., 2010). In another study of U.S. adults, only about one fourth agreed that people are caring and sympathetic to people with mental illness (Kobau, DiIorio, Chapman, & Delvecchio, 2010). When asked about how much it would be worth to avoid mental illness compared to general medical illnesses, the public was less willing to pay to avoid mental health treatment than they were to pay to avoid physical health treatment (Smith et al., 2012). These studies provide important snapshots of attitudes toward mental illness across the country; however, studies that examine attitudes in depth such as distinguishing between attitudes relative to perceived or experienced stigma, studies that link attitudes to actual behavior, or studies that track attitudes toward mental illness at the state level do not occur routinely. These limited, cross-sectional studies tell us little about how attitudes shift in relation to historical events (e.g., media over-sensationalization of the rare violence associated with a person with mental illness), how attitudes shift over time in the same people, and how these attitudes differ within a state relative to characteristics of the state such as the average unemployment rate or per capita expenditures on state mental health agencies.



METHODOLOGY

This section describes the source of questions used in this study, data collection methods, individual and state-level predictors used in the analysis, and the analytical strategy.


BRFSS

The BRFSS is a state-based system of health surveys that collects information on health risk behaviors, preventive health practices, and health care access primarily related to chronic disease and injury (CDC, 2012). BRFSS was established in 1984 by the CDC; currently data are collected monthly in all 50 states, the District of Columbia, Puerto Rico, the U.S. Virgin Islands, and Guam (<http://www.cdc.gov/brfss/>). The BRFSS Mental Illness and Stigma Module was developed by CDC and SAMHSA to track the prevalence of serious psychological distress, mental health treatment in past 12 months, and attitudes toward mental illness in U.S. states. In 2007, 35 states, the District of Columbia, and Puerto Rico included the Mental Illness and Stigma Module on the BRFSS. In 2009, 16 states did so. Fourteen states supported the Mental Illness and Stigma Module in both 2007 and 2009. The latest data available for each state were used for analysis (21 states, the District of Columbia, and Puerto Rico with data from 2007 and 16 states with data from 2009); (Tables 1-4). In both years, BRFSS sampled adults by landline telephone. Among the 35 states, the District of Columbia, and Puerto Rico surveyed in 2007, the median Council of American Survey Research Organization (CASRO) response rate was 51% and the CASRO cooperation rate was 71.4% (CDC, 2008). In 2009, among the 16 states surveyed, the CASRO median response rate was 55.3% and the CASRO cooperation rate was 77% (CDC, 2011).

Measures

Attitudes Toward Mental Illness

Attitudes were assessed by asking respondents to indicate their level of agreement with two statements. The first statement assessed attitudes toward the effectiveness of treatment: “Treatment can help people with mental illness lead normal lives.” The second statement assessed the respondent’s perception of public attitudes toward persons with mental illness: “People are generally caring and sympathetic to people with mental illness” (The Scottish Government, 2009). This method of asking respondents to indicate what other people think about a health condition has been previously used in assessing other health-related stigma (Green, 1995). For the two statements, participants were asked to answer whether they agreed strongly, agreed slightly, neither agreed nor disagreed, disagreed slightly, or disagreed strongly. Before inclusion in BRFSS, cognitive testing confirmed that adults understood these questions as intended. The question about attitudes toward treatment also demonstrated acceptable construct validity with expectations regarding mental illness recovery (Kobau, 2010). In this study, *total disagreement* (seen in graphs, Section 4) was defined with responses including either slightly or strongly disagree.



Serious Psychological Distress

The Kessler 6-scale asks respondents how often in the past 30 days they felt six symptoms of mental illness (i.e., feeling nervous, depressed, hopeless, restless, like a failure, like everything was an effort). Each item is scored on a 5-point scale indicating frequency, ranging from 0 (none of the time) to 4 (all of the time), and summed (score range: 0–24). Respondents scoring 13 or more on this scale were classified as having serious psychological distress (SPD) (Kessler et al., 2003). The K6 is a valid screen for serious mental illness in the general population (Kessler et al., 2003).

Mental Health Treatment

Another question was asked of respondents: “Are you now taking medicine or receiving treatment from a doctor or other health professional for any type of mental health condition or emotional problem?” Response options included yes, no, don’t know, and refused. Adults who responded yes were classified as currently receiving treatment for a mental health or emotional problem.

Frequent Mental Distress (FMD)

FMD was measured with the question, “For how many days in the past 30 days was your mental health (due to stress, depression, or problems with emotions) not good?” Respondents reporting 14 or more not good mental health days were identified as having frequent mental distress (Zahran et al., 2004).


Individual Level Predictors

The BRFSS is the source of all the individual-level predictors, which includes sociodemographic characteristics (sex, age, racial/ethnic group, education, income, and veteran status); self-reported doctor-diagnosed chronic disease (diabetes mellitus, cardiovascular disease [including myocardial infarction, stroke, coronary heart disease], asthma, and arthritis); and self-reported mental health symptoms (frequent mental distress [FMD], serious psychological distress [SPD], and being under medical care for an emotional issue).

State-level Predictors

State-level predictors were obtained from different sources: (1) State mental health agency expenditures per capita for fiscal year 2004 (latest data available) from the National Association of State Mental Health Program Directors Research Institute (NRI); (2) the percentage of the population living in mental health professionals shortage areas from the Health Resources and Services Administration (as of December 2010); (3) the annual average unemployment rate from the Bureau of Labor Statistics; d) health insurance coverage for each year from the U.S. Census Bureau; (4) the percentage of youth aged 18–24 for each year from the BRFSS; (5) per capita values of donated media (local broadcast TV) from 210 media markets across the U.S. for the What a Difference a Friend Makes campaign (cumulative since its inception in 2006 throughout the end of each year) provided by SAMHSA and the Ad Council. (Appendix A, Table A).¹

¹ State-level data were not available for Puerto Rico for all predictors (e.g., donated media) and, thus, are not presented in Table A.



What a Difference a Friend Makes Campaign Overview

In December 2006, the Ad Council, in partnership with SAMHSA, launched a national public service advertising campaign that asked 18- to 25-year-olds to step up and support friends they know are experiencing a mental health problem by demonstrating the roles they can play in their friend’s recovery. The public service announcement (PSA) campaign includes television, Web videos, radio, outdoor, print and Web elements, as well as an online community that young adults can join to share their stories and experiences with recovery and support. Viewers and listeners are encouraged to continue to support their friends who are living with mental health problems and to visit the campaign website to learn more (<http://www.whatadifference.samhsa.gov/>).

Analysis

All the analyses excluded persons who responded “did not know” or refused to answer these questions.²

Raw and Adjusted Percentages

For each jurisdiction, the proportions of responses in each attitude category were estimated by a set of subpopulations or domains of interest (defined by sociodemographic characteristics, self-reporting chronic disease, and mental health symptoms). Both point estimates and standard errors were computed taking into account the complex sample design (sample weights, stratification, and clustering). Adjusted proportions, or predictive marginals (Korn & Graubard, 1999), were obtained with SUDAAN multinomial logistic regression models. Level of agreement with each attitude statement served as dependent variables controlling for sex, age, racial/ethnic group, education, and household income. The analysis was performed using SUDAAN 10 (Research Triangle Institute, 2008). Within each subgroup of sex, age, and race/ethnicity, we examined all possible comparisons to identify groups that differed significantly from each other (e.g., State-Specific Findings overview paragraph).

Multilevel Models

To explore predictors (e.g., per capita expenditures on state mental health agencies) of differences in attitudes across states,³ a multilevel logistic regression model was fitted to the combined sample. A random intercept was used to account for clustering of observations within each jurisdiction year. Additional clustering within each jurisdiction across years was taken into account in the computation of standard errors. The multilevel models were fitted by pseudo-maximum likelihood to incorporate sample weights (Pfeffermann & Sverchkov, 2009). Ordinal and binary logistic regressions were used, the latter arising as a result of dichotomizing the attitudes variables depending on whether the respondent indicated agreement (either slight or strong). The analysis was performed using Mplus 6 (Muthén & Muthén, 1998–2011).

² For each question, about 2% of respondents answered “did not know,” and about 0.3% of respondents refused to answer each question.

³ Puerto Rico was omitted from multilevel analysis because not all state-level predictors examined (e.g., donated media) were available.



Multilevel model parameter estimates were examined to determine statistically significant associations. Figures 1 through 3 in Appendix A depict how much the percent disagreement that treatment can help those with mental illness, the percent disagreement that people are caring and sympathetic to people with mental illness, and the percent of adults receiving mental health treatment vary with the quantity of each state-level predictor (e.g., per capita expenditures, percentage of population living in mental health professional shortage areas), from lowest value to highest value. The three horizontal bars in each figure represent the average level of agreement for these three outcomes at the 25th, 50th, and the 75th percentile of the values of these state-level predictors. Each figure depicts the statistically significant associations found between each state-level predictor and the outcome.

Additional Analyses

Estimates for the “Overall” category in the state-specific results (i.e., graphs) were obtained by age and race standardization of the BRFSS datasets to the 2007 U.S. population distribution. The estimates are provided for comparison purposes. All of the states in the sample are used in the estimation, and sample weights are taken into account in the estimation.

Calls to Action boxes highlight significant differences ($p \leq 0.05$) between subgroups compared to the average state percent in the “**strongly disagree**” categories (potentially a high risk group) for each question on attitudes. (Supporting data are provided in Appendix B.)



KEY FINDINGS IN SURVEYED STATES⁴

This section briefly summarizes the main findings. Readers can refer to the sections that follow for supporting data.

Overall Study Results

Across states surveyed, most adults (>80%) agreed (slightly or strongly) that treatment can help people living with mental illness lead normal lives (Tables 1-2). Yet, fewer adults (35–67%) agreed that other people are caring and sympathetic to people living with mental illness (Tables 3-4). While no regional differences were identified in the analysis, states differed in overall level of agreement with both statements after controlling for individual level differences.⁵

Treatment Effectiveness

- ▶ The vast majority of adults (>80%) in the states surveyed agreed that treatment is effective (Tables 1-2).
- ▶ Adults who reported receiving mental health treatment were generally more likely to strongly agree that treatment is effective (e.g., CO, HI, KS, NH, TX; see Appendix B tables).
- ▶ The probability of adults receiving mental health treatment increased when states provided more funding for their state mental health agencies (see Appendix A, Figure 3).
- ▶ Some population subgroups (e.g., non-Hispanic blacks, Hispanics, those with less than a high school education,) in some states were more likely to strongly disagree that treatment is effective (e.g., GA, CA, MA, WA; see Appendix B tables).
- ▶ Higher per capita expenditure on state mental health agencies was associated with less disagreement with “Treatment can help people with mental illness lead normal lives.” (See Appendix A, Figure 1.)

People Are Generally Caring and Sympathetic to People With Mental Illness

In general, smaller percentages of people in the states surveyed agreed that people are generally caring and sympathetic to people with mental illness (range 35%–67%) [Tables 3-4] than that treatment is effective (Tables 1-2).

- ▶ Adults with mental illness symptoms (i.e., serious psychological distress [SPD], frequent mental distress [FMD]) and those reporting that they are currently receiving treatment for a mental illness or an emotional problem more often strongly disagreed that people are generally caring and sympathetic to people with mental illness (Appendix B, tables).
- ▶ Adults living with existing chronic conditions (e.g., arthritis, heart disease, asthma) were also more likely to strongly disagree that people are generally caring and sympathetic to those with mental illness than adults living without these conditions (Appendix B, tables).

⁴ Key findings reflect results from the surveyed states and are not national estimates.

⁵ Estimates may differ slightly from the CDC 2010 MMWR report due to the inclusion of additional control variables in this study.



- ▶ In some states, population subgroups such as women, adults who were unable to work or were out of work, adults living in households earning <\$20,000/year), and adults with less than a high school education were more likely to strongly disagree that people are caring and sympathetic to people with mental illness (e.g., CO, KS, NE, MT, WI; see Appendix B tables).
- ▶ Adults who lived in areas with fewer mental health professionals were more likely to disagree that “People are caring and sympathetic to people with mental illness.” (See Appendix A, Figure 2.)
- ▶ Higher values of donated media for the What a Difference a Friend Makes campaign were associated with less disagreement with “People are caring and sympathetic to people with mental illness,” in states with greater percentages of younger adults (ages 18–24). (See Appendix A, Figure 2.)




Table 1. Level of Agreement* with the Statement, *Treatment Can Help People with Mental Illness Lead Normal Lives*, by State and Territory—2007 BRFSS.

State	Unweighted Sample Size	Disagree Strongly	Disagree Slightly	Neither Agree Nor Disagree	Agree Slightly	Agree Strongly
Alaska	2,224	1.5%	4.0%	1.5%	26.6%	66.5%
Arkansas	5,119	2.1%	3.4%	0.6%	23.9%	70.0%
Colorado	5,241	1.4%	4.2%	0.8%	27.0%	66.5%
Connecticut	6,416	1.5%	3.0%	2.1%	23.8%	69.6%
District of Columbia	3,336	1.4%	2.8%	1.4%	20.2%	74.2%
Illinois	4,947	1.5%	4.2%	0.9%	27.9%	65.4%
Indiana	5,254	1.6%	3.6%	1.3%	26.8%	66.7%
Iowa	4,804	1.7%	3.5%	0.6%	26.3%	67.9%
Kentucky	5,916	2.0%	1.5%	5.3%	25.1%	66.1%
Louisiana	5,837	2.1%	3.4%	1.8%	17.1%	75.6%
Maine	3,734	1.8%	3.4%	1.0%	23.8%	70.0%
Minnesota	4,669	1.3%	3.5%	3.3%	28.7%	63.1%
Montana	5,233	2.0%	3.8%	3.4%	26.4%	64.3%
New Hampshire	5,310	1.3%	3.7%	2.1%	25.9%	67.0%
New Mexico	5,672	1.6%	3.9%	2.5%	25.1%	67.0%
Ohio	4,853	1.4%	3.9%	1.4%	23.9%	69.3%
Oklahoma	6,487	1.2%	3.0%	0.8%	24.1%	70.8%
Oregon	1,833	1.1%	2.5%	1.4%	22.1%	72.8%
Puerto Rico	3,717	0.8%	3.7%	8.9%	57.2%	29.4%
Rhode Island	3,820	1.7%	4.1%	3.0%	29.9%	61.3%
Texas	7,045	1.5%	4.2%	4.3%	26.4%	63.6%
Virginia	5,159	2.7%	3.2%	1.9%	22.0%	70.3%
Wisconsin	4,241	1.7%	4.4%	0.7%	30.3%	62.9%

*Adjusted for sex, age group, racial/ethnic group, education and household income level. Estimates are weighted; sample size is unweighted.

Table 2. Level of Agreement* with the Statement, *Treatment Can Help People with Mental Illness Lead Normal Lives*, by State and Territory—2009 BRFSS.

State	Unweighted Sample Size	Disagree Strongly	Disagree Slightly	Neither Agree Nor Disagree	Agree Slightly	Agree Strongly
California	3,627	1.4%	4.8%	0.9%	29.3%	63.5%
Georgia	5,181	1.3%	3.7%	2.0%	23.7%	69.3%
Hawaii	6,026	1.4%	3.4%	1.8%	27.5%	66.0%
Kansas	8,410	0.9%	3.1%	3.8%	27.1%	65.0%
Massachusetts	4,552	1.2%	3.4%	2.8%	23.7%	68.9%
Michigan	2,722	1.8%	3.8%	1.3%	23.5%	69.5%
Mississippi	9,911	1.9%	3.3%	1.3%	23.0%	70.4%
Missouri	4,499	1.4%	2.7%	1.4%	24.0%	70.5%
Nebraska	4,730	1.6%	5.7%	1.8%	29.5%	61.4%
Nevada	3,421	1.9%	4.3%	1.8%	27.2%	64.8%
South Carolina	8,761	1.4%	3.8%	1.1%	25.6%	68.1%
Tennessee	2,124	1.3%	2.9%	3.0%	17.2%	75.5%
Utah	2,363	1.2%	4.1%	1.0%	26.9%	66.8%
Vermont	6,096	1.2%	3.1%	1.5%	25.2%	69.0%
Washington	6,883	1.8%	3.5%	1.3%	22.8%	70.6%
Wyoming	5,482	2.1%	4.9%	0.9%	27.1%	65.1%

*Adjusted for sex, age group, racial/ethnic group, education and household income level. Estimates are weighted; sample size is unweighted.



Table 3. Level of Agreement* with the Statement, *People are Generally Caring and Sympathetic to People with Mental Illness*, by State and Territory—2007 BRFSS.

State	Unweighted Sample Size	Disagree Strongly	Disagree Slightly	Neither Agree Nor Disagree	Agree Slightly	Agree Strongly
Alaska	2,227	10.0%	26.4%	2.4%	38.7%	22.5%
Arkansas	5,083	12.9%	28.1%	1.4%	34.8%	22.8%
Colorado	5,234	9.1%	28.7%	1.2%	37.5%	23.5%
Connecticut	6,393	11.7%	28.0%	4.0%	34.0%	22.4%
District of Columbia	3,308	13.3%	28.8%	3.1%	33.3%	21.5%
Illinois	4,940	10.1%	26.6%	1.1%	38.6%	23.5%
Indiana	5,262	10.4%	25.5%	2.0%	38.1%	23.9%
Iowa	4,779	10.9%	24.9%	1.5%	40.0%	22.7%
Kentucky	5,899	16.4%	19.3%	10.6%	28.5%	25.2%
Louisiana	5,712	13.5%	20.1%	3.7%	27.8%	34.8%
Maine	3,734	9.2%	26.7%	2.1%	38.1%	23.9%
Minnesota	4,669	8.3%	23.8%	5.1%	40.8%	21.9%
Montana	5,247	9.4%	26.2%	4.8%	39.8%	19.8%
New Hampshire	5,296	10.6%	26.1%	3.6%	38.2%	21.5%
New Mexico	5,723	11.4%	28.8%	2.6%	33.4%	23.7%
Ohio	4,866	10.8%	27.7%	3.3%	38.0%	20.1%
Oklahoma	6,558	11.7%	25.1%	1.2%	34.7%	27.3%
Oregon	1,822	16.1%	31.4%	2.1%	32.8%	17.7%
Puerto Rico	3,684	6.6%	40.8%	17.3%	30.9%	4.4%
Rhode Island	3,823	9.7%	22.4%	4.4%	38.6%	24.9%
Texas	7,032	11.2%	21.9%	5.8%	35.9%	25.1%
Virginia	5,130	11.2%	27.3%	4.2%	35.9%	21.5%
Wisconsin	4,266	9.7%	28.0%	0.5%	39.5%	22.3%

*Adjusted for sex, age group, racial/ethnic group, education and household income level. Estimates are weighted; sample size is unweighted.

Table 4. Level of Agreement* with the Statement, *People are Generally Caring and Sympathetic to People with Mental Illness*, by State and Territory—2009 BRFSS.


State	Unweighted Sample Size	Disagree Strongly	Disagree Slightly	Neither Agree Nor Disagree	Agree Slightly	Agree Strongly
California	3,621	9.8%	25.3%	0.8%	43.6%	20.5%
Georgia	5,122	13.5%	23.5%	4.7%	33.7%	24.6%
Hawaii	6,010	8.5%	24.1%	2.5%	39.7%	25.2%
Kansas	8,414	9.2%	24.3%	6.1%	38.2%	22.2%
Massachusetts	4,535	10.0%	24.1%	4.3%	36.4%	25.1%
Michigan	2,736	10.9%	27.7%	1.4%	36.9%	23.0%
Mississippi	9,964	17.8%	25.2%	2.5%	29.0%	25.5%
Missouri	4,522	11.7%	28.5%	2.9%	35.3%	21.6%
Nebraska	4,745	7.6%	23.7%	2.2%	41.4%	25.1%
Nevada	3,420	12.2%	28.9%	2.3%	34.6%	22.0%
South Carolina	8,775	10.4%	26.4%	1.1%	37.2%	24.9%
Tennessee	2,100	7.7%	21.4%	7.1%	27.9%	35.9%
Utah	2,366	7.0%	26.2%	0.9%	42.7%	23.2%
Vermont	6,094	8.3%	25.5%	2.7%	38.9%	24.5%
Washington	6,853	16.9%	28.3%	2.3%	33.7%	18.9%
Wyoming	5,476	9.2%	26.5%	1.6%	39.3%	23.5%

*Adjusted for sex, age group, racial/ethnic group, education and household income level. Estimates are weighted; sample size is unweighted.

ALASKA, 2007


On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than adults 18-24 years old and more women than men disagreed that people are caring and sympathetic to people with mental illness (Figure 2). About 59% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 47% of adults with frequent mental distress and 51% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, or Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 27% of adults with diabetes **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 17% of adults with cardiovascular disease **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The American Diabetes Association offers tips for coping with stress for individuals living with diabetes. <http://www.diabetes.org/living-with-diabetes/complications/stress.html>
- ▶ The American Heart Association offers tips for coping with feelings associated with cardiovascular disease. http://www.heart.org/HEARTORG/Conditions/More/CardiacRehab/Coping-with-Feelings_UCM_307092_Article.jsp

See Appendix B for supporting data for highlighted groups

ARKANSAS, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 41% disagreed that people are caring and sympathetic to people with mental illness. More younger adults between 18 and 24 years old than adults 55 years old or older and more men than women disagreed that treatment helps (Figure 1). More women than men; and more white, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 63% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 54% of adults with frequent mental distress and just over 50% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 8% of adults with less than a high school education **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- About 17% of adults living in households earning less than \$20,000 annually **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>

See Appendix B for supporting data for highlighted groups

CALIFORNIA, 2009

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 35% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more black, non-Hispanic and Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 25 and 34 years old than adults between 35 and 54 years old and more white, non-Hispanic adults than other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 51% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 40% of adults with frequent mental distress and 43% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 8% of black, non-Hispanic adults strongly disagreed that treatment can help people with mental illness lead normal lives.
- About 14% of adults with arthritis strongly disagreed that people are caring and sympathetic to people with mental illness.
- ▶ **Stories that Heal** aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov
- ▶ The **Arthritis Foundation** offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

COLORADO, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 38% disagreed that people are caring and sympathetic to people with mental illness. More adults between 18 and 24 years old than adults 35-54 years old and more men than women disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults; more women than men; and more white, non-Hispanic and black, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 61% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 51% of adults with frequent mental distress and 46% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 15% of adults with cardiovascular disease **strongly disagreed** that people are caring and sympathetic to people with mental illness.
 - About 24% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The American Heart Association offers tips for coping with feelings associated with cardiovascular disease. http://www.heart.org/HEARTORG/Conditions/More/CardiacRehab/Coping-with-Feelings_UCM_307092_Article.jsp
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

CONNECTICUT, 2007

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 40% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults; more women than men; and more white, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 59% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 54% of adults with frequent mental distress and 49% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 18% of adults with asthma **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 20% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The American Lung Association offers programs for individuals living with asthma to help manage symptoms of the disease, to manage stress, and to facilitate relaxation. <http://www.lungusa.org/>
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

DISTRICT OF COLUMBIA, 2007

On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 42% disagreed that people are caring and sympathetic to people with mental illness. More black, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults aged 25–54 years old than older adults; more women than men; and more white, non-Hispanic and black, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 49% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic, as did 49% of adults with frequent mental distress and 51% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 25% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 25% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

GEORGIA, 2009

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 37% disagreed that people are caring and sympathetic to people with mental illness. More men than women; and more Hispanic adults than adults in any other racial/ethnic groups, and more black, non-Hispanics than white, non-Hispanics disagreed that treatment helps (Figure 1). More adults 25–34 years old than adults 55 years old or older and more women than men disagreed that people are caring and sympathetic (Figure 2). About 59% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 47% of adults with frequent mental distress and 52% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 17% of Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
 - ▶ The SAMHSA ADS Center provides resources for implementing culturally sensitive anti-stigma activities. <http://promoteacceptance.samhsa.gov/topic/culture/brochures.aspx>
 - ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements (PSAs) in Spanish. <http://www.aceptariignorar.samhsa.gov/>

See Appendix B for supporting data for highlighted groups

HAWAII, 2009


On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 33% disagreed that people are caring and sympathetic to people with mental illness. More Hispanic and other, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults; more women than men; and more white, non-Hispanic adults than other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 49% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 45% of adults with frequent mental distress and 48% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 5% of Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- ▶ The SAMHSA ADS Center provides resources for implementing culturally sensitive anti-stigma activities. <http://promoteacceptance.samhsa.gov/topic/culture/brochures.aspx>
- ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements (PSAs) in Spanish. <http://www.aceptarignorar.samhsa.gov/>

See Appendix B for supporting data for highlighted groups

ILLINOIS, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 37% disagreed that people are caring and sympathetic to people with mental illness. Adults in different age groups, both sexes, and different racial/ethnic groups disagreed to the same extent that treatment helps (Figure 1). More women than men and more white, non-Hispanic and black, non-Hispanic adults than adults in other racial/ethnic groups disagreed that people are caring and sympathetic (Figure 2). About 67% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 50% of adults with frequent mental distress and 46% of those receiving treatment for a mental health or emotional problem.

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 17% of adults who are unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 31% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress. <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

INDIANA, 2007

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than adults 18-24 years old and 55 years old or older; more women than men; and more black, non-Hispanic adults than white, non-Hispanic and Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 59% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 48% of adults with frequent mental distress and 45% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 5% of adults with less than a high school education **strongly disagreed** that treatment can help people with mental illness lead normal lives.
 - About 17% of non-Hispanic black adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov

See Appendix B for supporting data for highlighted groups

IOWA, 2007

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 45% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 49% of adults with frequent mental distress and 47% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 20% of adults who were out of work **strongly disagreed** that treatment can help people with mental illness lead normal lives.
 - About 16% of adults living with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ **Getting Through Tough Economic Times** provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

KANSAS, 2009

On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 34% disagreed that people are caring and sympathetic to people with mental illness. More adults 55 years old or older than adults 35 to 54 years old and more men than women disagreed that treatment helps (Figure 1). More adults between 25 and 34 years old than younger adults or adults 55 years old or older; more women than men; and more white, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 56% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 46% of adults with frequent mental distress and 43% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 14% of adults living in households earning <\$20,000 a year **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 12% of adults with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

KENTUCKY, 2007

On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More other, non-Hispanic adults than adults in any other racial/ethnic groups disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 55% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 49% of adults with frequent mental distress and 50% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 21% of women **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 31% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Womenshealth.gov provides resources and information to women about mental health and mental illnesses.
<http://www.womenshealth.gov/mental-health/index.cfm>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress.
<http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtm>

See Appendix B for supporting data for highlighted groups

LOUISIANA, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 34% disagreed that people are caring and sympathetic to people with mental illness. More younger adults between 18 and 34 years old than older adults; more men than women; and more black, non-Hispanic than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 25 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 36% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 41% of adults with frequent mental distress and 37% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 23% of adults with frequent mental distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 21% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress:
<http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

MAINE, 2007


On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more other, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 64% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 53% of adults with frequent mental distress and 48% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics and Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics and Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 16% of adults living in households earning <\$20,000 a year **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 29% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

MASSACHUSETTS, 2009

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 34% disagreed that people are caring and sympathetic to people with mental illness. More Hispanic adults than adults in any other racial/ethnic group disagreed that treatment helps (Figure 1). More women than men and more white, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 46% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 52% of adults with frequent mental distress and 42% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 6% of Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- About 3% of adults earning <\$20,000 a year **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements in Spanish. <http://www.aceptarignorar.samhsa.gov/>
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>

See Appendix B for supporting data for highlighted groups

MICHIGAN, 2009


On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 39% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more black, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 48% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic, as did 46% of adults with frequent mental distress and 44% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 6% of black, non-Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- About 19% of black, non-Hispanic adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community www.storiesthatheal.samhsa.gov
- ▶ The SAMHSA ADS Center provides guidance and toolkits for culturally sensitive anti-stigma activities. <http://promoteacceptance.samhsa.gov/topic/culture/brochures.aspx>

See Appendix B for supporting data for highlighted groups

MINNESOTA, 2007

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 32% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 63% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 53% of adults with frequent mental distress and 45% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 13% of adults with cardiovascular disease **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 11% of adults with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The American Heart Association offers resources for managing depression and anxiety in people with cardiovascular disease. <http://www.heart.org/HEARTORG/>
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

MISSISSIPPI, 2009

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 43% disagreed that people are caring and sympathetic to people with mental illness. More adults between 25 and 54 years old than older adults; more men than women; and more black, non-Hispanic than white and other non-Hispanic adults disagreed that treatment helps (Figure 1). More younger adults between 18 and 54 years old than older adults; more women than men; and more other, non-Hispanic adults than adults in other racial/ethnic groups disagreed that people are caring and sympathetic (Figure 2). About 56% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic, as did 56% of adults with frequent mental distress and 58% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 8% of black, non-Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
 - About 54% of other, non-Hispanic adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community www.storiesthatheal.samhsa.gov
- ▶ The SAMHSA ADS Center provides guidance and toolkits for culturally sensitive anti-stigma activities. <http://promoteacceptance.samhsa.gov/topic/culture/brochures.aspx>

See Appendix B for supporting data for highlighted groups

MISSOURI, 2009

On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 40% disagreed that people are caring and sympathetic to people with mental illness. More black, non-Hispanic adults than adults in any other racial/ethnic groups disagreed that treatment helps (Figure 1). More adults between 25 and 54 years old than older adults; more women than men; and more other, non-Hispanic adults than adults in other racial/ethnic groups disagreed that people are caring and sympathetic (Figure 2). About 60% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 57% of adults with frequent mental distress and 53% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 13% of black, non-Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
 - About 18% of adults living in households earning <\$20,000 a year **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>

See Appendix B for supporting data for highlighted groups

MONTANA, 2007


On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more other, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 58% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 48% of adults with frequent mental distress and 44% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 12% of women **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 25% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Womenshealth.gov provides resources and information to women about mental health and mental illnesses. <http://www.womenshealth.gov/mental-health/index.cfm>
- ▶ Getting Through Tough Economic Times™ provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

NEBRASKA, 2009

On average, about 7% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 31% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more other, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 47% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to mental illness, as did 46% of adults with frequent mental distress and 45% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 24% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 18% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

NEVADA, 2009

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 41% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more Hispanic adults than white, non-Hispanic and black, non-Hispanic adults disagreed that treatment helps (Figure 1). More women than men and more white, non-Hispanic adults than Hispanic adults or other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 64% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 47% of adults with frequent mental distress and 65% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 11% of Hispanic adults **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- About 28% of black, non-Hispanic adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements in Spanish. <http://www.aceptarignorar.samhsa.gov/>
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov

See Appendix B for supporting data for highlighted groups

NEW HAMPSHIRE, 2007


On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 37% disagreed that people are caring and sympathetic to people with mental illness. Adults in different age groups, both sexes, and different racial/ethnic groups disagreed to the same extent that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 64% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 54% of adults with frequent mental distress and 46% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 25% of Hispanic adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 21% of out-of-work adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.

▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements in Spanish.

<http://www.acceptarignorar.samhsa.gov/>

▶ *Getting Through Tough Economic Times* provides guidance on coping with financial difficulties. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

NEW MEXICO, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 40% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More women than men and more white, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 56% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 50% of adults with frequent mental distress and 54% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 14% of women **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 15% of adults with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Womenshealth.gov provides resources and information to women about mental health and mental illnesses. <http://www.womenshealth.gov/mental-health/index.cfm>
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

OHIO, 2007

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 39% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than adults 18-24 years old; more women than men; and more white, non-Hispanic adults and black, non-Hispanic adults than other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 64% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 52% of adults with frequent mental distress and 50% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 18% of black, non-Hispanic adults **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 16% of adults living in households earning <\$20,000 a year **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>

See Appendix B for supporting data for highlighted groups

OKLAHOMA, 2007

On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 37% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more black, non-Hispanic and Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 25 and 54 years old than adults in other age groups; more women than men; and more white, non-Hispanic adults and other, non-Hispanic adults than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 52% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 47% of adults with frequent mental distress and 52% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 14% of women **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 16% of adults with asthma **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Womenshealth.gov provides resources and information to women about mental health and mental illness.
- ▶ The American Lung Association offers programs for individuals living with asthma to help manage symptoms of the disease, to manage stress, and to facilitate relaxation. <http://www.lungusa.org/>

See Appendix B for supporting data for highlighted groups

OREGON, 2007


On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 48% disagreed that people are caring and sympathetic to people with mental illness. Adults in different age groups, both sexes, and different racial/ethnic groups disagreed to the same extent that treatment helps (Figure 1). More adults between 25 and 34 years old than adults 55 years old or older disagreed that people are caring and sympathetic (Figure 2). About 61% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 62% of adults with frequent mental distress and 51% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics and Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 32% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 41% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

PUERTO RICO, 2007


On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 47% disagreed that people are caring and sympathetic to people with mental illness. Adults in different age groups and both sexes disagreed to the same extent that treatment helps (Figure 1). More adults between 25 and 54 years old than older adults disagreed that people are caring and sympathetic (Figure 2). About 57% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 62% of adults with frequent mental distress, and 49% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for White, non-Hispanics, Black, non-Hispanics, and Other, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for White, non-Hispanics, Black, non-Hispanics, and Other, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 16% of adults with frequent mental distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements in Spanish. <http://www.aceptariqnorar.samhsa.gov/>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtm>

See Appendix B for supporting data for highlighted groups

RHODE ISLAND, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 32% disagreed that people are caring and sympathetic to people with mental illness. More adults between 25 and 34 years old than adults 55 years old or older; more men than women; and more Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 58% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 44% of adults with frequent mental distress, and 39% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 16% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 13% of adults with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups

SOUTH CAROLINA, 2009

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 37% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more black, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 58% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 51% of adults with frequent mental distress and 52% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 16% of adults with cardiovascular disease **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 24% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.

▶ The American Heart Association offers resources for managing depression and anxiety in people with cardiovascular disease. <http://www.heart.org/HEARTORG>


▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

TENNESSEE, 2009


On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 29% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 50% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 46% of adults with frequent mental distress and 38% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Hispanics and Other, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Hispanics and Other, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 19% of adults with frequent mental distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 22% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress:
<http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.html>

See Appendix B for supporting data for highlighted groups

TEXAS, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 33% disagreed that people are caring and sympathetic to people with mental illness. More non-Hispanic black, and Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults; more women than men; and more white, non-Hispanic adults than Hispanic adults and other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 52% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 46% of adults with frequent mental distress and 46% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 15% of adults with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 21% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

UTAH, 2009


On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 33% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 52% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 44% of adults with frequent mental distress and 46% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanic were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanic were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 11% of adults living with arthritis **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 20% of adults with serious psychological distress **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

VERMONT, 2009


On average, about 4% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 34% disagreed that people are caring and sympathetic to people with mental illness. Adults in different age groups, both sexes, and different racial/ethnic groups disagreed to the same extent that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults and more women than men disagreed that people are caring and sympathetic (Figure 2). About 62% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 53% of adults with frequent mental distress and 43% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 14% of adults out of work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- About 13% of adults with cardiovascular disease **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>
- ▶ The American Heart Association offers resources for managing depression and anxiety in people with cardiovascular disease. <http://www.heart.org/HEARTORG>

See Appendix B for supporting data for highlighted groups

VIRGINIA, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 39% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more other, non-Hispanic adults than white, non-Hispanic and black, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults in other age groups; more women than men; and more white, non-Hispanic adults and black, non-Hispanic adults than other, non-Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 52% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 49% of adults with frequent mental distress and 44% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 19% of adults with frequent mental distress and 19% of adults with serious psychological distress **strongly disagreed** people are caring and sympathetic to people with mental illness
- ▶ The National Institute of Mental Health provides tips on how to recognize and deal with stress: <http://www.nimh.nih.gov/health/publications/stress/fact-sheet-on-stress.shtml>

See Appendix B for supporting data for highlighted groups

WASHINGTON, 2009

On average, about 5% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 45% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more Hispanic than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than adults 18-24 years old or adults 55 years old or older; more women than men; and more white, non-Hispanic than Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 60% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 56% of adults with frequent mental distress and 52% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 11% of Hispanic adults in Washington **strongly disagreed** that treatment can help people with mental illness lead normal lives.
- About 37% of black, non-Hispanic adults in Washington **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ The Campaign for Social Inclusion uses proven social marketing strategies and public education methods to disseminate TV, radio, and print public service advertisements in Spanish. <http://www.aceptariqnorar.samhsa.gov>
- ▶ Stories that Heal aims to promote acceptance of mental health problems within the African American community. www.storiesthatheal.samhsa.gov

See Appendix B for supporting data for highlighted groups

WISCONSIN, 2007

On average, about 6% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 38% disagreed that people are caring and sympathetic to people with mental illness. More men than women disagreed that treatment helps (Figure 1). More women than men disagreed that people are caring and sympathetic (Figure 2). About 54% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 53% of adults with frequent mental distress and 49% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness





Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION


- About 19% of adults with household incomes <\$20,000 **strongly disagreed** that people are caring and sympathetic to people with mental illness.
 - About 19% of adults unable to work **strongly disagreed** that people are caring and sympathetic to people with mental illness.
- ▶ Information regarding access to low-cost health care, including mental health treatment, is available at: <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=2&lvlID=18>
- ▶ *Getting Through Tough Economic Times* provides practical advice on how to deal with the effects of financial difficulties on physical and mental health. <http://www.samhsa.gov/economy>

See Appendix B for supporting data for highlighted groups

WYOMING, 2009


On average, about 7% of survey respondents disagreed that treatment can help people with mental illness lead normal lives, while 36% disagreed that people are caring and sympathetic to people with mental illness. More men than women and more Hispanic and other, non-Hispanic adults than white, non-Hispanic adults disagreed that treatment helps (Figure 1). More adults between 35 and 54 years old than older adults; more women than men; and more other, non-Hispanic adults than white, non-Hispanic adults and Hispanic adults disagreed that people are caring and sympathetic (Figure 2). About 60% of adults experiencing serious psychological distress disagreed that people are caring and sympathetic to people with mental illness, as did 51% of adults with frequent mental distress and 50% of those receiving treatment for a mental health or emotional problem (Figure 3).

Figure 1. Percent Total Disagreement That Treatment Can Help People With Mental Illness Lead Normal Lives




*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 2. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness



*Results for Black, non-Hispanics were suppressed due to small sample size

Figure 3. Percent Total Disagreement That People Are Caring and Sympathetic To People With Mental Illness by Serious Psychological Distress, Frequent Mental Distress, and Receiving Treatment for a Mental Health or Emotional Problem



CALLS TO ACTION

- About 20% of adults with cardiovascular disease strongly disagreed that people are caring and sympathetic to people with mental illness
- About 12% of adults with arthritis strongly disagreed that people are caring and sympathetic to people with mental illness.
- ▶ The American Heart Association offers tips for coping with feelings associated with cardiovascular disease. http://www.heart.org/HEARTORG/Conditions/More/CardiacRehab/Coping-with-Feelings_UCM_307092_Article.jsp
- ▶ The Arthritis Foundation offers programs to improve the quality of life for individuals with arthritis, including how to manage pain and get support for dealing with the emotional impact of living with arthritis. <http://www.arthritis.org/programs.php>

See Appendix B for supporting data for highlighted groups



Discussion

Most adults in the 37 states and territories surveyed in 2007 and 2009 believe that mental illness treatment is effective, consistent with recent findings (Croghan et al., 2003; Pescosolido et al., 2010), but substantially fewer adults believe that people are caring and sympathetic to people with mental illness.

Adults with mental illness symptoms appear to perceive a less friendly social environment for people with mental illness. Adults who lived in states that had more resources allocated for mental health services had more positive attitudes regarding the effectiveness of mental illness treatment: They were more willing to seek treatment and more likely to perceive that others were caring and sympathetic to people with mental illness. Young adults who lived in states that had more resources to market the What a Difference a Friend Makes campaign—designed to show how to support a friend with mental illness—were more likely to believe that other people are caring and sympathetic to people with mental disorders, suggesting that the campaign had a positive effect in these states⁶ and contributes to evidence for such campaigns (Corrigan, 2012).

The findings regarding favorable attitudes toward treatment effectiveness might result from the multiple efforts of previous mental health public awareness campaigns and widespread marketing of medicines for common mental disorders (Rosenthal et al., 2002; Frosch, Grand, Tarn, & Kravitz, 2010). Although smaller percentages of adults disagreed that treatment is effective, even small percentages at the population level (e.g., 5%) can represent thousands of people possibly at risk of not seeking treatment. The state-specific findings also highlight some subgroups who strongly disagreed with statements about treatment effectiveness (e.g., blacks, Hispanics, adults with less education). Individuals in these subgroups with mental illness symptoms might be at increased risk of a number of negative individual and social outcomes if they choose not to seek treatment because they do not believe treatment is effective. States can use these findings to target these population groups to improve their mental health literacy and to reduce negative attitudes toward mental illness. Despite belief in treatment effectiveness, some people remain unwilling to take psychiatric medications (Croghan, et al., 2003). Reasons for not taking medications might include concerns about unwanted side effects and preferences for nonpharmacologic treatments such as cognitive behavioral therapy (Schnittker 2003; Riedel-Heller, 2005). Public education about the effectiveness of different types of treatment, including alternatives to psychiatric medications and the chronic nature of untreated mental illness for some individuals, may be beneficial for some groups.

That substantial percentages of adults, particularly those with mental illness symptoms, and those with chronic conditions did not believe that other people are caring and sympathetic to people with mental illness is of concern. These individuals might not be getting the emotional support they need for their own conditions. Some people respond negatively when confronted with a friend or family member's mental illness, contributing to avoidance and isolation—potentially interfering with recovery. The emotional and psychological aspects of mental illness make supportive friends and family even more important to a person's recovery. By offering reassurance, companionship, emotional strength, and acceptance, a friend, family member, or co-worker can help a person with mental illness through his or her treatment and recovery.

⁶ A concurrent (2007–2009) campaign targeting college students was supported by the Jed Foundation.



Strategies for Combating Stigma

Attitudes toward mental illness collected through the BRFSS identify areas for further study and population groups for intervention to reduce negative attitudes toward mental illness and promote social inclusion of those with mental illness symptoms. The cumulative findings in this report offer federal and state decision makers and other key stakeholders' insights about the public's attitudes toward persons living with mental illness as well as the relative impact of anti-stigma efforts on public attitudes toward mental illness.

The current data indicate the following strategies be considered by mental illness stakeholders and the general public to improve attitudes and behaviors toward persons with mental illness and promote social inclusion.

- ▶ Continue to explore how to monitor attitudes toward mental illness and evaluate innovative anti-stigma interventions.
- ▶ Implement culturally competent stigma reduction initiatives at local, regional, and state-wide levels. Guidance on event planning, partnership development, outreach to schools and businesses, mental health resources, and marketing to the general public is provided in SAMHSA's Developing a Stigma Reduction Initiative. (See <http://store.samhsa.gov/shin/content/SMA06-4176/SMA06-4176.pdf>.)
- ▶ Implement evidence-based mental health literacy programs such as Mental Health First-Aid USA in states and local communities. (See http://www.mentalhealthfirstaid.org/cs/program_overview/.)
- ▶ Offer technical assistance to local media regarding how they can reduce stigma by avoiding sensationalism about mental illness and ensure balance in broadcasting by encouraging stories about recovery, accomplishment, and contributions by people with mental illness.
- ▶ Support people with mental health problems by helping to develop community resources and by referring them to available community resources.
- ▶ Don't label people by their illness. Instead of saying, "she's bipolar," say, "she has bipolar disorder."
- ▶ Learn how to offer reassurance, companionship, emotional strength, and acceptance to a friend, family member, neighbor, or others with mental health problems.

Limitations

The findings in this report are subject to several limitations. First, because BRFSS surveys in 2007 and 2009 included only noninstitutionalized adults with landline telephones, persons in institutions and in households without telephones (which might include those with more serious mental illness) or with cell phones only are excluded, perhaps reducing the differences observed among groups. Second, because states commonly use only English- or Spanish-language surveys, persons who speak other primary languages are excluded. Findings from Puerto Rico did not closely match the distribution of responses in other states surveyed. This may be due to survey language or cultural factors that influence attitudes and response styles. The relatively small number of participants from Puerto Rico is unlikely to skew overall findings related to trends or media impact. Third, factors other than the sociodemographic characteristics studied here (e.g., cultural, social, environmental) also might account for a portion of differences in attitudes. Fourth, this study examined attitudes as indicators of stigma, but could not assess experienced stigma (e.g., exclusion) or its consequences (e.g., social isolation, lost employment) or differentiate between felt and experienced stigma in some groups (e.g., unemployed, people receiving treatment). Fifth, participants' responses to a public survey may be influenced by social-desirability biases or social comparison effects whereby respondents respond more favorably or believe they are more open-minded than the public (Green,



further examination regarding its interpretation and associations with other attitudes (The Scottish Government, 2009). Other state-level factors (e.g., average unemployment rate) were not statistically significant predictors of attitudes in this study, but future studies using additional data from other states and other attitudinal items might find associations between these variables and attitudes toward mental illness.

Acknowledgment

This report is based, in part, on data contributed by BRFSS state coordinators and state mental health services data infrastructure coordinators.

RESOURCES FOR STATES TO ADDRESS MENTAL ILLNESS STIGMA

Federal Resources

- ▶ The SAMHSA Resource Center to Promote Acceptance, Dignity and Social (ADS) Inclusion Associated with Mental Health: <http://www.stopstigma.samhsa.gov/>.
- ▶ Substance Abuse and Mental Health Services Administration. (2006). *Developing a Stigma Reduction Initiative*. SAMHSA Pub No. SMA-4176. Rockville, MD: Center for Mental Health Services, U.S. Department of Health and Human Services. From <http://store.samhsa.gov/shin/content/SMA06-4176/SMA06-4176.pdf> (accessed July 9, 2012).
- ▶ SAMHSA: What a Difference a Friend Makes: <http://www.whatadifference.samhsa.gov/>.
- ▶ The National Institute on Mental Health (NIMH): <http://www.nimh.nih.gov/index.shtml>.
- ▶ Office on Women's Health: <http://www.womenshealth.gov/mental-health/>.
- ▶ U.S. Department of Veterans Affairs: <http://www.mentalhealth.va.gov/>.
- ▶ CDC Mental Health Work Group: <http://www.cdc.gov/mentalhealth/>.

Nonfederal Resources

- ▶ Mental Health America: <http://www.mentalhealthamerica.net/>.
- ▶ The Rosalynn Carter Fellowships For Mental Health Journalism: http://www.cartercenter.org/health/mental_health/fellowships/index.html.



REFERENCES

- Ajzen, I., & Fishbein, M. (1980). *Understanding attitudes and predicting social behavior*. Englewood Cliffs, New Jersey: Prentice Hall.
- Centers for Disease Control and Prevention. (2010). Attitudes toward Mental Illness—35 States, District of Columbia, and Puerto Rico, 2007. *Morbidity and Mortality Weekly Report*, May 28, 2010. 59(20); 619-625. Available from <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5920a3.htm>. (accessed July 9, 2012).
- Centers for Disease Control and Prevention. (2012). *Behavioral Risk Factor Surveillance System*. Atlanta, GA: Author. Available from <http://www.cdc.gov/BRFSS> (accessed July 9, 2012).
- Centers for Disease Control and Prevention. (2008). *2007 Behavioral Risk Factor Surveillance System summary data quality report*. Atlanta, GA: Author. Available from <ftp://ftp.cdc.gov/pub/data/brfss/2007SummaryDataQualityReport.pdf> (accessed July 9, 2012).
- Centers for Disease Control and Prevention. (2011). *2009 Behavioral Risk Factor Surveillance System Summary Data Quality Report*. Atlanta, GA: Author. Retrieved February 28, 2012 Available from ftp://ftp.cdc.gov/pub/data/brfss/2009_Summary_Data_Quality_Report.pdf (accessed July 9, 2012).
- Chapman, D.P., Perry, G.S., & Strine, T.W. (2005). The vital link between chronic disease and depressive disorders. *Preventing Chronic Disease*, 2(1). Available from http://www.cdc.gov/pcd/issues/2005/jan/04_0066.htm (accessed July 9, 2012).
- Colton, C.W., & Manderscheid, R.W. (2006). Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Preventing Chronic Disease*, 3(2). Available from http://www.cdc.gov/pcd/issues/2006/apr/05_0180.htm (accessed July 9, 2012).
- Corrigan, P.W. (2012). Where is the evidence supporting public service announcements to eliminate mental illness stigma? *Psychiatric Services*, 63(1), 79–82.
- Corrigan, P.W., Markowitz, F.E., & Watson, A.C. (2004). Structural levels of mental illness: Stigma and discrimination. *Schizophrenia Bulletin*, 20(3), 481–491.
- Croghan, T.W., Tomlin, M., Pescosolido, B.A., Schnittker, J., Martin, J.K., Lubell, K. et al. (2003). American attitudes toward and willingness to use psychiatric medications. *The Journal of Nervous and Mental Disease*, 191(3), 166–174.
- El-Gabalawy, R., Cox, B., Clara, I., & Mackenzie, C. (2010). Assessing the validity of social anxiety disorder subtypes using a nationally representative sample. *Journal of Anxiety Disorders*, 24(2), 244-249.
- Frosch, D.L., Grande, D., Tarn, D.M., & Kravitz, R.L. (2010). A decade of controversy: Balancing policy with evidence in the regulation of prescription drug advertising. *American Journal of Public Health*, 100(1), 24–32.
- Goffman, E. (1963). *Stigma: Notes on the management of spoiled identity*. New York: Simon & Schuster, Inc.
- Green, G. (1995). Attitudes toward people with HIV: Are they as stigmatizing as people with HIV perceive them to be? *Social Science & Medicine*, 41(4), 557–568.
- Greene-Shortridge, T.M., Britt, T.W., & Castro, C.A. (2007). The stigma of mental health problems in the military. *Military Medicine*, 172(2), 157–161.



Hennessy, K.D., Green-Hennessy, S., Marshall, C. (2012). Datapoints: Receipt of reported mental health specialty care in the 2010 HealthStyles Survey. *Psychiatric Services*, 63(4), 306.

Institute of Medicine (IOM). (2012). *Living well with chronic illness: A call for public health action*. Washington, DC: The National Academies Press.

Jorm, A.F., Korten, A.E., Jacomb, P.A., Christensen, H., Rodgers, B., & Pollitt, P. (1997). “Mental health literacy”: A survey of the public’s ability to recognize mental disorders and their beliefs about the effectiveness of treatment. *Medical Journal of Australia*, 166(4), 182–186.

Kadri, N. & Sartorius, N. (2005). The global fight against the stigma of schizophrenia. *PLoS Medicine*, 2(7), 597–599.

Kessler, R.C., Barker, P.R., Colpe, L.J., Epstein, J.F., Gfroerer, J.C., Hiripi, E., Howes, M.J., Norman, S.T., Manderscheid, R.W., Walters, E.E., & Zaslavsky, A.M. (2003). Screening for serious mental illness in the general population. *Archives of General Psychiatry*, 60(2), 184–189.

Kessler, R.C., Chiu, W.T., Demler, O., & Walters, E.E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62(6), 617–627.

Kobau, R., DiIorio, C., Chapman, D., & Delvecchio, P. (2010). Substance Abuse and Mental Health Services Administration/CDC Mental Illness Stigma Panel Members. Attitudes about mental illness and its treatment: validation of a generic scale for public health surveillance of mental illness associated stigma. *Community Mental Health Journal*, 46(2), 164–176.

Korn, E.L. & Graubard, B.I. (1999). Predictive margins for survey data. *Biometrics*, 55(2), 652–659.

Link, B.G. & Phelan, J.C. (2001). Conceptualizing stigma. *Annual Review of Sociology*, 27, 363–385.

Link, B.G., Yang, L.H., Phelan, J.C., & Collins, P.Y. (2004). Measuring mental illness stigma. *Schizophrenia Bulletin*, 30(3), 512–541.

Link, B.G. & Phelan, J.C. (2006). Stigma and its public health implications. *The Lancet*, 367(9509), 528–529.

Muthén, L.K. & Muthén, B.O. (1998–2011). *Mplus User’s Guide. Sixth Edition*. Los Angeles, CA: Authors.

Nadeem, E., Lange, J.M., Edge, D., Fongwa, M., Belin, T., & Miranda, J. (2007). Does stigma keep poor young immigrant and U.S. born black and Latina women from seeking mental health care? *Psychiatric Services*, 58(12), 1547–1554.

Pescosolido, B.A., Martin, J.K., Long, J.S., Medina, T.R., Phelan, J.C., & Link, B.G. (2010). “A disease like any other?” A decade of change in public reactions to schizophrenia, depression, and alcohol dependence. *American Journal of Psychiatry*, 16(11), 1321–1330.

Pfeffermann, D. & Sverchkov, M. (2009). Inference under informative sampling. In *Sample Surveys: Inference and Analysis. Handbook of Statistics 29B* (D. Pfeffermann and C. R. Rao, eds.) 455–487. North-Holland: Amsterdam.

President’s New Freedom Commission on Mental Health. (2003). *Achieving the Promise: Transforming Mental Health Care in America*. Rockville, MD: Author. Available from <http://govinfo.library.unt.edu/mentalhealthcommission/reports/FinalReport/downloads/FinalReport.pdf> (accessed July 9, 2012).



- Riedel-Heller, S.G., Matschinger, H., & Angermeyer, M.C. (2005). Mental disorders—who and what might help? Help-seeking and treatment preferences of the lay public. *Social Psychiatry and Psychiatric Epidemiology*, 40(2), 167–174.
- Research Triangle Institute. (2008). *SUDAAN language manual, release 10.0*. Research Triangle Park, NC: Author.
- Rosenthal, M.B., Berndt, E.R., Frank, R.G., Donohue, J.M., & Epstein, A.M. (2002). Promotion of prescription drugs to consumers. *New England Journal of Medicine*, 346(7), 498–505.
- The Scottish Government. (2009). *Well what do you think? The fourth National Scottish Survey of public attitudes to mental wellbeing and mental health problems*. Scotland, UK: Author.
- Sayce, L. (1998). Stigma, discrimination and social exclusion: What's in a word? *Journal of Mental Health*, 7(4), 331–343.
- Sartorius, N. (2005). Stigma and discrimination because of schizophrenia: A summary of the WPA Global Programme Against Stigma and Discrimination because of Schizophrenia. *World Psychiatry*, 4(S1), 11–15.
- Schnittker, J. (2003). Misgivings of medicine?: African-Americans' skepticism of psychiatric medication. *Journal of Health and Social Behavior*, 44(4), 506–523.
- Sirey, J., Bruce, M.L., Alexopoulos, G.S., Perlick, D.A., Raue, P., Friedman, S.J., & Meyers, B.S. (2001). Perceived stigma as a predictor of treatment discontinuation in young and older outpatients with depression. *American Journal of Psychiatry*, 158(3), 479–481.
- Sirey, J.A., Bruce, M.L., Alexopoulos, G.S., Perlick, D.A., Friedman, S.J., & Meyers, B.S. (2001). Stigma as a barrier to recovery: Perceived stigma and patient-rated severity of illness as predictors of antidepressant drug adherence. *Psychiatric Services*, 53(9), 1179.
- Smith, D.M., Damschroder, L.J., Kim, S.Y., & Ubel, P.A. (2012). What's it worth? Public willingness to pay to avoid mental illnesses compared with general illnesses. *Psychiatric Services*, 63(4), 319–324.
- Stuart, H. (2005). Why stigma matters and why it should be beaten. *World Psychiatry*, 4 (Supplement 1), 4–5.
- Substance Abuse and Mental Health Services Administration. (2006). *Developing a stigma reduction initiative*. SAMHSA Pub No. SMA-4176. Rockville, MD: U.S. Department of Health and Human Services.
- U.S. Department of Health and Human Services. (1999). *Mental health: A report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health.
- U.S. Department of Health and Human Services. (2001). *Mental health: Culture, race, and ethnicity—A supplement to mental health: A report of the Surgeon General*. Rockville, MD: Author.
- Wahl, O.F. (2003). News media portrayal of mental illness. *American Behavioral Scientist*, 46, 1594–1600.
- Wang, P.S., Lane, M., Olfson, M., Pincus, H.A., Wells, K.B., & Kessler, R.C. (2005). Twelve-month use of mental health services in the United States: Results from the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62(6), 629–640.
- Weiss, M.G., Ramakrishna, J., & Somma, D. (2006). Health-related stigma: Rethinking concepts and interventions. *Psychological Health Medicine*, 11(3), 277–287.
- Zahran, H.S., Kobau, R., Moriarty, D.G., Zack, M.M., Giles, W.H., & Lando, J. (2004). Self-reported frequent mental distress among adults—United States, 1993–2001. *Morbidity and Mortality Weekly Report*, 54(41), 963–966.

APPENDIX A:

Mental Health-related and BRFSS Indicators



Table A. Mental Health-related and BRFSS Indicators for 37 States and the District of Columbia*

	Per Capita Expenditure of State Mental Health Agencies (2004)		Population in Mental Health Professional Shortage Area (2010)		Population with Health Insurance Coverage		Unemployment Rate		Per Capita Donated Media Value		Frequent Mental Distress		Serious Psychological Distress		Receiving Mental Health Medicine or Treatment		Population Aged 18–24	
	Dollars		Percent		Percent		Percent		Dollars		Percent		Percent		Percent		Percent	
State	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009
Alaska	287.56	287.56	38	38	82	82	6	8	36.89	36.86	9	N/A	2	N/A	10	N/A	15	N/A
Arkansas	32.95	32.95	48	48	84	81	5	7	0.83	2.89	11	N/A	5	N/A	12	N/A	13	N/A
California	113.61	113.61	10	10	82	80	5	11	4.95	8.08	10	12	4	3	8	6	14	14
Colorado	69.07	69.07	15	15	84	85	4	8	1.32	16.10	9	N/A	3	N/A	10	N/A	13	N/A
Connecticut	152.09	152.09	13	13	91	88	5	8	4.75	6.43	8	N/A	3	N/A	12	N/A	9	N/A
District of Columbia	409.92	409.92	14	14	91	88	5	10	0.09	96.89	8	N/A	2	N/A	11	N/A	6	N/A
Georgia	51.25	51.25	32	32	83	80	5	10	1.26	1.42	11	9	5	3	11	8	8	8
Hawaii	147.99	147.99	6	6	93	92	3	7	0.01	0.20	9	7	2	2	6	6	13	12
Illinois	68.51	68.51	35	35	87	85	5	10	3.12	6.37	8	N/A	3	N/A	9	N/A	11	N/A
Indiana	80.58	80.58	16	16	89	86	5	10	0.67	2.06	10	N/A	3	N/A	12	N/A	13	N/A
Iowa	76.14	76.14	44	44	91	89	4	6	4.16	10.54	7	N/A	2	N/A	12	N/A	13	N/A
Kansas	83.22	83.22	37	37	87	87	4	7	2.99	2.94	7	9	3	2	10	11	8	9
Kentucky	49.69	49.69	28	28	86	84	6	11	3.82	4.40	12	N/A	6	N/A	14	N/A	9	N/A
Louisiana	52.63	52.63	59	59	82	84	4	7	1.09	2.09	9	N/A	5	N/A	11	N/A	8	N/A
Maine	130.66	130.66	12	12	91	90	5	8	9.19	18.40	9	N/A	4	N/A	15	N/A	12	N/A
Massachusetts	103.99	103.99	6	6	95	96	4	8	0.07	1.21	9	8	3	3	14	12	9	7
Michigan	90.96	90.96	12	12	88	86	7	14	2.18	6.81	10	11	4	3	11	14	11	11
Minnesota	121.37	121.37	27	27	92	91	5	8	0.50	0.71	7	N/A	3	N/A	11	N/A	13	N/A
Mississippi	95.50	95.50	35	35	81	82	6	10	3.48	4.48	13	13	6	7	12	13	14	12
Missouri	69.33	69.33	38	38	87	85	5	9	4.60	8.91	11	12	5	5	12	14	12	9
Montana	126.49	126.49	46	46	84	85	4	6	26.16	41.18	9	N/A	3	N/A	13	N/A	12	N/A
Nebraska	58.28	58.28	43	43	87	89	3	5	1.27	5.69	8	7	2	3	10	9	10	6
Nevada	54.45	54.45	8	8	83	79	5	12	2.19	2.74	11	12	4	4	8	8	12	8
New Hampshire	117.21	117.21	4	4	90	90	4	6	23.12	37.88	10	N/A	3	N/A	13	N/A	8	N/A
New Mexico	27.78	27.78	47	47	78	78	4	7	6.82	13.06	10	N/A	4	N/A	10	N/A	14	N/A

	Per Capita Expenditure of State Mental Health Agencies (2004)		Population in Mental Health Professional Shortage Area (2010)		Population with Health Insurance Coverage		Unemployment Rate		Per Capita Donated Media Value		Frequent Mental Distress		Serious Psychological Distress		Receiving Mental Health Medicine or Treatment		Population Aged 18–24	
	Dollars		Percent		Percent		Percent		Dollars		Percent		Percent		Percent		Percent	
	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009
State	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009	2007	2009
Ohio	64.06	64.06	19	19	88	86	6	10	3.37	3.82	12	N/A	5	N/A	13	N/A	8	N/A
Oklahoma	39.79	39.79	36	36	82	82	4	6	0.06	1.00	12	N/A	5	N/A	12	N/A	11	N/A
Oregon	60.79	60.79	24	24	83	82	5	11	3.90	6.82	10	N/A	3	N/A	13	N/A	9	N/A
Rhode Island	92.92	92.92	15	15	89	88	5	11	0.00	2.13	10	N/A	5	N/A	14	N/A	12	N/A
South Carolina	67.31	67.31	27	27	84	83	6	12	0.19	1.78	11	11	5	4	11	12	9	10
Tennessee	88.16	88.16	34	34	86	85	5	11	1.19	1.42	N/A	13	N/A	8	N/A	14	N/A	7
Texas	36.70	36.70	27	27	75	74	4	8	1.42	4.15	11	N/A	5	N/A	11	N/A	7	N/A
Utah	73.56	73.56	37	37	87	85	3	7	4.42	5.19	N/A	9	N/A	2	N/A	13	N/A	13
Vermont	165.95	165.95	7	7	89	90	4	7	13.31	46.83	9	9	3	3	13	14	13	12
Virginia	69.79	69.79	17	17	85	87	3	7	3.53	5.35	8	N/A	3	N/A	11	N/A	9	N/A
Washington	93.60	93.60	30	30	89	87	5	9	1.55	3.11	9	9	3	3	12	13	13	8
Wisconsin	94.82	94.82	36	36	92	91	5	9	3.68	5.09	8	N/A	3	N/A	10	N/A	14	N/A
Wyoming	102.44	102.44	75	75	86	84	3	6	7.45	7.17	9	9	3	3	11	12	14	13

Figure 1. Percent Disagreement With Treatment Can Help People With Mental Illness Lead Normal Lives




Figure 2. Percent Disagreement With People Are Generally Caring and Sympathetic To People With Mental Illness





Figure 3. Percent of Adults Receiving Mental Health Treatment



APPENDIX B:

State-specific Data Tables



Alaska, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,224	66.5	(63.2-69.6)	26.6	(23.6-29.7)	1.5	(0.6-3.3)*	4.0	(2.7-5.9)	1.5	(0.9-2.5)
Age (years)	18-24	148	53.7	(41.6-65.4)	41.8	(30.5-54.0)	3.1	(0.5-18.6)*	1.1	(0.4-3.0)*	0.3	(0.1-1.3)*
	25-34	366	64.2	(55.8-71.8)	29.0	(22.1-37.1)	0.6	(0.2-2.1)*	5.2	(2.2-11.7)*	1.0	(0.3-2.8)*
	35-54	1,029	63.7	(59.1-68.1)	26.7	(22.8-31.1)	1.6	(0.8-3.0)*	6.4	(3.8-10.7)	1.6	(0.8-3.2)*
	55+	662	78.6	(74.0-82.6)	15.8	(12.4-20.0)	0.7	(0.3-1.4)*	1.9	(1.1-3.4)	3.0	(1.3-6.6)*
Sex	Men	975	59.6	(54.2-64.7)	30.4	(25.8-35.3)	1.8	(0.5-6.2)*	6.0	(3.7-9.8)	2.2	(1.2-4.0)*
	Women	1,249	73.2	(69.3-76.7)	23.1	(19.6-26.9)	1.1	(0.6-2.3)*	1.9	(1.2-2.9)	0.8	(0.3-1.7)*
Race/ Ethnicity	Hispanic or Latino	57	74.6	(60.5-84.8)	11.4	(5.4-22.6)*	2.8	(0.9-8.7)*	1.8	(0.4-7.4)*	9.4	(3.8-21.3)*
	Not Hispanic or Latino:	1,619	67.7	(64.0-71.3)	26.2	(23.0-29.8)	1.8	(0.7-4.5)*	3.5	(1.9-6.3)*	0.7	(0.2-1.9)*
	White											
	Black or African American	25	--	--	--	--	--	--	--	--	--	--
	All Other Races	500	62.9	(54.6-70.5)	28.7	(21.9-36.5)	0.4	(0.1-0.9)*	5.9	(3.9-8.9)	2.2	(1.2-4.0)*
Education	<High School	152	51.0	(37.2-64.6)	27.7	(16.5-42.5)	13.7	(4.6-34.2)*	4.8	(2.4-9.7)*	2.8	(1.1-7.2)*
	High School Graduate (or GED)	651	63.2	(56.8-69.2)	31.0	(25.5-37.2)	0.9	(0.4-2.3)*	3.8	(2.5-5.7)	1.0	(0.5-2.1)*
	Some College	651	65.4	(59.3-71.0)	27.2	(22.1-33.0)	1.5	(0.8-2.8)*	4.4	(1.8-10.4)*	1.5	(0.6-3.3)*
	College Graduate	767	72.8	(67.1-77.7)	21.7	(17.3-27.0)	0.5	(0.2-1.7)*	3.4	(1.4-8.1)*	1.6	(0.5-5.0)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Alaska, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	264	69.2	(56.5-79.6)	18.2	(12.1-26.3)	1.4	(0.4-5.2)*	7.8	(2.7-20.5)*	3.4	(1.6-7.2)*
	\$20,000-\$34,999	323	68.4	(59.4-76.2)	26.5	(19.2-35.4)	1.1	(0.4-2.6)*	3.3	(1.5-7.0)*	0.7	(0.2-2.8)*
	\$35,000-\$49,999	319	59.2	(50.8-67.2)	31.5	(23.6-40.7)	5.5	(1.3-20.3)*	3.3	(1.7-6.4)*	0.4	(0.1-2.0)*
	\$50,000-\$74,999	414	68.8	(61.2-75.4)	26.5	(20.3-33.8)	0.7	(0.2-3.0)*	3.3	(1.3-8.4)*	0.7	(0.2-2.1)*
	\$75,000 and more	693	70.0	(64.6-74.9)	25.9	(21.2-31.2)	0.3	(0.1-0.7)*	3.0	(1.2-7.4)*	0.8	(0.3-2.7)*
Employment Status	Employed	1,514	65.9	(62.1-69.5)	28.2	(24.8-32.0)	0.7	(0.4-1.4)*	3.9	(2.5-6.1)	1.2	(0.6-2.2)*
	Out of work	127	60.7	(41.5-77.0)	27.2	(14.9-44.4)	7.7	(1.5-31.0)*	2.7	(1.1-6.3)*	1.7	(0.5-5.9)*
	Homemaker	173	61.5	(50.3-71.6)	34.4	(24.2-46.3)	1.7	(0.4-6.3)*	1.2	(0.4-3.5)*	1.2	(0.2-7.6)*
	Student	51	80.5	(63.5-90.8)	17.7	(7.9-34.9)*	0.0	(0.0-0.0)	1.2	(0.2-7.8)*	0.6	(0.1-2.6)*
	Retired	248	71.8	(63.2-79.1)	20.3	(14.5-27.6)	0.9	(0.3-2.4)*	2.6	(1.1-6.3)*	4.4	(1.1-15.4)*
	Unable to work	101	68.9	(52.5-81.6)	13.2	(7.1-23.2)*	1.8	(0.4-7.5)*	14.9	(4.6-39.0)*	1.2	(0.4-4.0)*
Veteran Status	Yes	373	73.2	(65.0-80.1)	22.9	(16.6-30.8)	1.6	(0.7-3.3)*	1.7	(0.8-3.3)*	0.6	(0.2-1.8)*
	No	1,848	64.8	(61.3-68.2)	27.5	(24.3-30.9)	1.5	(0.5-4.0)*	4.5	(3.0-6.9)	1.7	(1.0-2.9)
Diabetes	Yes	124	69.3	(57.4-79.0)	17.8	(10.1-29.5)	1.4	(0.4-4.2)*	0.4	(0.2-1.1)*	11.2	(5.3-22.1)*
	No	2,096	66.1	(62.7-69.4)	27.0	(24.0-30.3)	1.5	(0.6-3.5)*	4.2	(2.8-6.3)	1.2	(0.7-1.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Alaska, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	128	72.2	(61.6-80.8)	21.8	(13.9-32.5)	0.4	(0.1-1.9)*	3.3	(1.4-7.3)*	2.3	(0.8-6.2)*
	No	2,066	66.2	(62.7-69.5)	26.9	(23.8-30.1)	1.6	(0.7-3.6)*	4.0	(2.6-6.0)	1.4	(0.8-2.4)
Asthma	Yes	304	72.4	(61.9-80.9)	23.6	(15.9-33.5)	0.8	(0.3-1.9)*	2.5	(1.1-5.7)*	0.6	(0.2-1.9)*
	No	1,908	65.6	(62.1-68.8)	26.8	(23.7-30.1)	1.6	(0.7-3.8)*	4.3	(2.8-6.5)	1.7	(1.0-2.9)
Arthritis	Yes	617	72.1	(66.5-77.2)	23.7	(18.9-29.2)	0.9	(0.4-2.0)*	2.1	(1.2-3.6)	1.2	(0.6-2.3)*
	No	1,594	64.6	(60.6-68.4)	27.7	(24.2-31.4)	1.7	(0.7-4.5)*	4.5	(2.9-7.1)	1.5	(0.8-2.8)*
Frequent Mental Distress	Yes	210	61.2	(49.7-71.6)	25.7	(17.4-36.2)	1.3	(0.3-4.8)*	10.9	(3.7-28.1)*	0.9	(0.4-2.4)*
	No	1,949	67.6	(64.1-70.9)	26	(23.0-29.3)	1.5	(0.6-3.7)*	3.3	(2.2-4.8)	1.6	(0.9-2.7)
Serious Psychological Distress	Yes	52	84.3	(22.7-99.0)	9.9	(3.0-28.1)*	0.0	(0.0-100.0)*	1.1	(0.0-100.0)*	4.7	(2.1-10.3)*
	No	2,115	66.4	(63.2-69.5)	27.1	(24.2-30.3)	1.5	(0.6-3.5)*	3.4	(2.4-4.9)	1.5	(0.9-2.5)
Receiving Treatment for Mental Health Condition	Yes	223	73.2	(61.1-82.5)	16	(9.5-25.5)	1.2	(0.3-4.3)*	9.5	(2.9-26.7)*	0.3	(0.0-1.7)*
	No	1,993	65.6	(62.1-68.9)	27.8	(24.7-31.2)	1.5	(0.6-3.7)*	3.4	(2.4-4.9)	1.6	(1.0-2.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Alaska, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,227	22.5	(19.9-25.5)	38.7	(35.5-41.9)	2.4	(1.3-4.5)*	26.4	(23.6-29.3)	10.0	(8.3-12.0)
Age (years)	18-24	150	20.3	(12.8-30.7)	47.8	(36.7-59.0)	6.0	(1.6-20.4)*	21.0	(14.0-30.3)	4.9	(1.8-12.6)*
	25-34	370	20.5	(15.0-27.5)	38.6	(31.3-46.4)	1.2	(0.5-2.8)*	30.2	(23.2-38.2)	9.5	(6.2-14.3)
	35-54	1,040	21.0	(17.6-24.9)	37.7	(33.6-42.1)	1.4	(0.6-3.3)*	28.6	(24.5-33.1)	11.2	(8.6-14.5)
	55+	651	26.2	(21.7-31.4)	37.2	(31.6-43.1)	2.3	(1.1-4.8)*	23.2	(18.6-28.4)	11.1	(7.9-15.4)
Sex	Men	977	24.8	(20.7-29.6)	39.2	(34.4-44.2)	3.7	(1.7-8.0)*	25.0	(20.9-29.6)	7.3	(5.3-10.0)
	Women	1,250	20.3	(17.3-23.7)	38.4	(34.4-42.4)	1.1	(0.5-2.3)*	27.6	(24.2-31.3)	12.6	(10.1-15.7)
Race/ Ethnicity	Hispanic or Latino	58	28.9	(17.0-44.7)	33.1	(18.8-51.4)	3.8	(0.5-22.0)*	13.9	(4.6-35.0)*	20.4	(9.5-38.3)*
	Not Hispanic or Latino:	1,611	19.9	(17.1-23.0)	41.0	(37.3-44.8)	2.2	(1.0-4.8)*	27.4	(24.2-30.9)	9.5	(7.6-11.8)
	White											
	Black or African American	24	--	--	--	--	--	--	--	--	--	--
	All Other Races	513	30.8	(24.1-38.4)	33.3	(26.9-40.4)	3.0	(0.7-12.4)*	22.6	(17.4-28.7)	10.4	(6.6-15.9)
Education	<High School	151	34.3	(23.2-47.5)	32.2	(20.2-47.0)	8.7	(1.5-37.9)*	19.6	(11.7-30.9)	5.2	(2.4-10.9)*
	High School Graduate (or GED)	664	30.4	(24.4-37.1)	35.3	(29.7-41.3)	2.3	(0.6-8.5)*	22.4	(18.2-27.4)	9.6	(6.8-13.4)
	Some College	649	20.3	(16.4-24.9)	42.1	(36.5-47.9)	1.7	(0.8-3.5)*	23.7	(18.8-29.5)	12.1	(8.7-16.8)
	College Graduate	759	13.5	(10.3-17.5)	40.6	(35.4-46.1)	1.9	(0.8-4.3)*	33.9	(28.8-39.4)	10.1	(7.4-13.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Alaska, 2007, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	272	27.2 (18.2-38.5)	34.1 (24.9-44.8)	1.2 (0.2-6.2)*	30.7 (21.1-42.4)	6.7 (4.3-10.4)				
	\$20,000-\$34,999	323	23.1 (17.6-29.6)	35.1 (27.5-43.6)	0.9 (0.3-2.8)*	25.6 (18.6-34.0)	15.3 (9.8-23.2)				
	\$35,000-\$49,999	318	16.9 (11.9-23.4)	36.5 (28.2-45.6)	4.7 (0.9-21.1)*	32.6 (25.3-40.8)	9.3 (5.7-14.9)				
	\$50,000-\$74,999	412	18.1 (14.0-23.0)	43.1 (36.0-50.6)	1.3 (0.5-3.0)*	27.4 (21.4-34.2)	10.2 (6.2-16.2)				
	\$75,000 and more	690	25.3 (20.4-31.0)	41.3 (35.9-46.9)	1.8 (0.8-4.0)*	23.7 (19.4-28.5)	7.9 (5.7-10.8)				
Employment Status	Employed	1,513	22.5 (19.4-26.0)	38.9 (35.2-42.7)	1.9 (1.1-3.2)	27.4 (24.1-31.0)	9.3 (7.4-11.7)				
	Out of work	130	24.3 (13.1-40.6)	34.3 (22.2-48.8)	18.4 (6.2-43.7)*	20.2 (12.7-30.7)	2.8 (1.1-7.1)*				
	Homemaker	175	27.2 (18.2-38.5)	41.8 (31.8-52.5)	0.6 (0.2-2.2)*	16.6 (11.6-23.3)	13.8 (7.8-23.2)				
	Student	51	13.2 (5.8-27.4)*	55.2 (37.2-71.9)	0.4 (0.3-0.6)	25.4 (13.4-43.0)*	5.8 (1.3-22.5)*				
	Retired	244	23.1 (16.3-31.8)	36.0 (27.4-45.6)	0.5 (0.2-1.0)*	26.1 (18.0-36.3)	14.3 (8.2-23.7)				
	Unable to work	101	15.5 (9.0-25.4)	34.9 (22.0-50.4)	0.3 (0.2-0.5)	36.7 (21.4-55.3)	12.6 (6.9-21.8)				
Veteran Status	Yes	366	28.1 (21.1-36.5)	31.9 (25.2-39.4)	1.4 (0.6-3.2)*	29.0 (22.1-37.0)	9.6 (5.5-16.3)				
	No	1,857	21.3 (18.6-24.3)	40.1 (36.6-43.6)	2.7 (1.3-5.3)*	25.9 (22.9-29.0)	10.1 (8.3-12.3)				
Diabetes	Yes	123	21.2 (12.1-34.5)	24.7 (14.7-38.4)	0.6 (0.1-2.8)*	26.1 (17.0-37.8)	27.4 (17.0-41.0)				
	No	2,100	22.6 (19.9-25.6)	39.5 (36.2-42.9)	2.5 (1.3-4.8)*	26.4 (23.5-29.4)	8.9 (7.3-10.9)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Alaska, 2007, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	130	25.6	(16.2-37.9)	32.6	(22.2-45.1)	1.0	(0.2-4.1)*	23.6	(13.9-37.2)	17.2	(10.6-26.6)
	No	2,068	22.5	(19.7-25.6)	39.0	(35.7-42.4)	2.5	(1.3-4.8)*	26.4	(23.6-29.4)	9.6	(7.8-11.7)
Asthma	Yes	305	22.8	(15.5-32.3)	32.9	(25.3-41.6)	0.1	(0.0-0.6)*	31.7	(24.6-39.7)	12.5	(8.2-18.6)
	No	1,911	22.5	(19.8-25.6)	39.6	(36.1-43.1)	2.8	(1.5-5.2)*	25.6	(22.6-28.8)	9.5	(7.7-11.7)
Arthritis	Yes	614	18.9	(15.2-23.3)	41.2	(35.3-47.3)	1.5	(0.7-3.3)*	25.7	(20.8-31.2)	12.7	(9.3-17.2)
	No	1,601	24.0	(20.7-27.7)	38.0	(34.3-41.8)	2.8	(1.3-5.7)*	26.3	(23.1-29.8)	8.9	(7.1-11.3)
Frequent Mental Distress	Yes	207	18.3	(11.4-28.1)	34	(24.9-44.5)	1.0	(0.3-2.7)*	26.7	(17.2-39.0)	20.0	(13.1-29.4)
	No	1,948	23.3	(20.5-26.4)	39.6	(36.2-43.0)	2.1	(1.1-4.1)*	26.2	(23.3-29.2)	8.8	(7.1-10.9)
Serious Psychological Distress	Yes	52	15.6	(7.5-29.7)*	25.1	(10.5-49.0)*	0.4	(0.1-3.4)*	48.7	(23.1-75.0)*	10.2	(4.5-21.5)*
	No	2,117	22.3	(19.6-25.1)	39.3	(36.1-42.6)	2.5	(1.3-4.7)*	26.0	(23.2-28.9)	10.0	(8.3-12.1)
Receiving Treatment for Mental Health Condition	Yes	218	11.2	(7.1-17.4)	37.3	(28.1-47.5)	0.1	(0.0-0.7)*	36.6	(26.4-48.1)	14.8	(9.3-22.7)
	No	2,001	23.9	(21.0-27.1)	39.1	(35.7-42.5)	2.7	(1.4-5.1)*	25.1	(22.4-28.1)	9.2	(7.5-11.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,119	70.0	(68.2-71.6)	23.9	(22.3-25.5)	0.6	(0.4-0.9)	3.4	(2.7-4.2)	2.1	(1.5-2.9)
Age (years)	18-24	163	56.7	(48.5-64.5)	33.9	(26.5-42.2)	0.0	(0.0-0.0)	6.3	(3.6-10.8)	3.1	(1.1-8.2)*
	25-34	512	67.1	(62.4-71.4)	25.6	(21.7-30.0)	0.6	(0.2-2.0)*	4.6	(2.8-7.4)	2.1	(0.9-4.7)*
	35-54	1,750	72.1	(69.6-74.5)	22.2	(20.0-24.6)	0.8	(0.5-1.5)	2.6	(1.9-3.6)	2.2	(1.5-3.4)
	55+	2,662	74.3	(72.4-76.1)	20.9	(19.2-22.6)	0.8	(0.5-1.2)	2.5	(1.9-3.3)	1.5	(1.0-2.2)
Sex	Men	1,810	63.4	(60.5-66.2)	28.6	(25.9-31.5)	0.9	(0.5-1.4)	4.7	(3.6-6.2)	2.5	(1.6-3.7)
	Women	3,309	76	(74.1-77.9)	19.5	(17.8-21.3)	0.4	(0.3-0.7)	2.3	(1.6-3.1)	1.8	(1.1-2.9)
Race/ Ethnicity	Hispanic or Latino	149	77.2	(68.2-84.3)	18.6	(12.3-27.3)	0.1	(0.1-0.1)	2.4	(1.1-5.3)*	1.6	(0.2-10.1)*
	Not Hispanic or Latino:	4,335	69.8	(68.0-71.6)	24.4	(22.8-26.2)	0.7	(0.5-1.0)	3.4	(2.6-4.3)	1.7	(1.2-2.5)
	White											
	Black or African American	374	65.3	(58.2-71.7)	25.8	(19.3-33.6)	0.3	(0.1-1.0)*	4.6	(2.5-8.3)*	4.0	(1.9-8.1)*
	All Other Races	201	76.5	(67.8-83.3)	17.2	(11.3-25.2)	1.3	(0.3-5.0)*	2.9	(1.2-6.7)*	2.2	(0.8-6.0)*
Education	<High School	581	58.2	(52.8-63.4)	28.2	(23.0-34.0)	1.2	(0.5-2.9)*	4.4	(2.7-7.1)	8.0	(5.3-12.0)
	High School Graduate (or GED)	1,812	65.6	(62.5-68.6)	27.9	(25.1-30.9)	0.8	(0.5-1.4)	3.6	(2.5-5.1)	2.1	(1.2-3.5)
	Some College	1,240	70.6	(67.0-74.1)	22.8	(19.6-26.4)	0.6	(0.3-1.0)*	4.0	(2.6-6.1)	2.0	(1.2-3.4)
	College Graduate	1,477	76.7	(73.9-79.3)	19.3	(17.0-22.0)	0.5	(0.3-1.0)*	2.5	(1.6-4.0)	0.9	(0.4-2.1)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	959	65.8 (61.5-69.8)	25.8 (22.0-30.0)	0.8 (0.4-1.4)*	4.9 (3.2-7.2)	2.9 (1.4-5.6)*				
	\$20,000-\$34,999	1,053	69.7 (65.9-73.3)	24.3 (21.0-27.9)	0.9 (0.4-2.0)*	4.1 (2.7-6.2)	1.0 (0.5-1.9)*				
	\$35,000-\$49,999	718	72.0 (67.5-76.0)	20.1 (16.5-24.3)	0.7 (0.3-2.0)*	4.6 (2.8-7.6)	2.5 (1.2-5.1)*				
	\$50,000-\$74,999	784	74.4 (70.6-77.9)	21.4 (18.1-25.1)	0.4 (0.1-1.2)*	1.9 (1.1-3.5)*	1.9 (0.9-3.9)*				
	\$75,000 and more	953	71.0 (67.2-74.6)	25.2 (21.7-29.0)	0.2 (0.0-0.6)*	1.5 (0.8-2.8)*	2.1 (1.1-4.0)*				
Employment Status	Employed	2,509	69.7 (67.3-72.0)	23.4 (21.3-25.6)	0.6 (0.3-1.1)*	3.6 (2.6-4.9)	2.7 (1.7-4.1)				
	Out of work	160	67.3 (57.1-76.1)	29.4 (21.1-39.5)	0.2 (0.0-1.3)*	2.5 (1.1-5.3)*	0.6 (0.1-4.2)*				
	Homemaker	480	67.5 (62.2-72.5)	25.9 (21.4-31.0)	0.3 (0.1-1.0)*	3.6 (1.7-7.2)*	2.7 (1.2-5.7)*				
	Student	77	75.8 (62.7-85.4)	20 (10.7-34.4)*	0.0 (0.0-0.0)	3.6 (1.2-10.5)*	0.5 (0.1-3.9)*				
	Retired	1,414	69.2 (66.3-71.9)	25 (22.5-27.7)	0.9 (0.6-1.5)	3.3 (2.4-4.5)	1.6 (0.9-2.8)				
	Unable to work	458	73.1 (68.0-77.7)	21.5 (17.3-26.4)	1.1 (0.5-2.5)*	2.9 (1.7-5.0)	1.4 (0.7-2.8)*				
Veteran Status	Yes	744	67.7 (63.7-71.5)	25.7 (22.2-29.4)	0.8 (0.3-1.7)*	3.2 (2.2-4.8)	2.6 (1.4-4.8)*				
	No	4,373	70.3 (68.4-72.1)	23.6 (21.9-25.4)	0.6 (0.4-0.9)	3.4 (2.7-4.4)	2.0 (1.4-2.9)				
Diabetes	Yes	587	71.5 (67.0-75.6)	22.2 (18.4-26.5)	1.2 (0.6-2.5)*	3.1 (1.9-4.9)	2.0 (1.0-4.0)*				
	No	4,527	69.7 (67.9-71.5)	24.0 (22.3-25.8)	0.7 (0.5-0.9)	3.4 (2.7-4.3)	2.2 (1.6-3.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	681	70.4	(66.3-74.2)	23.2	(19.6-27.2)	1.4	(0.8-2.6)*	3.0	(1.9-4.7)	2.0	(1.1-3.7)*
	No	4,381	69.9	(68.0-71.7)	24	(22.3-25.8)	0.5	(0.4-0.8)	3.4	(2.7-4.3)	2.1	(1.5-3.0)
Asthma	Yes	607	68.7	(63.4-73.5)	22.7	(18.3-27.8)	0.5	(0.2-1.0)*	3.9	(2.2-6.7)	4.3	(2.3-8.1)*
	No	4,498	69.8	(68.0-71.6)	24.1	(22.4-25.8)	0.8	(0.6-1.1)	3.4	(2.7-4.2)	2.0	(1.4-2.8)
Arthritis	Yes	2,108	71.4	(68.9-73.7)	23.4	(21.2-25.8)	0.6	(0.4-1.1)	2.7	(1.9-3.6)	1.9	(1.3-3.0)
	No	2,988	69.3	(67.1-71.5)	24.1	(22.1-26.2)	0.6	(0.4-1.0)	3.7	(2.9-4.8)	2.2	(1.5-3.3)
Frequent Mental Distress	Yes	546	65.0	(59.3-70.2)	25.0	(20.4-30.2)	0.6	(0.2-1.7)*	6.5	(4.3-9.6)	2.9	(1.4-6.3)*
	No	4,463	70.5	(68.7-72.3)	23.8	(22.1-25.5)	0.6	(0.4-0.9)	3.1	(2.4-4.0)	2.0	(1.4-2.9)
Serious Psychological Distress	Yes	274	57.8	(49.7-65.5)	32.6	(25.4-40.7)	1.4	(0.5-3.6)*	6.9	(4.2-11.2)	1.4	(0.6-3.3)*
	No	4,733	70.7	(68.9-72.4)	23.2	(21.6-24.9)	0.7	(0.5-1.0)	3.2	(2.5-4.1)	2.2	(1.6-3.1)
Receiving Treatment for Mental Health Condition	Yes	717	75.3	(71.1-79.1)	19.6	(16.2-23.6)	0.6	(0.2-1.6)*	2.9	(1.7-5.0)	1.6	(0.7-3.4)*
	No	4,387	69.3	(67.5-71.1)	24.3	(22.6-26.2)	0.6	(0.4-1.0)	3.5	(2.8-4.4)	2.2	(1.5-3.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,083	22.8	(21.3-24.3)	34.8	(33.1-36.6)	1.4	(1.0-1.8)	28.1	(26.5-29.8)	12.9	(11.8-14.1)
Age (years)	18-24	164	16.4	(11.1-23.4)	44.9	(36.7-53.4)	0.0	(0.0-0.0)	29.8	(22.9-37.8)	8.9	(5.4-14.2)
	25-34	512	19.8	(16.3-23.8)	34.8	(30.4-39.4)	2.5	(1.4-4.7)*	30.8	(26.7-35.2)	12.1	(9.3-15.6)
	35-54	1,741	22.0	(19.9-24.2)	34.6	(32.1-37.2)	1.5	(0.9-2.3)	27.5	(25.2-29.8)	14.5	(12.8-16.5)
	55+	2,636	28.3	(26.5-30.3)	31.2	(29.2-33.2)	1.3	(0.9-1.8)	26.4	(24.5-28.3)	12.8	(11.5-14.2)
Sex	Men	1,812	24.6	(22.2-27.2)	38	(35.1-40.9)	1.5	(1.0-2.4)	25.4	(23.1-27.9)	10.4	(8.8-12.3)
	Women	3,271	21.0	(19.4-22.7)	31.8	(29.8-33.9)	1.2	(0.9-1.7)	30.7	(28.6-32.9)	15.2	(13.7-16.8)
Race/ Ethnicity	Hispanic or Latino	152	35.7	(27.0-45.5)	32.8	(23.1-44.2)	0.9	(0.2-4.2)*	22.1	(14.0-33.1)	8.5	(3.9-17.5)*
	Not Hispanic or Latino:	4,298	19.2	(17.8-20.6)	36.2	(34.3-38.1)	1.5	(1.1-2.0)	29.9	(28.2-31.8)	13.2	(12.0-14.5)
	White											
	Black or African American	375	27	(20.6-34.5)	34.1	(26.9-42.2)	0.6	(0.2-1.3)*	25.8	(20.0-32.5)	12.6	(9.1-17.1)
	All Other Races	197	29.6	(21.7-38.9)	24.6	(17.3-33.7)	3.1	(1.1-8.1)*	23.7	(17.4-31.5)	19.0	(13.0-26.8)
Education	<High School	565	32.6	(27.7-37.9)	36.1	(30.2-42.6)	1.2	(0.6-2.4)*	18.2	(14.4-22.9)	11.8	(8.7-15.7)
	High School Graduate (or GED)	1,812	25.9	(23.5-28.6)	35.9	(33.0-39.0)	1.4	(0.9-2.2)	24.7	(22.0-27.5)	12.1	(10.3-14.1)
	Some College	1,224	21.1	(17.9-24.7)	32.5	(28.9-36.3)	2.0	(1.2-3.3)	29.3	(26.0-32.9)	15.1	(12.7-17.8)
	College Graduate	1,475	17.9	(15.7-20.4)	35.2	(32.4-38.2)	0.9	(0.5-1.5)	33.4	(30.7-36.3)	12.5	(10.7-14.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	957	26.9 (23.2-31.0)	27.6 (23.7-32.0)	1.1 (0.6-2.0)*	27.1 (23.1-31.6)	17.3 (14.2-20.8)				
	\$20,000-\$34,999	1,050	23.2 (20.2-26.5)	35.3 (31.5-39.3)	1.2 (0.7-2.2)*	26.1 (22.7-29.8)	14.2 (11.7-17.3)				
	\$35,000-\$49,999	715	21.3 (18.2-24.8)	34.8 (30.5-39.5)	1.8 (0.8-4.1)*	29.4 (25.4-33.8)	12.7 (9.9-16.0)				
	\$50,000-\$74,999	790	20.2 (17.2-23.7)	36.9 (33.1-40.9)	1.1 (0.6-2.1)*	29.5 (26.0-33.4)	12.2 (9.9-14.9)				
	\$75,000 and more	945	20.1 (17.0-23.6)	37.0 (33.3-40.8)	1.7 (1.0-2.9)	30.0 (26.7-33.4)	11.3 (9.2-13.7)				
Employment Status	Employed	2,503	21.9 (19.9-24.0)	36.4 (34.0-38.8)	1.7 (1.2-2.5)	28.1 (26.0-30.4)	11.9 (10.4-13.5)				
	Out of work	158	21.5 (13.7-32.2)	35.4 (26.0-46.2)	0.7 (0.2-2.3)*	26.1 (17.9-36.5)	16.2 (10.4-24.5)				
	Homemaker	471	22.0 (18.1-26.5)	36.4 (31.3-41.8)	1.3 (0.6-2.8)*	26.8 (22.3-31.9)	13.4 (10.2-17.3)				
	Student	77	27.3 (15.6-43.3)	30.8 (18.3-46.8)	1.1 (0.3-3.8)*	24.2 (15.2-36.3)	16.6 (9.4-27.6)				
	Retired	1,403	25.8 (23.4-28.3)	32.6 (29.8-35.5)	1.4 (0.9-2.1)	29.5 (26.8-32.4)	10.8 (9.1-12.7)				
	Unable to work	450	21.0 (17.2-25.4)	26.6 (21.4-32.5)	0.5 (0.3-1.0)*	30.0 (24.4-36.2)	21.9 (17.5-27.2)				
Veteran Status	Yes	743	21.5 (18.6-24.6)	32.7 (28.8-36.8)	0.9 (0.4-1.9)*	30.4 (26.6-34.6)	14.6 (11.5-18.3)				
	No	4,338	22.9 (21.3-24.7)	35.1 (33.2-37.1)	1.5 (1.2-2.0)	27.7 (26.0-29.5)	12.7 (11.5-13.9)				
Diabetes	Yes	580	25.5 (22.0-29.4)	28.9 (24.3-33.9)	2.4 (1.2-4.7)*	28.8 (24.2-33.8)	14.4 (11.6-17.9)				
	No	4,499	22.5 (20.9-24.1)	35.4 (33.5-37.3)	1.3 (1.0-1.8)	28.1 (26.4-29.8)	12.7 (11.5-14.0)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Arkansas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	671	24.6	(21.2-28.2)	29.6	(25.7-33.9)	0.7	(0.3-1.8)*	27.9	(23.7-32.5)	17.2	(14.1-20.8)
	No	4,352	22.5	(20.9-24.2)	35.4	(33.5-37.4)	1.5	(1.1-2.0)	28.1	(26.4-29.9)	12.5	(11.3-13.8)
Asthma	Yes	602	19.8	(16.2-24.0)	30.8	(25.6-36.4)	0.7	(0.3-1.6)*	32.7	(27.7-38.1)	16.1	(12.9-19.8)
	No	4,467	23.1	(21.5-24.7)	35.5	(33.6-37.4)	1.4	(1.0-1.8)	27.5	(25.9-29.3)	12.5	(11.3-13.8)
Arthritis	Yes	2,090	22.3	(20.4-24.4)	33.4	(30.9-36.0)	1.1	(0.8-1.7)	28.1	(25.8-30.6)	15.0	(13.3-16.8)
	No	2,970	22.9	(21.0-24.9)	35.4	(33.1-37.7)	1.5	(1.1-2.1)	28.2	(26.2-30.3)	12.1	(10.7-13.7)
Frequent Mental Distress	Yes	549	18.5	(14.5-23.3)	26.6	(21.8-32.1)	0.5	(0.3-0.8)	33.4	(28.2-39.0)	21.0	(17.3-25.3)
	No	4,427	23.4	(21.8-25.0)	35.6	(33.8-37.6)	1.6	(1.2-2.1)	27.4	(25.7-29.1)	12.0	(10.9-13.3)
Serious Psychological Distress	Yes	270	15.4	(9.8-23.2)	21.6	(15.9-28.6)	0.0	(0.0-0.0)	35.7	(28.1-44.0)	27.4	(20.7-35.2)
	No	4,707	23.2	(21.7-24.7)	35.5	(33.6-37.3)	1.4	(1.1-1.8)	27.8	(26.2-29.5)	12.2	(11.0-13.4)
Receiving Treatment for Mental Health Condition	Yes	709	20.0	(16.7-23.9)	27.8	(23.8-32.2)	1.5	(0.6-3.9)*	30.8	(26.6-35.3)	19.8	(16.5-23.7)
	No	4,360	23.1	(21.5-24.7)	35.7	(33.8-37.7)	1.4	(1.0-1.8)	27.9	(26.2-29.7)	12.0	(10.8-13.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,627	63.5	(61.2-65.7)	29.3	(27.2-31.5)	0.9	(0.5-1.8)*	4.8	(3.9-5.9)	1.4	(1.0-2.0)
Age (years)	18-24	147	51.8	(42.5-61.0)	41.2	(32.4-50.7)	3.1	(0.9-9.9)*	3.6	(1.7-7.4)*	0.2	(0.0-1.2)*
	25-34	348	58.7	(52.4-64.8)	32.2	(26.2-38.7)	1.0	(0.4-2.5)*	7.1	(4.3-11.6)	1.0	(0.4-3.0)*
	35-54	1,270	66.4	(63.1-69.5)	27.8	(24.9-30.9)	0.3	(0.1-0.8)*	4.1	(2.9-5.7)	1.4	(0.8-2.5)
	55+	1,862	66.0	(63.1-68.9)	25.2	(22.7-28.0)	0.7	(0.3-1.3)*	5.2	(3.9-6.9)	2.8	(1.9-4.2)
Sex	Men	1,424	59.0	(55.4-62.6)	31.9	(28.5-35.5)	1.3	(0.5-3.6)*	6.2	(4.6-8.2)	1.5	(0.9-2.6)
	Women	2,203	67.6	(64.9-70.2)	26.9	(24.3-29.6)	0.6	(0.3-1.2)*	3.6	(2.7-4.7)	1.4	(0.9-2.1)
Race/ Ethnicity	Hispanic or Latino	778	57.0	(52.4-61.5)	31.0	(26.8-35.6)	1.7	(0.4-6.5)*	7.2	(5.3-9.6)	3.1	(2.1-4.7)
	Not Hispanic or Latino:	2,431	64.3	(61.6-66.9)	28.5	(26.0-31.2)	0.9	(0.6-1.2)	4.9	(3.9-6.2)	1.4	(1.0-1.8)
	White											
	Black or African American	111	60.7	(47.4-72.6)	19.9	(11.2-33.1)	2.6	(0.8-8.2)*	8.9	(4.8-15.9)*	7.9	(4.6-13.2)
	All Other Races	301	55.4	(48.0-62.6)	35.5	(28.6-43.0)	0.9	(0.2-3.6)*	4.5	(2.4-8.3)*	3.7	(1.8-7.4)*
Education	<High School	425	58.9	(51.6-65.7)	35.2	(28.4-42.6)	0.3	(0.1-1.3)*	4.4	(2.3-8.0)*	1.3	(0.6-2.8)*
	High School Graduate (or GED)	706	57.0	(51.5-62.4)	34.1	(29.1-39.4)	0.3	(0.1-1.4)*	6.2	(4.2-9.1)	2.4	(1.3-4.4)*
	Some College	919	62.5	(57.9-66.9)	30.0	(25.6-34.8)	2.3	(0.7-7.2)*	4.3	(2.9-6.3)	0.9	(0.5-1.7)*
	College Graduate	1,575	69.9	(66.7-72.9)	23.7	(20.9-26.7)	0.8	(0.4-1.8)*	4.3	(2.8-6.3)	1.3	(0.7-2.5)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	762	57.1	(51.8-62.2)	33.4	(28.5-38.7)	2.3	(0.6-8.4)*	5.9	(3.7-9.1)	1.4	(0.8-2.5)
	\$20,000-\$34,999	550	63.2	(56.8-69.2)	28.1	(22.7-34.2)	0.5	(0.1-1.9)*	6.9	(4.3-10.7)	1.4	(0.6-2.8)*
	\$35,000-\$49,999	424	64.1	(57.1-70.6)	28.9	(22.8-35.8)	1.4	(0.5-4.1)*	4.4	(2.5-7.6)	1.2	(0.5-2.6)*
	\$50,000-\$74,999	513	66.7	(60.7-72.3)	28.3	(23.0-34.2)	0.5	(0.1-1.6)*	3.4	(1.7-6.6)*	1.1	(0.4-3.3)*
	\$75,000 and more	1,177	68.3	(64.6-71.8)	27.1	(23.7-30.7)	0.3	(0.2-0.7)*	3.4	(2.2-5.2)	0.9	(0.4-1.9)*
Employment Status	Employed	1,746	65.8	(62.7-68.7)	28.5	(25.6-31.5)	0.6	(0.3-1.3)*	3.8	(2.6-5.4)	1.4	(0.8-2.3)
	Out of work	292	59.2	(51.3-66.7)	33.9	(26.5-42.1)	0.5	(0.1-2.3)*	6.0	(3.3-10.6)	0.4	(0.1-2.1)*
	Homemaker	344	57.7	(51.0-64.1)	34.6	(28.4-41.4)	0.6	(0.1-3.2)*	5.3	(3.0-9.3)	1.7	(0.6-4.8)*
	Student	92	65.7	(54.2-75.6)	23.8	(15.6-34.4)	4.5	(1.2-15.1)*	4.5	(1.5-12.1)*	1.6	(0.3-7.3)*
	Retired	936	59.1	(54.7-63.3)	32.0	(27.9-36.3)	0.6	(0.3-1.4)*	6.1	(4.2-8.9)	2.2	(1.1-4.1)*
	Unable to work	217	65.3	(56.5-73.1)	25.9	(19.0-34.1)	0.2	(0.0-1.3)*	7.5	(4.5-12.2)	1.2	(0.5-2.6)*
Veteran Status	Yes	439	57.4	(50.7-63.8)	35.0	(28.5-42.0)	0.6	(0.2-2.0)*	5.2	(2.5-10.4)*	1.8	(0.8-4.3)*
	No	3,177	64.2	(61.8-66.6)	28.6	(26.4-31.0)	1.0	(0.5-2.0)*	4.8	(3.8-6.0)	1.4	(0.9-2.0)
Diabetes	Yes	363	65.6	(58.7-71.9)	26.3	(20.6-32.9)	0.3	(0.1-1.0)*	5.8	(3.5-9.4)	2.0	(0.9-4.4)*
	No	3,261	63.3	(61.0-65.6)	29.6	(27.4-31.9)	1.0	(0.5-1.9)*	4.7	(3.8-5.9)	1.4	(0.9-2.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	360	67.4	(60.5-73.5)	24.4	(19.1-30.8)	1.0	(0.3-3.7)*	5.7	(3.5-9.3)	1.5	(0.6-3.6)*
	No	3,262	63.2	(60.9-65.5)	29.7	(27.5-32.0)	0.9	(0.5-1.8)*	4.7	(3.8-5.9)	1.4	(1.0-2.1)
Asthma	Yes	504	66.2	(60.7-71.3)	25.3	(20.4-31.0)	0.2	(0.1-0.8)*	7.6	(4.9-11.6)	0.7	(0.3-1.6)*
	No	3,121	63.0	(60.5-65.4)	30.0	(27.8-32.4)	1.1	(0.6-2.0)*	4.4	(3.5-5.5)	1.5	(1.1-2.2)
Arthritis	Yes	1,150	61.4	(57.4-65.4)	31.1	(27.3-35.1)	0.8	(0.3-1.8)*	4.8	(3.4-6.8)	1.8	(1.1-3.2)
	No	2,473	64.1	(61.5-66.7)	28.7	(26.3-31.3)	1.0	(0.5-2.0)*	4.8	(3.8-6.1)	1.3	(0.8-2.0)
Frequent Mental Distress	Yes	398	59.6	(52.5-66.4)	31.7	(25.5-38.7)	0.3	(0.0-2.1)*	7.3	(4.6-11.4)	1.0	(0.5-2.4)*
	No	3,213	63.9	(61.6-66.2)	29.1	(26.9-31.4)	1.0	(0.5-2.0)*	4.4	(3.5-5.6)	1.5	(1.0-2.2)
Serious Psychological Distress	Yes	132	62.6	(51.5-72.5)	28.0	(19.2-38.8)	0.0	(0.0-0.0)	8.3	(4.0-16.4)*	1.1	(0.4-2.9)*
	No	3,479	63.7	(61.4-65.9)	29.6	(27.4-31.8)	0.6	(0.4-1.0)	4.7	(3.8-5.8)	1.4	(1.0-2.1)
Receiving Treatment for Mental Health Condition	Yes	407	72.7	(66.3-78.3)	16.6	(12.0-22.5)	0.4	(0.1-2.9)*	9.8	(6.2-15.4)	0.4	(0.1-1.2)*
	No	3,213	62.3	(59.9-64.7)	30.8	(28.6-33.2)	1.0	(0.5-1.9)*	4.3	(3.4-5.5)	1.5	(1.1-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,621	20.5	(18.8-22.4)	43.6	(41.3-45.8)	0.8	(0.5-1.3)	25.3	(23.4-27.3)	9.8	(8.5-11.2)
Age (years)	18-24	146	15.6	(10.5-22.6)	50.8	(41.1-60.3)	0.7	(0.1-4.4)*	29.8	(22.0-39.0)	3.2	(1.4-7.1)*
	25-34	347	18.6	(14.2-23.9)	38.8	(32.6-45.4)	0.3	(0.1-1.4)*	31.9	(25.8-38.8)	10.3	(6.6-15.8)
	35-54	1,271	21.3	(18.7-24.2)	45.4	(42.1-48.7)	0.8	(0.4-1.8)*	23.4	(20.8-26.3)	9.1	(7.3-11.2)
	55+	1,857	22.9	(20.3-25.6)	41	(38.0-43.9)	1.1	(0.7-1.8)	22.4	(20.1-24.8)	12.7	(10.5-15.3)
Sex	Men	1,421	23.4	(20.5-26.5)	43	(39.4-46.7)	1.1	(0.6-2.1)*	23.9	(20.9-27.2)	8.6	(6.7-11.0)
	Women	2,200	17.9	(15.8-20.1)	44.1	(41.2-47.0)	0.5	(0.3-0.9)*	26.7	(24.2-29.4)	10.9	(9.2-12.8)
Race/ Ethnicity	Hispanic or Latino	776	24.9	(21.3-28.8)	42.2	(37.4-47.1)	1.1	(0.5-2.0)*	23.8	(19.4-28.8)	8.2	(6.0-11.1)
	Not Hispanic or Latino:	2,430	19.2	(17.2-21.4)	42.5	(39.8-45.3)	1.4	(1.0-2.1)	26.6	(24.2-29.1)	10.3	(8.9-11.9)
	White											
	Black or African American	110	25.5	(16.1-37.8)	39.8	(28.3-52.6)	1.2	(0.3-4.3)*	18.3	(12.0-26.9)	15.2	(8.6-25.4)
	All Other Races	299	27.2	(21.2-34.2)	42.3	(34.9-50.1)	1.5	(0.7-3.0)*	19.0	(14.4-24.6)	10.0	(6.1-16.0)
Education	<High School	419	28.9	(23.0-35.6)	50.9	(43.4-58.5)	0.1	(0.0-0.4)*	14.0	(9.8-19.7)	6.0	(3.0-11.6)*
	High School Graduate (or GED)	710	25.3	(21.0-30.2)	38.9	(33.7-44.3)	0.6	(0.2-2.5)*	24.5	(20.1-29.6)	10.7	(7.9-14.2)
	Some College	920	18.2	(15.1-21.7)	44.3	(39.5-49.2)	0.9	(0.5-1.8)*	26.5	(22.3-31.2)	10.1	(7.9-12.9)
	College Graduate	1,570	15.8	(13.6-18.3)	44.2	(41.0-47.6)	1.2	(0.7-2.2)	28.4	(25.5-31.6)	10.3	(8.5-12.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	762	22.1 (18.4-26.4)	43.8 (38.3-49.5)	0.7 (0.2-2.6)*	22.8 (18.3-28.0)	10.6 (7.5-14.7)				
	\$20,000-\$34,999	546	22.7 (18.1-28.0)	36.5 (30.6-42.7)	0.6 (0.2-1.8)*	29.3 (23.6-35.8)	10.9 (7.6-15.4)				
	\$35,000-\$49,999	423	23.4 (18.2-29.7)	45.2 (38.2-52.4)	0.4 (0.2-1.0)*	21.7 (16.8-27.6)	9.2 (6.3-13.1)				
	\$50,000-\$74,999	513	17.8 (13.5-23.1)	42.8 (37.1-48.7)	1.2 (0.6-2.6)*	28.9 (24.0-34.3)	9.3 (6.6-12.9)				
	\$75,000 and more	1,176	19.4 (16.5-22.7)	46.3 (42.5-50.1)	0.8 (0.3-1.9)*	23.9 (20.9-27.1)	9.7 (7.7-12.1)				
Employment Status	Employed	1,744	21.8 (19.3-24.6)	42.4 (39.4-45.5)	0.9 (0.5-1.8)*	25.1 (22.6-27.9)	9.7 (7.9-11.9)				
	Out of work	294	19.2 (13.9-26.0)	35.8 (28.3-44.0)	0.3 (0.1-1.2)*	35.7 (27.8-44.6)	9.0 (5.2-15.1)				
	Homemaker	344	25.5 (20.2-31.7)	48.4 (41.6-55.2)	0.1 (0.0-0.9)*	20.1 (15.6-25.5)	5.9 (3.7-9.4)				
	Student	91	9.6 (5.1-17.6)*	53.4 (40.5-65.8)	0.4 (0.1-3.3)*	28.9 (18.5-42.1)	7.6 (3.2-17.1)*				
	Retired	932	20.3 (17.2-23.9)	46.6 (42.2-51.1)	1.4 (0.7-2.6)*	20.6 (17.5-24.0)	11.1 (8.5-14.3)				
	Unable to work	216	18.9 (12.8-27.2)	37.9 (29.6-47.0)	0.7 (0.1-5.3)*	26.2 (19.4-34.3)	16.3 (11.1-23.3)				
Veteran Status	Yes	439	21 (16.5-26.5)	37.4 (31.2-44.0)	1.0 (0.4-2.4)*	28.7 (21.5-37.1)	11.9 (8.1-17.3)				
	No	3,170	20.4 (18.6-22.4)	44.3 (41.9-46.8)	0.8 (0.4-1.3)	24.9 (22.9-27.1)	9.5 (8.2-11.0)				
Diabetes	Yes	361	18.3 (13.6-24.1)	47.6 (40.4-54.9)	2.4 (0.8-6.9)*	21.7 (16.7-27.6)	10.0 (6.5-15.1)				
	No	3,257	20.8 (19.0-22.7)	43.3 (40.9-45.7)	0.6 (0.4-1.1)	25.5 (23.5-27.7)	9.8 (8.4-11.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

California, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	358	20.0	(15.2-25.8)	43.9	(36.6-51.4)	1.5	(0.6-3.6)*	24.0	(18.6-30.3)	10.7	(7.7-14.7)
	No	3,258	20.6	(18.7-22.5)	43.6	(41.2-45.9)	0.7	(0.4-1.2)	25.4	(23.4-27.5)	9.7	(8.4-11.2)
Asthma	Yes	504	15.3	(11.6-20.1)	38.4	(32.6-44.5)	1.3	(0.5-3.3)*	34.2	(29.0-39.9)	10.8	(7.9-14.6)
	No	3,115	21.4	(19.5-23.4)	44.5	(42.1-47.0)	0.7	(0.4-1.2)	23.7	(21.7-25.9)	9.6	(8.3-11.2)
Arthritis	Yes	1,146	17.5	(14.7-20.8)	39.8	(35.9-43.8)	0.6	(0.3-1.3)*	27.9	(24.4-31.6)	14.2	(11.6-17.2)
	No	2,471	21.5	(19.5-23.7)	44.8	(42.2-47.5)	0.8	(0.5-1.5)	24.5	(22.3-26.9)	8.3	(6.9-9.9)
Frequent Mental Distress	Yes	401	14.7	(10.8-19.6)	44.9	(38.1-51.8)	0.6	(0.2-2.0)*	27.8	(22.3-34.1)	12.1	(8.5-17.0)
	No	3,205	21.4	(19.5-23.4)	43.4	(41.0-45.8)	0.8	(0.5-1.4)	25.0	(23.0-27.2)	9.3	(8.0-10.8)
Serious Psychological Distress	Yes	133	15.7	(9.9-23.8)	32.5	(22.9-43.8)	0.4	(0.1-3.1)*	32.9	(23.4-44.1)	18.5	(10.8-29.7)
	No	3,472	20.8	(19.0-22.7)	43.9	(41.6-46.2)	0.8	(0.5-1.3)	25.1	(23.2-27.2)	9.4	(8.2-10.9)
Receiving Treatment for Mental Health Condition	Yes	409	17.1	(12.9-22.3)	39.1	(32.6-46.0)	0.7	(0.4-1.4)*	25.3	(20.1-31.4)	17.7	(13.0-23.7)
	No	3,205	21.3	(19.5-23.3)	43.6	(41.2-46.0)	1.1	(0.7-1.6)	25.1	(23.1-27.3)	8.9	(7.6-10.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,241	66.5	(64.8-68.2)	27.0	(25.4-28.7)	0.8	(0.5-1.1)	4.2	(3.4-5.3)	1.4	(1.1-1.9)
Age (years)	18-24	190	51.8	(44.1-59.5)	37.0	(29.7-45.0)	0.5	(0.1-3.4)*	9.3	(5.5-15.4)	1.4	(0.5-3.8)*
	25-34	650	62.0	(57.7-66.1)	31.4	(27.5-35.5)	1.0	(0.4-2.3)*	4.2	(2.8-6.3)	1.5	(0.8-3.0)*
	35-54	2,156	69.5	(67.2-71.7)	25.7	(23.6-27.9)	0.6	(0.3-1.0)*	3.0	(2.2-4.1)	1.2	(0.8-1.9)
	55+	2,216	71.4	(69.2-73.5)	22.4	(20.5-24.4)	1.1	(0.7-1.6)	3.3	(2.6-4.4)	1.8	(1.2-2.6)
Sex	Men	2,076	60.5	(57.7-63.1)	31.4	(28.8-34.1)	0.9	(0.5-1.5)*	5.9	(4.5-7.9)	1.4	(0.9-2.2)
	Women	3,165	72.2	(70.2-74.2)	22.9	(21.0-24.9)	0.7	(0.5-1.1)	2.6	(1.9-3.6)	1.5	(1.0-2.2)
Race/ Ethnicity	Hispanic or Latino	589	68.7	(64.0-73.1)	23.9	(19.9-28.5)	0.3	(0.1-1.0)*	5.6	(3.5-8.9)	1.4	(0.8-2.7)*
	Not Hispanic or Latino:	4,295	66.6	(64.7-68.4)	27.6	(25.8-29.4)	1.1	(0.7-1.5)	3.4	(2.7-4.3)	1.4	(1.0-2.0)
	White											
	Black or African American	121	66.5	(56.5-75.2)	24.4	(16.8-34.0)	0.5	(0.1-3.4)*	5.3	(1.8-14.3)*	3.3	(1.3-8.2)*
	All Other Races	199	65.4	(56.4-73.4)	30.4	(22.8-39.3)	0.2	(0.1-0.9)*	3.2	(1.1-8.9)*	0.7	(0.3-2.0)*
Education	<High School	332	61.9	(54.7-68.6)	28.9	(22.7-36.0)	0.3	(0.1-0.8)*	7.3	(4.0-13.0)*	1.6	(0.8-3.3)*
	High School Graduate (or GED)	1,175	61.7	(57.8-65.4)	29.5	(26.1-33.3)	1.3	(0.7-2.5)*	5.9	(4.3-8.0)	1.6	(0.9-2.7)
	Some College	1,344	65.2	(61.8-68.4)	29.2	(26.1-32.5)	0.9	(0.5-1.7)*	3.2	(2.1-4.8)	1.6	(0.9-2.8)
	College Graduate	2,386	72.9	(70.6-75.0)	22.7	(20.6-24.9)	0.7	(0.4-1.1)	2.5	(1.8-3.6)	1.2	(0.8-1.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	554	66.4	(60.4-71.9)	25.4	(20.4-31.3)	0.7	(0.3-1.4)*	5.3	(2.9-9.5)*	2.1	(1.1-3.9)*
	\$20,000-\$34,999	803	63.2	(58.6-67.6)	30.5	(26.3-35.1)	0.5	(0.3-1.1)*	4.5	(2.5-8.0)	1.2	(0.6-2.6)*
	\$35,000-\$49,999	781	65.9	(61.5-70.0)	27.2	(23.4-31.4)	1.0	(0.4-3.1)*	4.3	(2.6-7.0)	1.6	(0.7-3.5)*
	\$50,000-\$74,999	854	67.1	(63.3-70.8)	26.2	(22.7-29.9)	1.4	(0.6-3.3)*	4.1	(2.6-6.6)	1.2	(0.5-2.6)*
	\$75,000 and more	1,740	71.1	(68.3-73.8)	24.7	(22.2-27.5)	0.6	(0.3-1.1)*	2.6	(1.8-3.9)	0.9	(0.5-1.7)*
Employment Status	Employed	3,193	66.8	(64.6-68.9)	26.7	(24.7-28.8)	0.9	(0.5-1.4)	4.4	(3.3-5.8)	1.3	(0.9-1.9)
	Out of work	169	70.5	(60.3-79.0)	26.3	(18.3-36.2)	0.1	(0.0-1.0)*	0.7	(0.2-1.9)*	2.4	(0.7-8.2)*
	Homemaker	499	64.3	(58.9-69.3)	32.4	(27.4-37.8)	0.5	(0.2-1.3)*	1.8	(0.8-3.7)*	1.1	(0.4-2.9)*
	Student	73	63.8	(50.7-75.1)	27.1	(16.9-40.4)	1.2	(0.2-8.2)*	5.8	(2.2-14.6)*	2.1	(0.4-11.2)*
	Retired	1,100	66.1	(62.8-69.2)	26.2	(23.2-29.4)	0.8	(0.4-1.5)*	4.7	(3.4-6.6)	2.2	(1.3-3.7)
	Unable to work	201	69.7	(60.8-77.3)	21.4	(15.0-29.7)	0.5	(0.2-1.7)*	7.3	(3.3-15.6)*	1.0	(0.4-2.6)*
Veteran Status	Yes	745	65.4	(61.2-69.3)	26.2	(22.6-30.1)	1.2	(0.4-3.2)*	5.8	(4.0-8.3)	1.5	(0.8-2.7)
	No	4,495	66.7	(64.8-68.6)	27.1	(25.4-29.0)	0.7	(0.5-1.1)	4.0	(3.1-5.1)	1.4	(1.0-2.0)
Diabetes	Yes	357	65.7	(59.5-71.4)	29.2	(23.7-35.3)	1.0	(0.4-2.3)*	3.4	(1.9-5.9)	0.8	(0.2-2.5)*
	No	4,884	66.5	(64.7-68.2)	26.9	(25.2-28.6)	0.8	(0.6-1.2)	4.3	(3.4-5.4)	1.5	(1.1-2.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	409	63.2	(57.5-68.5)	30.2	(25.0-36.0)	0.2	(0.0-1.5)*	4.8	(2.6-8.8)*	1.6	(0.7-3.8)*
	No	4,790	66.7	(64.9-68.5)	26.9	(25.2-28.6)	0.8	(0.6-1.2)	4.2	(3.3-5.2)	1.4	(1.0-1.9)
Asthma	Yes	660	67.0	(61.9-71.7)	24.9	(20.7-29.8)	0.7	(0.3-1.8)*	6.0	(3.5-10.0)	1.4	(0.6-2.9)*
	No	4,569	66.3	(64.5-68.1)	27.3	(25.6-29.1)	0.9	(0.6-1.3)	4.0	(3.2-5.1)	1.5	(1.1-2.0)
Arthritis	Yes	1,613	67.0	(64.1-69.7)	26.7	(24.1-29.6)	0.8	(0.4-1.5)*	3.7	(2.7-5.1)	1.9	(1.2-3.0)
	No	3,603	66.4	(64.3-68.4)	27.1	(25.2-29.1)	0.8	(0.5-1.3)	4.4	(3.4-5.6)	1.3	(0.9-1.9)
Frequent Mental Distress	Yes	453	68.9	(62.7-74.6)	25.9	(20.6-32.0)	0.2	(0.0-0.6)*	3.8	(2.3-6.2)	1.3	(0.5-3.4)*
	No	4,735	66.3	(64.5-68.1)	27.2	(25.5-28.9)	0.8	(0.6-1.2)	4.3	(3.3-5.4)	1.4	(1.0-1.9)
Serious Psychological Distress	Yes	144	66.5	(55.5-76.0)	23.1	(15.5-33.0)	0.4	(0.1-1.2)*	9.0	(4.8-16.4)*	0.9	(0.3-3.1)*
	No	5,020	66.4	(64.7-68.1)	27.1	(25.5-28.8)	0.9	(0.6-1.2)	4.1	(3.2-5.1)	1.5	(1.1-2.0)
Receiving Treatment for Mental Health Condition	Yes	619	73.9	(68.8-78.5)	21.4	(17.0-26.5)	0.2	(0.1-0.7)*	3.6	(2.0-6.4)	0.9	(0.4-2.0)*
	No	4,606	65.6	(63.8-67.4)	27.7	(26.0-29.5)	0.9	(0.6-1.3)	4.3	(3.4-5.4)	1.5	(1.1-2.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,234	23.5	(22.0-25.1)	37.5	(35.8-39.2)	1.2	(0.9-1.5)	28.7	(27.3-30.3)	9.1	(8.2-10.1)
Age (years)	18-24	191	20.2	(14.7-26.9)	41.4	(33.9-49.3)	0.0	(0.0-0.0)	30.6	(23.9-38.3)	7.8	(4.3-13.7)
	25-34	652	22.2	(18.9-25.9)	40.7	(36.7-44.9)	0.4	(0.1-1.3)*	29.9	(26.4-33.7)	6.8	(4.9-9.2)
	35-54	2,164	22.8	(20.7-25.0)	35.9	(33.6-38.1)	1.4	(1.0-2.2)	29.6	(27.6-31.7)	10.3	(9.0-11.8)
	55+	2,197	26.4	(24.3-28.6)	36.1	(33.9-38.4)	1.8	(1.3-2.4)	26.1	(24.2-28.2)	9.6	(8.3-11.0)
Sex	Men	2,091	26.6	(24.2-29.2)	40.3	(37.7-43.0)	1.0	(0.7-1.5)	25.4	(23.2-27.6)	6.7	(5.4-8.3)
	Women	3,143	20.5	(18.7-22.4)	34.7	(32.7-36.9)	1.4	(1.0-1.9)	32.0	(30.0-34.1)	11.4	(10.2-12.8)
Race/ Ethnicity	Hispanic or Latino	595	35.6	(31.0-40.5)	31.1	(26.5-36.2)	1.3	(0.7-2.6)*	21.8	(18.0-26.2)	10.1	(7.2-14.0)
	Not Hispanic or Latino:	4,277	20.0	(18.5-21.7)	39.6	(37.8-41.5)	1.2	(0.9-1.5)	31.2	(29.5-32.9)	8.0	(7.1-8.9)
	White											
	Black or African American	119	21.1	(13.8-30.8)	30.6	(21.9-40.9)	0.6	(0.1-3.0)*	31.8	(22.8-42.4)	16.0	(9.6-25.4)
	All Other Races	203	24.3	(18.2-31.7)	39.2	(31.1-47.9)	0.6	(0.1-3.0)*	21.3	(15.1-29.2)	14.6	(9.7-21.4)
Education	<High School	330	39.7	(32.9-46.9)	30.1	(23.5-37.6)	1.8	(0.8-4.1)*	21.6	(16.1-28.3)	6.8	(4.5-10.1)
	High School Graduate (or GED)	1,175	25.0	(22.0-28.3)	37.3	(33.6-41.1)	1.4	(0.9-2.1)	24.3	(21.2-27.6)	12.0	(9.4-15.2)
	Some College	1,342	23.4	(20.5-26.6)	34.6	(31.5-37.9)	0.9	(0.6-1.6)	32.9	(29.8-36.2)	8.1	(6.7-9.9)
	College Graduate	2,383	16.2	(14.4-18.2)	42.0	(39.6-44.4)	1.1	(0.8-1.7)	31.8	(29.6-34.0)	8.9	(7.7-10.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	552	24.5	(20.0-29.6)	31.9	(26.5-37.9)	1.4	(0.8-2.4)	30.0	(24.6-36.1)	12.2	(8.9-16.5)
	\$20,000-\$34,999	800	28.9	(24.8-33.3)	34.1	(29.9-38.6)	1.0	(0.5-1.8)*	26.0	(22.2-30.2)	10.1	(7.7-13.0)
	\$35,000-\$49,999	775	21.6	(18.1-25.6)	36.4	(32.1-40.8)	1.5	(0.9-2.6)	30.7	(26.6-35.2)	9.8	(7.8-12.4)
	\$50,000-\$74,999	861	22.7	(19.4-26.4)	34.4	(30.8-38.2)	0.4	(0.2-1.0)*	32.9	(29.3-36.7)	9.6	(7.6-12.0)
	\$75,000 and more	1,748	20.1	(17.7-22.8)	43.6	(40.7-46.5)	0.9	(0.6-1.5)	28.3	(26.0-30.7)	7.1	(5.6-8.8)
Employment Status	Employed	3,201	23.5	(21.5-25.6)	37.7	(35.6-39.9)	1.1	(0.8-1.5)	29.5	(27.6-31.4)	8.3	(7.2-9.6)
	Out of work	169	21.3	(14.1-30.7)	35.9	(27.2-45.7)	1.4	(0.4-4.6)*	29.8	(21.7-39.4)	11.7	(6.8-19.3)
	Homemaker	486	29.3	(24.5-34.6)	38.1	(33.1-43.4)	1.7	(0.9-3.2)*	21.6	(18.0-25.7)	9.3	(6.9-12.4)
	Student	76	15.2	(7.4-28.5)*	41.2	(28.7-54.9)	0.9	(0.2-3.8)*	34.1	(21.8-48.9)	8.7	(3.6-19.7)*
	Retired	1,097	24.8	(22.1-27.8)	39.1	(35.9-42.4)	1.9	(1.3-2.9)	26.7	(23.8-29.7)	7.4	(6.0-9.2)
	Unable to work	199	14.9	(10.0-21.6)	24.0	(17.1-32.5)	1.2	(0.4-3.4)*	35.6	(27.4-44.6)	24.4	(16.7-34.2)
Veteran Status	Yes	748	22.1	(18.8-25.7)	37.0	(33.1-41.1)	1.4	(0.9-2.4)	30.0	(26.1-34.2)	9.5	(7.2-12.5)
	No	4,485	23.7	(22.0-25.4)	37.5	(35.7-39.4)	1.2	(1.0-1.6)	28.5	(26.9-30.2)	9.0	(8.0-10.2)
Diabetes	Yes	356	25.0	(20.1-30.5)	32.5	(27.1-38.3)	2.6	(1.4-4.8)*	29.3	(24.2-35.1)	10.6	(7.5-14.9)
	No	4,877	23.4	(21.8-25.0)	37.8	(36.0-39.5)	1.2	(0.9-1.5)	28.7	(27.1-30.3)	9.0	(8.0-10.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Colorado, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	406	23.0	(18.6-28.0)	32.2	(26.9-37.9)	1.3	(0.7-2.6)*	28.9	(23.8-34.6)	14.6	(10.8-19.4)
	No	4,787	23.6	(22.0-25.3)	37.8	(36.0-39.5)	1.2	(0.9-1.6)	28.7	(27.1-30.3)	8.8	(7.8-9.8)
Asthma	Yes	649	17.6	(13.9-22.1)	36.4	(31.6-41.5)	0.5	(0.2-1.3)*	31.6	(27.0-36.5)	13.9	(10.0-19.0)
	No	4,571	24.4	(22.7-26.1)	37.7	(35.9-39.5)	1.3	(1.0-1.7)	28.2	(26.6-29.8)	8.4	(7.5-9.4)
Arthritis	Yes	1,608	19.2	(17.1-21.5)	37.3	(34.4-40.3)	1.5	(1.0-2.1)	30.4	(27.8-33.1)	11.6	(10.0-13.6)
	No	3,601	25.0	(23.1-27.0)	37.5	(35.5-39.5)	1.1	(0.8-1.5)	28.2	(26.4-30.0)	8.2	(7.1-9.5)
Frequent Mental Distress	Yes	445	17.3	(13.0-22.8)	30.8	(25.3-36.9)	0.6	(0.4-1.1)	33.3	(27.8-39.3)	17.9	(14.0-22.6)
	No	4,735	24.3	(22.6-25.9)	38.1	(36.4-39.9)	1.3	(1.1-1.7)	28.2	(26.7-29.8)	8.1	(7.2-9.1)
Serious Psychological Distress	Yes	145	10.8	(6.2-18.0)	28.0	(19.7-38.1)	0.6	(0.1-2.5)*	30.2	(21.5-40.6)	30.5	(21.9-40.7)
	No	5,012	24.0	(22.5-25.7)	37.9	(36.1-39.6)	1.2	(0.9-1.5)	28.7	(27.2-30.3)	8.2	(7.3-9.2)
Receiving Treatment for Mental Health Condition	Yes	617	17.6	(14.0-21.9)	36.1	(31.2-41.3)	0.8	(0.4-1.6)*	29.9	(25.7-34.4)	15.7	(12.8-19.1)
	No	4,601	24.2	(22.6-25.9)	37.6	(35.9-39.4)	1.3	(1.0-1.6)	28.6	(27.0-30.3)	8.3	(7.3-9.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,416	69.6	(67.9-71.2)	23.8	(22.3-25.4)	2.1	(1.6-2.8)	3.0	(2.5-3.8)	1.5	(1.0-2.1)
Age (years)	18-24	178	62.9	(53.9-71.0)	32.0	(24.3-40.8)	1.2	(0.3-4.6)*	2.6	(1.0-6.7)*	1.4	(0.3-5.5)*
	25-34	583	63.2	(58.1-68.0)	28.1	(23.8-32.9)	3.7	(1.7-7.8)*	3.7	(2.0-6.7)*	1.3	(0.3-5.2)*
	35-54	2,334	72.3	(69.9-74.6)	21.1	(19.0-23.4)	1.9	(1.3-2.9)	3.0	(2.1-4.2)	1.6	(1.0-2.6)
	55+	3,249	72.5	(70.6-74.4)	21.2	(19.5-23.1)	2.0	(1.5-2.7)	2.9	(2.3-3.7)	1.3	(0.9-2.0)
Sex	Men	2,408	63.2	(60.5-65.9)	28.6	(26.1-31.3)	2.6	(1.8-3.7)	3.3	(2.4-4.4)	2.3	(1.4-3.6)
	Women	4,008	75.5	(73.5-77.4)	19.2	(17.5-21.0)	1.6	(1.0-2.6)	2.8	(2.1-3.8)	0.8	(0.5-1.2)
Race/ Ethnicity	Hispanic or Latino	420	67.4	(59.7-74.3)	23.5	(17.9-30.1)	4.0	(1.3-11.3)*	4.1	(2.0-8.3)*	1.1	(0.6-2.1)*
	Not Hispanic or Latino:	5,341	70.7	(69.0-72.5)	23.5	(21.9-25.2)	1.8	(1.4-2.4)	2.6	(2.0-3.3)	1.3	(0.8-2.0)
	White											
	Black or African American	329	70.1	(62.2-77.0)	24.1	(17.8-31.8)	1.4	(0.5-4.5)*	3.0	(1.5-5.9)*	1.3	(0.5-3.2)*
	All Other Races	248	62.2	(52.9-70.7)	26.8	(19.2-36.2)	3.7	(0.8-16.3)*	6.2	(2.5-14.4)*	1.0	(0.2-5.2)*
Education	<High School	403	62.3	(54.6-69.3)	27.8	(21.5-35.2)	3.4	(0.9-11.9)*	4.2	(2.5-7.1)	2.3	(1.2-4.3)*
	High School Graduate (or GED)	1,560	65.1	(61.4-68.5)	26.7	(23.5-30.1)	2.5	(1.4-4.6)*	3.2	(2.3-4.5)	2.5	(1.4-4.6)*
	Some College	1,524	69.9	(66.4-73.1)	25.2	(22.0-28.6)	1.4	(0.8-2.3)	2.4	(1.7-3.5)	1.2	(0.7-2.2)*
	College Graduate	2,919	74.9	(72.6-77.1)	19.7	(17.7-21.9)	1.8	(1.3-2.6)	2.8	(1.9-4.2)	0.8	(0.4-1.5)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	685	64.7 (58.3-70.6)	26.9 (21.9-32.7)	3.0 (0.9-10.0)*	2.8 (1.7-4.7)	2.5 (1.3-4.8)*				
	\$20,000-\$34,999	876	66.1 (61.2-70.7)	23.8 (20.0-28.1)	2.3 (0.9-5.9)*	4.8 (2.9-7.9)	3.0 (1.3-6.8)*				
	\$35,000-\$49,999	764	69.4 (64.3-74.0)	25.0 (20.7-29.9)	2.1 (1.0-4.3)*	3.0 (1.7-5.2)	0.6 (0.2-1.7)*				
	\$50,000-\$74,999	940	70.4 (66.3-74.3)	24.5 (20.8-28.6)	2.3 (1.4-3.9)	2.0 (1.2-3.3)	0.8 (0.3-2.0)*				
	\$75,000 and more	2,339	74.7 (72.2-77.1)	20.4 (18.1-22.8)	1.8 (1.2-2.8)	2.4 (1.5-3.7)	0.7 (0.3-1.4)*				
Employment Status	Employed	3,609	69.6 (67.4-71.7)	24.2 (22.3-26.3)	2.0 (1.4-2.7)	3.0 (2.2-4.2)	1.2 (0.7-1.9)				
	Out of work	260	72.0 (63.4-79.3)	22.9 (15.9-31.8)	1.6 (0.6-4.2)*	2.1 (0.9-4.8)*	1.3 (0.5-3.7)*				
	Homemaker	501	71.4 (65.5-76.7)	22.3 (17.7-27.7)	3.2 (0.8-12.7)*	2.0 (0.9-4.1)*	1.1 (0.3-4.0)*				
	Student	123	79.7 (70.1-86.9)	18.5 (11.7-28.0)	0.0 (0.0-0.0)	1.7 (0.5-5.7)*	0.0 (0.0-0.0)				
	Retired	1,661	67.0 (64.1-69.8)	24.5 (22.0-27.3)	2.3 (1.5-3.4)	4.2 (3.1-5.7)	2.0 (1.2-3.4)				
	Unable to work	249	59.5 (51.1-67.4)	25.7 (18.5-34.4)	5.0 (1.2-18.6)*	4.0 (2.0-7.6)*	5.8 (1.7-17.9)*				
Veteran Status	Yes	766	68.2 (64.1-72.1)	24.2 (20.7-28.1)	2.8 (1.6-4.6)	2.7 (1.7-4.4)	2.1 (1.1-4.2)*				
	No	5,649	69.7 (67.9-71.5)	23.8 (22.1-25.5)	2.0 (1.4-2.8)	3.1 (2.5-3.9)	1.4 (0.9-2.1)				
Diabetes	Yes	606	71.5 (66.1-76.3)	20.5 (16.1-25.6)	1.5 (0.7-2.8)*	4.0 (2.1-7.3)*	2.6 (1.2-5.5)*				
	No	5,806	69.4 (67.6-71.1)	24.0 (22.4-25.7)	2.2 (1.6-2.9)	3.0 (2.4-3.8)	1.4 (1.0-2.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	581	64.9	(59.9-69.7)	26.5	(22.0-31.5)	2.2	(1.2-4.0)*	4.0	(2.5-6.5)	2.4	(1.2-4.7)*
	No	5,782	70.0	(68.2-71.7)	23.5	(21.9-25.2)	2.1	(1.5-2.9)	3.0	(2.4-3.8)	1.4	(1.0-2.1)
Asthma	Yes	844	71.1	(66.5-75.3)	22.9	(19.0-27.3)	1.2	(0.5-2.5)*	3.5	(1.9-6.2)*	1.4	(0.6-3.2)*
	No	5,555	69.2	(67.4-70.9)	23.9	(22.3-25.7)	2.3	(1.7-3.1)	3.0	(2.4-3.8)	1.6	(1.1-2.3)
Arthritis	Yes	2,182	71.4	(68.6-74.0)	22.0	(19.6-24.6)	1.5	(1.0-2.4)	3.7	(2.7-5.1)	1.4	(0.7-2.6)*
	No	4,216	69.0	(66.9-71.0)	24.3	(22.5-26.2)	2.3	(1.6-3.3)	2.9	(2.2-3.8)	1.5	(1.0-2.3)
Frequent Mental Distress	Yes	541	68.7	(62.9-74.0)	23.6	(18.8-29.2)	2.5	(0.7-8.5)*	2.3	(1.3-4.2)*	2.9	(0.8-9.8)*
	No	5,806	69.7	(67.9-71.4)	23.8	(22.2-25.4)	2.1	(1.6-2.8)	3.1	(2.5-3.9)	1.3	(1.0-1.9)
Serious Psychological Distress	Yes	188	61.9	(50.7-71.9)	23.1	(15.5-33.0)	1.4	(0.3-5.9)*	8.7	(3.6-19.4)*	4.9	(1.7-13.4)*
	No	6,132	70.2	(68.5-71.8)	23.8	(22.2-25.4)	1.9	(1.4-2.5)	2.9	(2.3-3.6)	1.3	(0.9-1.9)
Receiving Treatment for Mental Health Condition	Yes	837	78.9	(74.6-82.6)	15.8	(12.6-19.7)	2.7	(1.0-6.8)*	1.4	(0.7-2.6)*	1.3	(0.6-2.8)*
	No	5,561	68.4	(66.6-70.1)	24.8	(23.2-26.6)	2.0	(1.5-2.7)	3.3	(2.6-4.1)	1.5	(1.0-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,393	22.4	(21.0-23.8)	34.0	(32.4-35.6)	4.0	(3.4-4.7)	28.0	(26.5-29.6)	11.7	(10.6-12.8)
Age (years)	18-24	181	20.1	(14.0-28.0)	40.0	(31.6-49.0)	2.2	(0.8-6.3)*	27.0	(19.7-35.8)	10.7	(6.2-17.9)
	25-34	584	19.3	(15.7-23.6)	36.1	(31.6-41.0)	4.4	(2.6-7.5)	28.6	(24.5-33.2)	11.4	(8.5-15.3)
	35-54	2,327	19.6	(17.5-21.8)	32.7	(30.5-35.1)	4.1	(3.3-5.2)	30.5	(28.2-32.8)	13.1	(11.4-14.9)
	55+	3,232	28.3	(26.4-30.3)	32.2	(30.3-34.2)	4.1	(3.3-5.0)	24.7	(22.9-26.5)	10.7	(9.5-12.0)
Sex	Men	2,404	24.7	(22.4-27.2)	36.3	(33.8-38.9)	4.0	(3.2-5.0)	25.8	(23.6-28.2)	9.1	(7.5-11.0)
	Women	3,989	20.2	(18.6-21.9)	31.7	(29.7-33.7)	4.0	(3.2-5.0)	30.2	(28.3-32.1)	14.0	(12.6-15.5)
Race/ Ethnicity	Hispanic or Latino	410	38.0	(31.3-45.1)	26.8	(21.1-33.5)	5.4	(2.1-13.1)*	19.2	(14.6-24.9)	10.6	(7.0-15.7)
	Not Hispanic or Latino:	5,328	18.5	(17.1-19.9)	35.8	(34.0-37.6)	4.0	(3.4-4.7)	30.3	(28.6-32.0)	11.5	(10.3-12.7)
	White											
	Black or African American	327	32.1	(25.2-39.9)	25.4	(19.4-32.5)	3.9	(1.5-9.6)*	25.6	(19.3-33.2)	13.0	(7.7-21.2)
	All Other Races	249	27.0	(19.6-36.0)	30.1	(22.5-38.9)	3.6	(1.3-9.5)*	21.8	(14.3-31.6)	17.5	(11.0-26.7)
Education	<High School	400	38.3	(31.4-45.7)	26.9	(20.8-34.0)	3.0	(1.5-5.8)*	15.5	(10.9-21.6)	16.3	(10.0-25.5)
	High School Graduate (or GED)	1,556	26.5	(23.5-29.7)	34.6	(31.3-38.1)	3.2	(2.0-5.1)	25.4	(22.5-28.5)	10.4	(8.3-12.9)
	Some College	1,517	20.6	(17.9-23.6)	32.4	(29.1-35.8)	4.9	(3.6-6.6)	29.4	(26.1-33.0)	12.8	(10.8-15.1)
	College Graduate	2,911	15.8	(14.1-17.7)	36.1	(33.8-38.5)	4.1	(3.3-5.0)	32.1	(29.9-34.4)	11.9	(10.4-13.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	693	29.5 (24.7-34.8)	30.4 (25.2-36.1)	6.7 (3.3-13.2)*	20.6 (16.3-25.7)	12.7 (8.9-17.9)				
	\$20,000-\$34,999	864	27.1 (23.2-31.4)	31.8 (27.4-36.5)	3.0 (1.7-5.2)	24.9 (20.6-29.7)	13.2 (10.0-17.2)				
	\$35,000-\$49,999	753	23.2 (19.0-28.0)	35.9 (31.0-41.3)	3.5 (2.3-5.2)	27.0 (22.7-31.7)	10.4 (7.9-13.6)				
	\$50,000-\$74,999	935	18.9 (15.7-22.5)	39.3 (35.2-43.5)	3.8 (2.6-5.5)	27.6 (24.1-31.4)	10.4 (7.8-13.7)				
	\$75,000 and more	2,336	18.4 (16.2-20.7)	35.0 (32.6-37.6)	4.1 (3.2-5.2)	31.2 (28.9-33.6)	11.3 (9.8-13.0)				
Employment Status	Employed	3,602	22.1 (20.2-24.1)	33.8 (31.8-35.9)	4.1 (3.3-4.9)	27.9 (26.1-29.8)	12.1 (10.8-13.6)				
	Out of work	257	19.8 (14.1-26.9)	31.6 (23.9-40.6)	2.9 (1.2-6.9)*	32.7 (23.5-43.5)	13.0 (7.7-21.2)				
	Homemaker	506	19.6 (15.6-24.3)	37.1 (31.8-42.7)	6.2 (3.1-12.0)*	29.2 (24.6-34.2)	7.9 (5.6-11.2)				
	Student	123	20.4 (11.6-33.4)	39.7 (28.9-51.5)	2.1 (0.3-12.1)*	29.9 (20.7-41.1)	7.9 (3.1-18.4)*				
	Retired	1,642	27.4 (24.9-30.0)	34.0 (31.1-36.9)	3.9 (2.9-5.1)	25.2 (22.7-27.9)	9.5 (8.0-11.3)				
	Unable to work	250	17.3 (11.9-24.6)	28.3 (20.1-38.4)	2.8 (1.3-5.9)*	31.2 (23.4-40.2)	20.4 (12.9-30.5)				
Veteran Status	Yes	763	25.8 (22.2-29.7)	33.2 (29.3-37.3)	3.0 (2.0-4.6)	25.2 (21.5-29.3)	12.8 (9.6-16.8)				
	No	5,629	21.9 (20.4-23.5)	34.1 (32.4-35.9)	4.1 (3.5-4.9)	28.3 (26.7-30.0)	11.6 (10.5-12.8)				
Diabetes	Yes	608	23.7 (19.2-28.9)	30.5 (25.6-35.9)	4.1 (2.6-6.3)	29.7 (25.0-34.9)	12.0 (9.0-15.8)				
	No	5,780	22.2 (20.8-23.8)	34.3 (32.6-36.0)	4.0 (3.4-4.7)	27.9 (26.4-29.5)	11.6 (10.5-12.8)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Connecticut, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics		Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	573	23.9	(20.2-28.1)	32.9	(28.1-38.0)	2.4	(1.4-4.2)	28.4	(23.8-33.5)	12.4	(9.3-16.5)
	No	5,768	22.2	(20.7-23.7)	34.2	(32.5-35.9)	4.1	(3.5-4.9)	28.0	(26.5-29.6)	11.6	(10.4-12.8)
Asthma	Yes	852	16.9	(14.0-20.4)	31.4	(27.2-36.0)	4.2	(2.7-6.4)	29.7	(25.6-34.1)	17.8	(14.3-21.8)
	No	5,524	23.2	(21.6-24.8)	34.4	(32.7-36.2)	4.0	(3.3-4.7)	27.8	(26.2-29.4)	10.6	(9.5-11.8)
Arthritis	Yes	2,179	21.5	(19.5-23.8)	31.3	(28.7-34.0)	3.9	(3.1-5.0)	30.7	(28.2-33.3)	12.5	(10.8-14.4)
	No	4,197	22.6	(20.8-24.4)	34.8	(32.9-36.8)	4.0	(3.3-4.9)	27.2	(25.4-29.0)	11.4	(10.1-12.8)
Frequent Mental Distress	Yes	535	16.6	(12.6-21.5)	25.6	(20.6-31.4)	4.1	(2.5-6.5)	32.2	(27.1-37.8)	21.5	(16.6-27.4)
	No	5,785	23.1	(21.7-24.7)	34.7	(33.1-36.5)	3.9	(3.3-4.7)	27.7	(26.1-29.3)	10.5	(9.5-11.6)
Serious Psychological Distress	Yes	188	11.9	(7.6-18.2)	25.6	(17.3-36.2)	3.6	(1.4-9.4)*	31.0	(21.6-42.4)	27.8	(18.6-39.3)
	No	6,111	22.8	(21.4-24.3)	34.3	(32.7-36.0)	3.9	(3.3-4.7)	28.0	(26.5-29.6)	10.9	(9.9-12.0)
Receiving Treatment for Mental Health Condition	Yes	833	17.3	(13.8-21.5)	28.4	(24.4-32.8)	5.1	(3.5-7.3)	31.0	(26.9-35.3)	18.2	(14.6-22.6)
	No	5,542	23.1	(21.6-24.6)	34.7	(32.9-36.4)	3.8	(3.2-4.6)	27.7	(26.1-29.4)	10.7	(9.7-11.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,336	74.2	(72.1-76.3)	20.2	(18.3-22.3)	1.4	(0.9-2.2)	2.8	(2.1-3.6)	1.4	(1.0-2.0)
Age (years)												
	18-24	98	58.0	(46.8-68.3)	29.6	(20.1-41.2)	5.2	(1.6-15.4)*	5.6	(2.5-12.2)*	1.6	(0.4-6.4)*
	25-34	480	71.8	(66.4-76.6)	23.8	(19.3-29.0)	1.2	(0.5-3.0)*	1.8	(0.9-3.8)*	1.4	(0.6-3.0)*
	35-54	1,216	76.7	(73.5-79.7)	18.4	(15.7-21.6)	1.1	(0.6-2.1)*	2.2	(1.4-3.5)	1.5	(0.9-2.6)
	55+	1,504	77.6	(74.9-80.2)	17.2	(14.8-19.8)	1.0	(0.6-1.7)	3.1	(2.1-4.5)	1.1	(0.7-1.9)
Sex												
	Men	1,291	73.7	(70.1-76.9)	20.9	(17.9-24.3)	1.7	(0.8-3.3)*	2.5	(1.6-3.9)	1.2	(0.7-2.1)
	Women	2,045	75.2	(72.7-77.6)	19.2	(17.0-21.6)	1.2	(0.6-2.1)	2.9	(2.1-4.0)	1.5	(0.9-2.3)
Race/Ethnicity												
	Hispanic or Latino	183	69.8	(61.0-77.4)	24.4	(17.3-33.2)	1.0	(0.2-3.9)*	2.0	(0.7-5.2)*	2.8	(1.2-6.2)*
	Not Hispanic or Latino:	1,656	80.8	(78.1-83.2)	15.9	(13.7-18.4)	0.9	(0.5-1.8)*	1.6	(1.0-2.7)	0.8	(0.3-1.9)*
	White											
	Black or African American	1,283	69.0	(65.6-72.2)	22.0	(19.2-25.1)	2.0	(1.1-3.7)*	5.2	(3.8-7.1)	1.7	(1.1-2.7)
	All Other Races	158	71.0	(62.1-78.5)	22.5	(15.6-31.3)	1.5	(0.4-4.8)*	2.4	(0.7-7.7)*	2.7	(1.0-7.1)*
Education												
	<High School	234	74.1	(65.9-81.0)	18.7	(12.7-26.7)	0.7	(0.1-3.5)*	3.2	(1.7-5.9)*	3.2	(2.0-5.1)
	High School Graduate (or GED)	442	66.4	(60.4-72.0)	26.1	(20.8-32.3)	1.7	(0.8-3.8)*	3.8	(2.3-6.3)	1.9	(0.9-3.6)*
	Some College	532	70.1	(64.9-74.8)	23.2	(18.8-28.3)	2.7	(1.1-6.5)*	3.1	(1.9-5.3)	0.9	(0.3-2.3)*
	College Graduate	2,124	80.6	(78.2-82.8)	16.0	(13.9-18.2)	0.9	(0.5-1.6)*	1.6	(1.0-2.6)	0.9	(0.5-1.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	409	74.2 (68.3-79.3)	18.4 (13.8-24.0)	0.4 (0.1-1.5)*	4.7 (2.9-7.5)	2.3 (1.3-3.9)				
	\$20,000-\$34,999	386	66.2 (59.5-72.3)	27.1 (21.4-33.7)	0.7 (0.2-2.0)*	3.6 (2.1-6.0)	2.4 (1.3-4.4)*				
	\$35,000-\$49,999	344	68.9 (62.0-75.0)	25.8 (20.0-32.6)	0.8 (0.1-4.6)*	3.4 (1.7-6.7)*	1.1 (0.3-3.9)*				
	\$50,000-\$74,999	432	74.6 (68.4-79.9)	21.4 (16.4-27.5)	1.1 (0.4-3.1)*	2.2 (0.8-5.4)*	0.8 (0.2-3.5)*				
	\$75,000 and more	1,485	82.4 (79.5-84.9)	14.3 (12.0-16.9)	1.3 (0.6-2.6)*	1.3 (0.6-2.7)*	0.8 (0.3-1.8)*				
Employment Status	Employed	2,060	76.3 (73.7-78.7)	18.7 (16.5-21.2)	1.6 (0.9-2.9)	2.2 (1.5-3.1)	1.2 (0.7-2.0)				
	Out of work	153	62.6 (51.5-72.6)	29.2 (20.3-40.0)	3.2 (0.9-11.2)*	2.8 (1.1-6.9)*	2.2 (0.9-5.3)*				
	Homemaker	117	76.1 (64.4-84.9)	19.4 (11.8-30.4)	0.5 (0.1-3.2)*	3.6 (0.9-12.6)*	0.4 (0.1-1.7)*				
	Student	98	71.2 (59.8-80.5)	24.1 (15.5-35.5)	0.0 (0.0-0.0)	3.7 (1.2-10.9)*	1.0 (0.1-6.6)*				
	Retired	765	71.5 (67.2-75.5)	22.9 (19.2-27.1)	0.8 (0.4-1.8)*	3.3 (2.0-5.6)	1.4 (0.7-2.6)*				
	Unable to work	133	75.5 (64.7-83.8)	15.6 (8.5-27.0)	0.9 (0.2-4.2)*	5.0 (2.3-10.5)*	3.0 (1.4-6.3)*				
Veteran Status	Yes	369	73.5 (67.2-79.0)	21.0 (16.1-27.0)	1.6 (0.7-3.5)*	3.3 (1.4-7.7)*	0.6 (0.2-1.9)*				
	No	2,964	74.4 (72.1-76.6)	20.0 (18.0-22.2)	1.4 (0.8-2.3)	2.7 (2.1-3.6)	1.5 (1.0-2.1)				
Diabetes	Yes	285	73.2 (66.0-79.3)	21.3 (15.5-28.5)	0.7 (0.2-2.5)*	4.0 (1.9-8.0)*	0.9 (0.3-2.4)*				
	No	3,048	74.2 (72.0-76.4)	20.1 (18.1-22.3)	1.5 (1.0-2.4)	2.7 (2.0-3.5)	1.5 (1.0-2.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	241	74.4	(66.9-80.6)	20.2	(14.5-27.4)	0.4	(0.0-2.5)*	4.0	(1.8-8.8)*	1.1	(0.4-3.1)*
	No	3,062	74.1	(71.9-76.3)	20.3	(18.3-22.5)	1.5	(0.9-2.4)	2.7	(2.0-3.6)	1.4	(1.0-2.0)
Asthma	Yes	498	76.6	(71.6-80.9)	17.9	(14.0-22.6)	0.4	(0.2-0.8)*	3.0	(1.7-5.3)	2.1	(1.0-4.5)*
	No	2,829	73.7	(71.3-75.9)	20.6	(18.5-22.9)	1.7	(1.1-2.6)	2.8	(2.1-3.7)	1.3	(0.9-1.9)
Arthritis	Yes	1,026	74.1	(70.4-77.5)	20.2	(17.1-23.7)	1.2	(0.5-2.6)*	3.2	(2.1-5.0)	1.3	(0.8-2.2)
	No	2,298	74.2	(71.6-76.6)	20.3	(18.0-22.7)	1.5	(0.9-2.6)	2.6	(1.9-3.6)	1.4	(0.9-2.1)
Frequent Mental Distress	Yes	264	73.6	(65.9-80.1)	20.0	(14.2-27.5)	1.7	(0.5-5.2)*	3.4	(1.8-6.4)*	1.4	(0.5-3.5)*
	No	3,040	74.4	(72.1-76.5)	20.2	(18.2-22.4)	1.4	(0.9-2.3)	2.7	(2.0-3.6)	1.3	(0.9-1.9)
Serious Psychological Distress	Yes	77	73.7	(60.1-83.9)	16.7	(8.9-29.3)*	0.6	(0.1-5.2)*	2.2	(0.8-6.1)*	6.7	(3.5-12.5)*
	No	3,217	74.1	(71.9-76.2)	20.3	(18.4-22.5)	1.5	(0.9-2.3)	2.8	(2.1-3.6)	1.3	(0.9-1.9)
Receiving Treatment for Mental Health Condition	Yes	425	81.0	(75.1-85.8)	14.4	(10.0-20.1)	0.5	(0.1-2.2)*	2.1	(0.9-4.7)*	2.0	(1.0-4.2)*
	No	2,900	73.2	(70.9-75.4)	21.1	(19.0-23.3)	1.5	(0.9-2.5)	2.9	(2.2-3.8)	1.3	(0.9-1.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,308	21.5	(19.6-23.6)	33.3	(31.0-35.6)	3.1	(2.5-3.9)	28.8	(26.8-30.8)	13.3	(11.9-14.8)
Age (years)	18-24	98	20.1	(12.6-30.6)	46.3	(35.1-57.9)	1.3	(0.3-6.2)*	27.3	(18.3-38.8)	4.9	(2.3-10.2)*
	25-34	478	15.8	(11.9-20.6)	34.6	(29.5-40.1)	3.3	(2.0-5.5)	31.6	(27.2-36.3)	14.8	(11.3-19.1)
	35-54	1,215	20.9	(18.0-24.1)	31.3	(28.1-34.6)	3.0	(2.1-4.3)	29.6	(26.7-32.7)	15.3	(13.1-17.7)
	55+	1,478	26.9	(24.0-30.0)	31.1	(28.2-34.2)	3.6	(2.7-4.7)	26.3	(23.7-29.1)	12.2	(10.4-14.2)
Sex	Men	1,278	22.2	(19.2-25.5)	36.4	(32.8-40.1)	3.1	(2.2-4.3)	28.6	(25.6-31.7)	9.8	(7.8-12.1)
	Women	2,030	20.8	(18.5-23.3)	30.3	(27.8-33.0)	3.2	(2.4-4.2)	29.2	(26.8-31.7)	16.5	(14.6-18.6)
Race/ Ethnicity	Hispanic or Latino	180	32.9	(25.5-41.2)	36.5	(28.1-45.7)	1.3	(0.4-4.3)*	22.3	(16.0-30.1)	7.0	(4.1-11.8)
	Not Hispanic or Latino:	1,641	13.4	(11.0-16.1)	33.0	(30.2-35.8)	4.9	(3.8-6.3)	34.4	(31.7-37.2)	14.4	(12.4-16.6)
	White											
	Black or African American	1,276	26.5	(23.5-29.7)	31.1	(27.8-34.6)	1.6	(1.1-2.5)	23.9	(21.1-26.9)	16.9	(14.4-19.7)
	All Other Races	157	21.2	(14.5-29.9)	31.6	(23.7-40.7)	6.5	(3.1-13.1)*	24.4	(17.6-32.9)	16.3	(11.2-23.1)
Education	<High School	227	38.7	(31.0-47.1)	38.4	(29.3-48.4)	1.5	(0.5-4.4)*	14.9	(10.1-21.3)	6.5	(3.2-12.6)*
	High School Graduate (or GED)	442	28.4	(23.3-34.2)	36.1	(29.9-42.8)	1.0	(0.4-2.5)*	23.1	(18.0-29.0)	11.4	(8.5-15.2)
	Some College	530	19.7	(15.7-24.5)	37.8	(32.4-43.5)	2.8	(1.5-4.9)	26.2	(21.6-31.3)	13.5	(10.4-17.3)
	College Graduate	2,105	13.2	(11.3-15.4)	31.8	(29.3-34.4)	4.1	(3.2-5.2)	35.3	(32.7-37.9)	15.6	(13.7-17.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	397	27.8 (22.6-33.6)	29.2 (23.3-36.0)	1.8 (1.0-3.2)*	22.4 (17.4-28.3)	18.9 (13.7-25.4)				
	\$20,000-\$34,999	384	29.5 (24.0-35.7)	36.4 (29.7-43.6)	0.5 (0.2-1.4)*	23.0 (18.2-28.5)	10.6 (7.6-14.6)				
	\$35,000-\$49,999	342	20.8 (15.6-27.1)	39.0 (32.4-46.0)	1.3 (0.5-3.3)*	27.7 (22.0-34.1)	11.2 (7.8-15.8)				
	\$50,000-\$74,999	433	17.3 (13.0-22.8)	28.6 (23.1-34.8)	3.9 (2.2-6.9)	35.5 (29.8-41.6)	14.7 (11.4-18.8)				
	\$75,000 and more	1,472	15.0 (12.5-17.9)	35.8 (32.7-39.0)	4.4 (3.3-5.8)	31.7 (28.9-34.6)	13.2 (11.3-15.3)				
Employment Status	Employed	2,055	19.8 (17.5-22.3)	33.2 (30.5-35.9)	3.6 (2.8-4.7)	29.9 (27.6-32.4)	13.5 (11.8-15.4)				
	Out of work	149	15.1 (9.5-23.1)	36.3 (26.3-47.7)	0.7 (0.2-2.2)*	31.7 (22.8-42.2)	16.2 (10.2-24.9)				
	Homemaker	111	26.3 (17.3-37.7)	40.8 (29.1-53.6)	2.9 (1.0-8.3)*	16.9 (10.5-26.2)	13.2 (7.7-21.6)				
	Student	98	29.8 (18.9-43.7)	34.4 (23.9-46.8)	1.6 (0.2-10.1)*	24.4 (16.6-34.3)	9.7 (5.3-17.1)				
	Retired	750	25.5 (21.9-29.4)	33.3 (29.2-37.8)	2.9 (1.9-4.4)	28.8 (24.8-33.0)	9.6 (7.6-12.0)				
	Unable to work	134	23.5 (15.1-34.7)	24.2 (14.6-37.3)	2.0 (0.8-5.2)*	25.2 (17.0-35.8)	25.1 (16.5-36.3)				
Veteran Status	Yes	362	19.5 (14.9-25.1)	35.2 (29.1-41.8)	3.3 (1.7-6.3)*	28.6 (23.1-34.8)	13.3 (9.5-18.4)				
	No	2,943	21.7 (19.6-23.9)	33.1 (30.8-35.5)	3.1 (2.5-3.9)	28.8 (26.7-30.9)	13.3 (11.8-14.9)				
Diabetes	Yes	284	23.9 (18.6-30.2)	34.2 (27.1-42.1)	1.3 (0.6-2.5)*	27.7 (21.6-34.8)	12.9 (9.1-17.8)				
	No	3,021	21.3 (19.2-23.4)	33.2 (30.9-35.6)	3.3 (2.6-4.1)	28.9 (26.9-31.0)	13.3 (11.9-15.0)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

District of Columbia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	241	26.0	(19.2-34.1)	29.7	(22.8-37.8)	2.2	(1.0-5.0)*	25.9	(19.9-33.0)	16.1	(11.6-22.0)
	No	3,034	21.1	(19.1-23.2)	33.6	(31.3-36.0)	3.2	(2.5-4.0)	29.0	(27.0-31.1)	13.1	(11.6-14.7)
Asthma	Yes	501	24.0	(19.0-29.7)	32.1	(27.1-37.7)	2.9	(1.6-5.3)*	26.0	(21.8-30.7)	15.0	(11.8-18.8)
	No	2,799	21.1	(19.0-23.3)	33.5	(31.1-36.0)	3.2	(2.5-4.0)	29.3	(27.2-31.6)	13.0	(11.5-14.6)
Arthritis	Yes	1,012	23.5	(20.1-27.3)	33.6	(29.7-37.7)	2.6	(1.7-4.1)	27.3	(24.0-30.9)	12.9	(10.7-15.5)
	No	2,284	20.8	(18.6-23.3)	33.3	(30.7-36.0)	3.3	(2.6-4.3)	29.2	(26.9-31.5)	13.4	(11.7-15.2)
Frequent Mental Distress	Yes	266	13.8	(9.0-20.6)	35.1	(27.3-43.9)	2.0	(0.9-4.4)*	27.1	(21.4-33.6)	22.0	(15.9-29.6)
	No	3,013	22.4	(20.3-24.6)	33.2	(30.9-35.6)	3.2	(2.6-4.0)	28.9	(26.9-31.1)	12.3	(10.9-13.8)
Serious Psychological Distress	Yes	78	26.1	(14.6-42.1)	24.1	(12.6-41.2)*	0.8	(0.1-5.5)*	24.3	(14.8-37.2)	24.7	(15.1-37.7)
	No	3,192	21.5	(19.6-23.6)	33.6	(31.3-35.9)	3.2	(2.6-4.0)	28.7	(26.8-30.8)	12.9	(11.5-14.5)
Receiving Treatment for Mental Health Condition	Yes	418	16.1	(11.3-22.3)	30.5	(24.8-36.8)	2.4	(1.3-4.3)*	29.9	(24.9-35.3)	21.2	(16.5-26.8)
	No	2,880	22.2	(20.2-24.4)	33.9	(31.5-36.3)	3.3	(2.6-4.1)	28.6	(26.5-30.7)	12.1	(10.7-13.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,181	69.3	(67.2-71.4)	23.7	(21.8-25.8)	2.0	(1.4-2.8)	3.7	(2.7-4.8)	1.3	(0.9-1.8)
Age (years)	18-24	117	67.9	(56.1-77.8)	20.0	(12.4-30.6)	2.9	(0.9-9.1)*	8.7	(4.2-16.9)*	0.5	(0.1-2.3)*
	25-34	476	63.9	(57.8-69.5)	27.9	(22.6-34.0)	1.7	(0.9-3.1)*	5.3	(3.0-9.1)	1.2	(0.5-2.8)*
	35-54	1,890	72.1	(69.3-74.7)	21.2	(18.8-23.8)	2.6	(1.7-3.9)	2.6	(1.7-4.0)	1.6	(0.9-2.8)*
	55+	2,661	70.2	(67.9-72.4)	24.5	(22.4-26.8)	1.3	(0.9-1.9)	2.4	(1.8-3.3)	1.6	(1.1-2.4)
Sex	Men	1,820	63.2	(59.4-66.8)	28.2	(24.8-31.9)	1.9	(1.2-3.2)	5.3	(3.6-7.7)	1.3	(0.8-2.4)
	Women	3,361	75.1	(72.8-77.2)	19.5	(17.5-21.6)	2.1	(1.4-3.1)	2.1	(1.5-3.0)	1.2	(0.8-1.9)
Race/ Ethnicity	Hispanic or Latino	147	58.3	(45.2-70.4)	15.8	(7.2-31.3)*	0.7	(0.1-3.3)*	8.7	(4.0-17.7)*	16.5	(11.3-23.5)
	Not Hispanic or Latino:	3,716	70.2	(68.0-72.4)	23.6	(21.6-25.7)	2.0	(1.4-2.8)	3.1	(2.4-4.1)	1.0	(0.8-1.4)
	White											
	Black or African American	1,091	64.0	(58.6-69.0)	24.4	(20.1-29.4)	3.3	(2.1-5.2)	6.3	(4.1-9.6)	2.0	(1.3-3.2)
	All Other Races	165	63.3	(53.1-72.3)	25.3	(16.6-36.4)	2.4	(1.0-5.5)*	5.8	(2.4-13.7)*	3.2	(1.5-7.0)*
Education	<High School	531	67.7	(60.3-74.3)	22.0	(16.7-28.6)	0.7	(0.3-1.9)*	6.2	(3.2-11.6)*	3.4	(1.9-6.0)
	High School Graduate (or GED)	1,485	65.6	(61.1-69.9)	24.9	(21.0-29.3)	3.3	(2.0-5.4)	4.6	(2.8-7.4)	1.5	(0.8-3.0)*
	Some College	1,358	69.2	(65.1-73.0)	24.2	(20.6-28.2)	2.0	(1.2-3.5)	3.9	(2.4-6.3)	0.6	(0.3-1.4)*
	College Graduate	1,790	73.2	(69.8-76.3)	23.2	(20.1-26.5)	1.5	(0.8-2.9)*	1.6	(0.8-2.9)*	0.5	(0.2-1.2)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	844	69.9	(64.2-75.0)	20.2	(16.0-25.2)	2.4	(1.3-4.6)*	5.7	(3.3-9.6)	1.8	(0.9-3.6)*
	\$20,000-\$34,999	959	63.9	(58.3-69.2)	27.2	(22.1-32.9)	2.6	(1.3-5.0)*	4.8	(2.9-7.9)	1.5	(0.7-3.0)*
	\$35,000-\$49,999	657	71.9	(65.3-77.6)	21.7	(16.9-27.4)	1.6	(0.8-3.3)*	4.0	(1.6-9.9)*	0.8	(0.3-2.3)*
	\$50,000-\$74,999	765	70.6	(65.0-75.6)	26.2	(21.3-31.8)	1.0	(0.5-2.0)*	1.9	(0.7-4.6)*	0.4	(0.1-1.0)*
	\$75,000 and more	1,300	73.8	(69.9-77.4)	22.1	(18.7-25.8)	0.9	(0.5-1.5)	2.2	(1.1-4.6)*	1.0	(0.4-2.9)*
Employment Status	Employed	2,646	69.8	(67.0-72.6)	22.7	(20.3-25.4)	2.2	(1.4-3.4)	3.8	(2.6-5.5)	1.4	(0.8-2.4)
	Out of work	307	69.0	(60.3-76.5)	21.6	(14.8-30.4)	2.6	(1.2-5.3)*	6.7	(3.3-13.5)*	0.1	(0.0-0.5)*
	Homemaker	383	68.0	(59.6-75.4)	28.5	(20.8-37.8)	1.1	(0.3-4.3)*	0.9	(0.2-3.4)*	1.5	(0.4-5.9)*
	Student	77	74.6	(59.3-85.6)	22.3	(12.0-37.6)	1.3	(0.4-4.3)*	1.3	(0.3-5.0)*	0.4	(0.0-2.9)*
	Retired	1,388	66.8	(63.3-70.1)	27.8	(24.5-31.2)	1.7	(1.0-2.8)	2.9	(1.9-4.6)	0.8	(0.4-1.6)*
	Unable to work	367	69.2	(62.6-75.1)	23.4	(18.1-29.7)	1.2	(0.5-3.0)*	2.9	(1.4-5.6)*	3.4	(1.8-6.2)*
Veteran Status	Yes	757	68.7	(63.0-74.0)	24.9	(20.2-30.3)	1.9	(1.1-3.2)	3.0	(1.8-4.9)	1.5	(0.7-3.4)*
	No	4,422	69.4	(67.0-71.6)	23.6	(21.4-25.8)	2.0	(1.4-3.0)	3.8	(2.8-5.2)	1.2	(0.8-1.9)
Diabetes	Yes	717	73.3	(67.6-78.4)	21.5	(17.1-26.7)	1.8	(0.8-3.8)*	2.5	(1.3-4.5)*	0.9	(0.4-2.3)*
	No	4,458	68.9	(66.6-71.1)	24.0	(21.9-26.2)	2.1	(1.4-2.9)	3.8	(2.8-5.1)	1.3	(0.9-2.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	595	73.5 (66.6-79.3)	20.4 (15.4-26.4)	1.6 (0.7-3.5)*	2.3 (1.2-4.2)*	2.3 (1.0-5.5)*				
	No	4,536	69.2 (66.9-71.3)	23.8 (21.8-26.0)	2.1 (1.4-2.9)	3.8 (2.8-5.1)	1.2 (0.8-1.7)				
Asthma	Yes	567	73.8 (67.4-79.3)	20.1 (15.3-26.0)	2.1 (1.0-4.4)*	2.9 (1.2-6.7)*	1.1 (0.4-3.1)*				
	No	4,599	68.5 (66.2-70.7)	24.4 (22.3-26.6)	2.0 (1.4-2.9)	3.8 (2.8-5.1)	1.3 (0.9-1.9)				
Arthritis	Yes	1,852	68.5 (65.1-71.8)	25.2 (21.8-29.0)	2.1 (1.2-3.4)	2.9 (1.9-4.5)	1.3 (0.8-2.3)				
	No	3,309	69.5 (66.9-72.0)	23.3 (21.0-25.7)	2.0 (1.4-2.9)	3.9 (2.8-5.4)	1.3 (0.8-2.0)				
Frequent Mental Distress	Yes	491	67.6 (61.5-73.2)	24.2 (19.1-30.2)	0.8 (0.4-1.7)*	4.3 (2.0-8.9)*	3.1 (1.4-6.5)*				
	No	4,595	69.6 (67.3-71.8)	23.6 (21.5-25.8)	2.2 (1.5-3.1)	3.7 (2.7-5.0)	1.0 (0.7-1.6)				
Serious Psychological Distress	Yes	187	61.5 (50.9-71.1)	24.7 (16.8-34.6)	1.3 (0.4-4.5)*	8.4 (3.5-18.7)*	4.2 (1.9-9.2)*				
	No	4,846	69.8 (67.5-71.9)	23.6 (21.6-25.8)	1.9 (1.4-2.6)	3.5 (2.6-4.8)	1.2 (0.8-1.7)				
Receiving Treatment for Mental Health Condition	Yes	578	77.3 (72.9-81.3)	17.0 (13.5-21.1)	0.8 (0.3-2.1)*	2.6 (1.5-4.6)	2.2 (1.0-4.9)*				
	No	4,588	68.6 (66.3-70.9)	24.3 (22.2-26.5)	2.1 (1.5-3.0)	3.7 (2.8-5.0)	1.2 (0.8-1.8)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,122	24.6	(22.7-26.5)	33.7	(31.7-35.8)	4.7	(3.7-5.9)	23.5	(21.8-25.4)	13.5	(11.9-15.2)
Age (years)	18-24	115	19.0	(11.2-30.3)	37.0	(26.7-48.6)	3.9	(1.2-11.8)*	27.2	(17.8-39.2)	12.9	(5.8-26.3)*
	25-34	470	20.2	(15.8-25.5)	32.1	(27.0-37.8)	6.7	(3.9-11.2)	25.5	(20.6-31.2)	15.5	(11.2-20.9)
	35-54	1,889	24.6	(22.1-27.3)	34.2	(31.4-37.1)	3.9	(2.9-5.2)	23.6	(21.3-26.0)	13.7	(11.8-15.9)
	55+	2,613	29.1	(26.9-31.4)	33.3	(30.9-35.8)	4.3	(3.4-5.5)	21.4	(19.5-23.4)	11.9	(10.4-13.6)
Sex	Men	1,810	26.8	(23.6-30.1)	36.2	(32.8-39.8)	4.6	(3.0-6.8)	22.3	(19.3-25.5)	10.2	(7.8-13.2)
	Women	3,312	22.6	(20.6-24.7)	31.4	(29.1-33.7)	4.8	(3.6-6.2)	24.8	(22.8-27.0)	16.5	(14.6-18.6)
Race/ Ethnicity	Hispanic or Latino	147	20.8	(12.4-32.8)	15.6	(8.7-26.2)	28.0	(19.4-38.6)	12.1	(5.3-25.1)*	23.6	(12.4-40.2)*
	Not Hispanic or Latino:	3,672	23.0	(21.2-24.9)	35.1	(32.8-37.4)	4.5	(3.7-5.5)	25.3	(23.4-27.3)	12.1	(10.7-13.8)
	White											
	Black or African American	1,082	27.8	(23.3-32.8)	30.6	(26.1-35.5)	5.9	(3.8-8.9)	19.5	(15.6-24.1)	16.3	(12.7-20.6)
	All Other Races	164	31.3	(22.1-42.1)	29.0	(20.1-39.9)	2.6	(0.9-7.2)*	18.6	(10.9-29.8)	18.6	(12.5-26.7)
Education	<High School	525	31.7	(25.8-38.2)	34.4	(27.7-41.8)	8.0	(3.5-17.4)*	12.7	(8.3-19.1)	13.2	(8.6-19.7)
	High School Graduate (or GED)	1,475	31.9	(27.8-36.3)	31.0	(27.1-35.2)	4.0	(2.6-6.2)	18.9	(15.9-22.3)	14.1	(10.4-18.8)
	Some College	1,336	20.9	(17.9-24.3)	35.2	(31.3-39.3)	3.5	(2.5-4.9)	26.5	(22.5-30.8)	13.9	(11.5-16.6)
	College Graduate	1,771	19.4	(16.7-22.5)	34.7	(31.5-38.2)	5.1	(3.8-7.0)	27.6	(24.7-30.8)	13.1	(10.9-15.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	834	27.4 (22.7-32.7)	32.6 (26.9-38.7)	7.1 (3.5-13.5)*	18.9 (14.2-24.7)	14.1 (10.5-18.6)				
	\$20,000-\$34,999	948	26.9 (22.1-32.4)	31.5 (26.8-36.6)	3.8 (2.3-6.2)	21.0 (17.0-25.7)	16.8 (12.2-22.7)				
	\$35,000-\$49,999	647	24.6 (19.7-30.2)	28.5 (23.2-34.4)	2.9 (1.7-4.7)	27.8 (22.2-34.2)	16.3 (10.8-23.8)				
	\$50,000-\$74,999	756	22.4 (18.5-26.7)	38.8 (33.5-44.3)	5.1 (2.4-10.5)*	22.9 (19.1-27.1)	10.9 (8.0-14.8)				
	\$75,000 and more	1,294	22.8 (19.4-26.7)	35.9 (32.2-39.8)	4.0 (2.8-5.7)	24.8 (21.7-28.1)	12.5 (10.3-15.0)				
Employment Status	Employed	2,636	23.2 (20.8-25.8)	34.5 (31.8-37.3)	5.5 (4.0-7.6)	23.2 (20.9-25.7)	13.6 (11.6-15.8)				
	Out of work	301	28.1 (20.6-37.0)	33.4 (25.4-42.4)	2.3 (1.1-4.7)*	24.3 (17.4-32.9)	12.0 (7.1-19.5)				
	Homemaker	376	23.5 (17.3-31.1)	32.6 (26.3-39.6)	2.8 (1.4-5.7)*	23.8 (19.1-29.4)	17.3 (9.7-29.0)				
	Student	75	26.0 (13.2-44.9)*	25.6 (15.1-39.9)	4.7 (1.3-16.0)*	34.8 (20.5-52.5)	8.9 (3.9-18.9)*				
	Retired	1,363	28.4 (25.4-31.7)	35.2 (31.8-38.8)	4.1 (2.9-5.7)	21.4 (18.6-24.4)	10.9 (8.8-13.4)				
	Unable to work	360	21.7 (17.1-27.2)	28.5 (22.4-35.6)	2.8 (1.4-5.5)*	26.1 (19.9-33.4)	20.8 (15.2-27.9)				
Veteran Status	Yes	755	23.2 (19.1-28.0)	32.5 (27.7-37.7)	3.2 (2.1-4.8)	23.4 (19.2-28.1)	17.7 (12.0-25.4)				
	No	4,365	24.8 (22.7-26.9)	33.9 (31.7-36.2)	4.9 (3.8-6.3)	23.5 (21.6-25.6)	12.9 (11.3-14.7)				
Diabetes	Yes	707	27.5 (22.1-33.7)	36.4 (30.6-42.5)	2.9 (1.7-4.9)	19.5 (15.9-23.6)	13.7 (10.6-17.6)				
	No	4,409	24.3 (22.3-26.3)	33.6 (31.4-35.8)	4.8 (3.8-6.2)	23.9 (22.0-25.9)	13.4 (11.7-15.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Georgia, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	590	26.9	(20.7-34.1)	31.5	(25.8-37.8)	2.5	(1.3-4.7)*	23.0	(18.2-28.5)	16.1	(12.0-21.3)
	No	4,480	24.3	(22.4-26.4)	34.0	(31.8-36.2)	4.9	(3.8-6.2)	23.6	(21.7-25.6)	13.3	(11.6-15.1)
Asthma	Yes	549	25.3	(19.4-32.2)	32.4	(26.5-38.9)	5.0	(2.8-8.9)	19.0	(14.8-24.2)	18.2	(14.1-23.2)
	No	4,558	24.6	(22.6-26.6)	33.8	(31.6-36.0)	4.6	(3.6-6.0)	24.2	(22.3-26.3)	12.8	(11.1-14.7)
Arthritis	Yes	1,821	23.0	(20.5-25.8)	34.6	(31.3-38.1)	3.3	(2.4-4.5)	22.9	(20.2-25.7)	16.2	(12.9-20.2)
	No	3,281	25.1	(22.8-27.5)	33.7	(31.3-36.2)	5.1	(3.9-6.7)	23.7	(21.5-26.0)	12.4	(10.7-14.4)
Frequent Mental Distress	Yes	482	18.3	(14.1-23.5)	31.7	(25.3-38.8)	3.1	(1.8-5.5)	24.9	(19.2-31.6)	22.0	(17.7-27.0)
	No	4,545	25.2	(23.2-27.3)	33.8	(31.6-36.0)	4.9	(3.8-6.2)	23.4	(21.5-25.4)	12.8	(11.1-14.7)
Serious Psychological Distress	Yes	180	16.9	(11.2-24.7)	23.7	(14.9-35.4)	0.5	(0.1-2.0)*	30.5	(19.5-44.5)	28.4	(20.1-38.6)
	No	4,800	24.7	(22.8-26.7)	34.1	(32.0-36.3)	4.8	(3.8-6.2)	23.5	(21.6-25.4)	12.9	(11.3-14.7)
Receiving Treatment for Mental Health Condition	Yes	564	16.0	(12.5-20.3)	27.8	(22.5-33.7)	4.5	(2.6-7.7)	21.7	(17.3-26.8)	30.0	(24.6-36.0)
	No	4,541	25.4	(23.4-27.5)	33.7	(31.5-35.9)	5.2	(4.2-6.5)	23.4	(21.5-25.4)	12.3	(10.7-14.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,026	66.0	(63.1-68.7)	27.5	(24.8-30.3)	1.8	(1.1-2.8)	3.4	(2.5-4.5)	1.4	(0.9-2.2)
Age (years)	18-24	246	55.9	(47.7-63.8)	37.3	(29.8-45.5)	1.2	(0.4-3.6)*	4.6	(2.4-8.5)*	1.0	(0.4-2.7)*
	25-34	522	59.6	(53.6-65.2)	33.9	(28.5-39.7)	1.8	(0.6-5.4)*	3.3	(1.9-5.6)	1.4	(0.6-3.4)*
	35-54	2,078	65.6	(62.0-69.1)	27.5	(24.3-31.0)	1.9	(1.1-3.4)	3.1	(2.0-4.6)	1.8	(1.1-3.0)
	55+	3,149	72.6	(69.7-75.3)	21.0	(18.5-23.6)	2.0	(1.2-3.3)	3.1	(2.2-4.3)	1.4	(0.9-2.2)
Sex	Men	2,500	61.1	(57.4-64.6)	31.1	(27.7-34.7)	1.9	(1.1-3.5)*	3.8	(2.6-5.4)	2.1	(1.3-3.4)
	Women	3,526	70.5	(67.4-73.5)	24.0	(21.2-27.1)	1.6	(1.0-2.7)	3.0	(2.1-4.3)	0.8	(0.5-1.4)
Race/ Ethnicity	Hispanic or Latino	392	61.7	(54.5-68.4)	25.6	(19.9-32.3)	3.6	(1.4-9.1)*	4.5	(2.6-7.5)	4.6	(2.6-8.2)
	Not Hispanic or Latino:	2,079	68.9	(65.9-71.8)	24.0	(21.3-26.9)	3.2	(2.2-4.8)	2.8	(1.9-4.0)	1.1	(0.7-1.5)
	White											
	Black or African American	37	--	--	--	--	--	--	--	--	--	--
	All Other Races	3,466	58.4	(56.0-60.8)	32.5	(30.2-34.8)	1.4	(1.0-2.0)	4.4	(3.5-5.4)	3.3	(2.6-4.2)
Education	<High School	258	56.3	(47.0-65.1)	32.7	(24.6-42.0)	1.7	(0.5-5.4)*	6.4	(3.6-11.2)	2.9	(1.4-6.0)*
	High School Graduate (or GED)	1,519	63.2	(58.8-67.4)	30.7	(26.7-35.0)	1.2	(0.7-2.2)*	3.3	(2.2-5.0)	1.5	(0.9-2.7)
	Some College	1,718	64.8	(60.6-68.8)	28.0	(24.2-32.1)	2.3	(1.1-4.7)*	3.7	(2.4-5.7)	1.2	(0.6-2.1)*
	College Graduate	2,527	71.2	(67.7-74.4)	23.3	(20.3-26.6)	2.0	(1.0-4.0)*	2.4	(1.6-3.8)	1.1	(0.6-2.2)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	748	61.7	(55.7-67.4)	28.9	(23.7-34.6)	1.5	(0.6-3.5)*	5.7	(3.6-8.9)	2.2	(1.1-4.2)*
	\$20,000-\$34,999	1,037	68.5	(63.6-73.1)	25.7	(21.4-30.5)	1.2	(0.6-2.6)*	3.2	(1.8-5.6)	1.3	(0.7-2.4)
	\$35,000-\$49,999	1,001	66.6	(61.7-71.1)	29.4	(25.0-34.2)	1.0	(0.4-2.2)*	1.8	(1.0-3.2)	1.2	(0.6-2.3)*
	\$50,000-\$74,999	1,044	68.9	(64.1-73.3)	24.9	(20.7-29.5)	2.1	(1.1-4.0)*	3.0	(1.8-5.0)	1.1	(0.5-2.6)*
	\$75,000 and more	1,755	67.6	(63.7-71.2)	26.7	(23.3-30.5)	1.7	(0.8-3.6)*	2.9	(1.8-4.6)	1.1	(0.5-2.3)*
Employment Status	Employed	3,417	64.9	(61.6-68.1)	28.2	(25.2-31.4)	2.1	(1.2-3.8)*	3.4	(2.4-4.8)	1.4	(0.8-2.3)
	Out of work	333	64.4	(56.2-71.8)	28.2	(21.5-36.2)	1.4	(0.3-6.4)*	3.0	(1.3-7.1)*	2.9	(1.2-7.2)*
	Homemaker	292	68.3	(61.1-74.6)	24.0	(18.2-31.0)	2.4	(0.8-7.1)*	4.2	(1.8-9.6)*	1.0	(0.4-2.9)*
	Student	135	69.7	(59.7-78.1)	27.2	(19.2-37.0)	0.0	(0.0-0.0)	2.5	(0.8-8.1)*	0.6	(0.1-2.9)*
	Retired	1,635	68.6	(64.7-72.3)	24.8	(21.3-28.6)	2.0	(1.0-3.7)*	3.5	(2.2-5.4)	1.2	(0.6-2.2)*
	Unable to work	208	72.4	(60.3-81.9)	22.2	(13.8-33.6)	1.9	(0.6-5.5)*	2.3	(1.1-4.9)*	1.2	(0.4-3.7)*
Veteran Status	Yes	986	70.5	(66.1-74.5)	25.3	(21.5-29.6)	1.3	(0.6-2.8)*	2.1	(1.2-3.5)	0.8	(0.4-1.6)*
	No	5,037	65.0	(61.9-68.0)	27.9	(25.0-30.9)	1.9	(1.1-3.1)	3.7	(2.7-5.0)	1.6	(1.0-2.5)
Diabetes	Yes	577	67.6	(61.6-73.2)	27.3	(22.0-33.3)	2.0	(0.8-5.3)*	1.7	(0.8-3.4)*	1.4	(0.7-2.9)*
	No	5,440	65.8	(62.9-68.6)	27.5	(24.8-30.3)	1.7	(1.1-2.8)	3.5	(2.6-4.7)	1.4	(0.9-2.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	473	68.5 (62.0-74.3)	27.6 (21.9-34.2)	0.8 (0.3-2.3)*	1.9 (0.9-3.9)*	1.2 (0.6-2.4)*				
	No	5,492	65.7 (62.8-68.6)	27.5 (24.8-30.4)	1.8 (1.1-2.9)	3.5 (2.6-4.7)	1.5 (0.9-2.3)				
Asthma	Yes	932	68.2 (63.1-72.9)	25.6 (21.2-30.5)	2.6 (1.1-5.8)*	2.3 (1.2-4.3)*	1.3 (0.6-3.1)*				
	No	5,077	65.5 (62.5-68.4)	27.8 (25.1-30.7)	1.6 (1.0-2.5)	3.6 (2.7-4.9)	1.4 (0.9-2.2)				
Arthritis	Yes	1,730	69.3 (65.5-72.9)	24.3 (20.9-28.0)	1.8 (0.9-3.5)*	2.9 (1.8-4.6)	1.7 (0.9-3.1)*				
	No	4,273	65.2 (62.1-68.2)	28.1 (25.3-31.2)	1.7 (1.1-2.9)	3.5 (2.5-4.8)	1.4 (0.9-2.2)				
Frequent Mental Distress	Yes	489	63.2 (56.4-69.4)	31.2 (25.2-38.0)	0.6 (0.1-2.2)*	2.7 (1.3-5.3)*	2.4 (1.1-5.0)*				
	No	5,508	66.3 (63.4-69.2)	27.1 (24.4-30.0)	1.9 (1.2-3.1)	3.3 (2.5-4.5)	1.3 (0.8-2.1)				
Serious Psychological Distress	Yes	160	59.5 (48.9-69.2)	33.4 (24.3-44.0)	1.3 (0.2-8.9)*	4.1 (1.9-8.6)*	1.7 (0.8-3.9)*				
	No	5,806	66.2 (63.3-69.0)	27.4 (24.7-30.3)	1.7 (1.0-2.7)	3.3 (2.4-4.5)	1.4 (0.9-2.2)				
Receiving Treatment for Mental Health Condition	Yes	491	77.3 (71.2-82.4)	19.3 (14.3-25.4)	0.3 (0.0-2.2)*	1.9 (0.9-3.8)*	1.2 (0.6-2.4)*				
	No	5,529	65.0 (62.0-67.9)	28.3 (25.5-31.2)	1.9 (1.2-3.0)	3.4 (2.5-4.6)	1.5 (0.9-2.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,010	25.2	(22.8-27.8)	39.7	(36.8-42.7)	2.5	(1.8-3.3)	24.1	(21.8-26.6)	8.5	(7.2-10.1)
Age (years)	18-24	247	20.1	(15.0-26.4)	50.9	(42.5-59.2)	1.1	(0.4-3.0)*	21.7	(14.9-30.6)	6.2	(3.5-10.9)
	25-34	524	16.9	(13.2-21.3)	46.4	(40.5-52.4)	1.7	(0.8-3.5)*	27.4	(22.5-33.0)	7.6	(5.1-11.1)
	35-54	2,083	24.1	(21.1-27.4)	37.8	(34.3-41.4)	2.6	(1.8-4.0)	25.9	(22.9-29.2)	9.5	(7.7-11.8)
	55+	3,124	33.3	(30.3-36.4)	33.9	(31.0-36.9)	3.1	(2.2-4.4)	21.2	(18.9-23.7)	8.5	(7.1-10.2)
Sex	Men	2,502	26.4	(23.4-29.6)	42.3	(38.7-46.0)	2.6	(1.8-3.8)	20.7	(18.1-23.5)	8.0	(6.4-10.1)
	Women	3,508	24.2	(21.6-27.0)	37.0	(33.8-40.4)	2.3	(1.6-3.3)	27.5	(24.5-30.7)	9.0	(7.4-10.8)
Race/ Ethnicity	Hispanic or Latino	396	31.6	(25.7-38.2)	33.2	(26.7-40.4)	2.5	(1.1-5.6)*	21.8	(16.2-28.7)	10.9	(7.6-15.4)
	Not Hispanic or Latino:	2,073	24.9	(22.3-27.7)	36.4	(33.5-39.4)	2.4	(1.7-3.3)	24.5	(22.1-27.1)	11.8	(10.0-13.9)
	White											
	Black or African American	37	--	--	--	--	--	--	--	--	--	--
	All Other Races	3,456	31.2	(29.1-33.4)	38.5	(36.2-40.9)	2.2	(1.7-2.8)	19.8	(18.0-21.8)	8.3	(7.1-9.7)
Education	<High School	259	26.8	(20.2-34.6)	35.8	(26.8-45.9)	2.5	(0.7-9.0)*	21.6	(14.5-30.8)	13.4	(8.2-21.1)
	High School Graduate (or GED)	1,527	29.8	(25.9-33.9)	36.5	(32.2-41.0)	2.4	(1.5-3.8)	23.5	(19.8-27.6)	7.9	(6.0-10.4)
	Some College	1,711	23.9	(20.7-27.5)	38.4	(34.2-42.8)	2.1	(1.4-3.3)	26.6	(22.7-30.8)	8.9	(7.0-11.3)
	College Graduate	2,509	20.5	(17.9-23.5)	43.2	(39.6-46.8)	2.9	(2.0-4.2)	24.8	(22.0-27.8)	8.6	(6.9-10.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	738	25.8 (21.2-30.9)	35.5 (29.6-41.8)	2.3 (1.1-5.1)*	25.3 (19.7-31.9)	11.1 (8.0-15.1)				
	\$20,000-\$34,999	1,045	28.7 (24.6-33.3)	39.9 (34.8-45.2)	2.3 (1.2-4.3)*	19.0 (15.7-22.9)	10.1 (7.4-13.7)				
	\$35,000-\$49,999	993	23.4 (19.8-27.5)	41.7 (36.6-46.9)	1.6 (0.8-3.1)*	25.0 (20.7-29.8)	8.4 (6.0-11.5)				
	\$50,000-\$74,999	1,049	27.2 (23.2-31.5)	36.5 (31.7-41.5)	2.6 (1.6-4.0)	25.0 (21.0-29.5)	8.8 (6.6-11.7)				
	\$75,000 and more	1,746	22.2 (19.2-25.6)	42.2 (38.3-46.2)	2.5 (1.7-3.9)	25.7 (22.5-29.2)	7.3 (5.7-9.3)				
Employment Status	Employed	3,409	24.4 (21.7-27.3)	39.0 (35.7-42.3)	2.5 (1.7-3.5)	25.7 (23.0-28.7)	8.4 (6.9-10.3)				
	Out of work	329	20.7 (15.4-27.2)	43.1 (35.0-51.6)	2.8 (0.9-8.1)*	22.6 (16.2-30.7)	10.7 (6.3-17.7)				
	Homemaker	288	31.8 (25.6-38.7)	36.6 (29.5-44.4)	2.9 (1.3-6.4)*	20.9 (15.6-27.3)	7.8 (4.5-13.3)				
	Student	134	15.0 (9.1-23.8)	54.4 (43.0-65.4)	0.7 (0.1-5.1)*	20.5 (12.4-31.9)	9.3 (4.7-17.7)*				
	Retired	1,636	30.4 (26.8-34.3)	39.2 (35.1-43.4)	2.8 (1.8-4.3)	20.3 (17.3-23.6)	7.4 (5.8-9.4)				
	Unable to work	208	21.0 (14.1-30.2)	29.7 (18.7-43.8)	2.6 (0.9-6.8)*	32.8 (18.6-51.0)	14.0 (9.1-20.9)				
Veteran Status	Yes	986	25.8 (22.1-29.9)	39.7 (35.0-44.5)	2.1 (1.3-3.6)	23.2 (19.3-27.6)	9.2 (6.8-12.3)				
	No	5,021	25.0 (22.5-27.8)	39.7 (36.6-42.9)	2.5 (1.9-3.5)	24.3 (21.8-27.0)	8.4 (7.0-10.1)				
Diabetes	Yes	572	27.4 (22.6-32.9)	34.3 (28.4-40.7)	4.8 (2.4-9.3)*	23.7 (18.5-29.9)	9.7 (6.8-13.8)				
	No	5,429	25.0 (22.5-27.6)	40.2 (37.2-43.2)	2.3 (1.7-3.1)	24.1 (21.8-26.7)	8.5 (7.1-10.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Hawaii, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	469	24.1	(19.0-30.1)	36.1	(29.8-43.0)	3.4	(1.4-7.8)*	25.6	(20.3-31.8)	10.8	(7.5-15.4)
	No	5,483	25.4	(22.9-28.0)	39.7	(36.7-42.8)	2.4	(1.8-3.2)	24.1	(21.7-26.6)	8.4	(7.0-10.0)
Asthma	Yes	921	20.9	(17.1-25.3)	38.0	(32.9-43.4)	3.2	(1.9-5.2)	28.8	(24.3-33.9)	9.1	(6.7-12.2)
	No	5,071	26.0	(23.5-28.7)	39.9	(36.9-43.0)	2.3	(1.7-3.2)	23.3	(20.9-25.8)	8.5	(7.1-10.1)
Arthritis	Yes	1,719	23.9	(20.8-27.2)	36.6	(32.8-40.6)	2.7	(1.7-4.3)	26.1	(22.8-29.7)	10.7	(8.6-13.1)
	No	4,268	25.6	(22.9-28.4)	40.5	(37.3-43.7)	2.4	(1.7-3.3)	23.6	(21.1-26.3)	8.0	(6.6-9.7)
Frequent Mental Distress	Yes	495	21.0	(16.3-26.6)	32.5	(26.4-39.3)	1.3	(0.6-3.1)*	27.3	(21.6-33.9)	17.9	(13.1-23.9)
	No	5,486	25.6	(23.1-28.3)	40.3	(37.2-43.4)	2.6	(1.9-3.5)	23.9	(21.5-26.4)	7.7	(6.4-9.2)
Serious Psychological Distress	Yes	163	16.5	(11.0-23.9)	32.5	(22.6-44.2)	1.9	(0.4-7.6)*	30.4	(20.7-42.2)	18.8	(12.5-27.4)
	No	5,787	25.6	(23.1-28.3)	39.8	(36.8-42.8)	2.5	(1.8-3.3)	23.9	(21.6-26.5)	8.2	(6.8-9.7)
Receiving Treatment for Mental Health Condition	Yes	485	19.0	(15.0-23.7)	30.8	(24.9-37.5)	2.0	(1.0-3.8)*	30.1	(24.0-37.1)	18.0	(13.3-23.9)
	No	5,519	26.3	(23.7-29.0)	39.7	(36.7-42.9)	2.5	(1.9-3.3)	23.6	(21.2-26.2)	7.9	(6.6-9.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,947	65.4	(63.6-67.2)	27.9	(26.2-29.7)	0.9	(0.7-1.3)	4.2	(3.4-5.1)	1.5	(1.2-2.1)
Age (years)	18-24	194	52.7	(44.8-60.5)	40.4	(32.8-48.5)	0.7	(0.2-2.7)*	6.2	(3.6-10.7)	0.0	(0.0-0.0)
	25-34	526	61.1	(56.0-65.9)	31.5	(26.9-36.4)	0.5	(0.1-1.5)*	5.6	(3.3-9.3)	1.4	(0.5-3.5)*
	35-54	1,843	68.7	(66.1-71.3)	25.3	(22.9-27.9)	0.6	(0.3-1.2)*	3.5	(2.5-4.8)	1.8	(1.2-2.8)
	55+	2,380	68.7	(66.5-70.9)	24.3	(22.3-26.5)	1.7	(1.2-2.4)	3.2	(2.5-4.1)	2.0	(1.4-2.9)
Sex	Men	1,908	58.9	(55.9-61.8)	33.4	(30.6-36.4)	1.1	(0.7-1.7)	5.1	(3.7-6.9)	1.6	(1.1-2.3)
	Women	3,039	71.6	(69.5-73.6)	22.8	(20.9-24.8)	0.8	(0.5-1.2)	3.3	(2.6-4.3)	1.5	(1.0-2.3)
Race/ Ethnicity	Hispanic or Latino	275	65.5	(58.3-72.0)	27.5	(21.6-34.3)	0.4	(0.1-1.7)*	4.5	(2.2-9.0)*	2.1	(0.9-4.7)*
	Not Hispanic or Latino:	4,033	67.3	(65.5-69.1)	26.5	(24.8-28.3)	1.1	(0.8-1.5)	3.9	(3.1-4.8)	1.2	(0.9-1.7)
	White											
	Black or African American	448	58.6	(53.1-63.9)	33.5	(28.3-39.3)	0.7	(0.2-1.8)*	5.5	(3.5-8.4)	1.7	(0.9-3.1)*
	All Other Races	174	57.1	(48.0-65.7)	33.5	(25.3-42.9)	1.3	(0.3-5.2)*	4.5	(1.9-10.0)*	3.6	(1.8-7.3)*
Education	<High School	322	58.4	(50.5-65.9)	32.8	(25.9-40.6)	1.1	(0.5-2.6)*	5.5	(3.2-9.3)	2.1	(0.9-5.1)*
	High School Graduate (or GED)	1,380	61.6	(58.0-65.0)	30.4	(27.1-33.9)	1.2	(0.8-1.9)	5.4	(3.6-7.9)	1.4	(0.9-2.3)
	Some College	1,383	65.5	(62.1-68.8)	27.8	(24.6-31.1)	1.0	(0.5-1.9)*	4.1	(2.9-5.7)	1.6	(1.0-2.6)
	College Graduate	1,859	70.8	(68.0-73.4)	24.6	(22.1-27.3)	0.7	(0.4-1.1)	2.6	(1.7-3.9)	1.3	(0.7-2.5)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	665	61.7	(56.1-67.0)	32.0	(26.8-37.6)	1.0	(0.5-1.8)*	3.6	(2.3-5.7)	1.8	(0.9-3.3)*
	\$20,000-\$34,999	810	64.7	(59.9-69.1)	26.2	(22.1-30.7)	1.0	(0.5-2.0)*	5.3	(3.1-9.1)	2.8	(1.6-5.0)*
	\$35,000-\$49,999	708	67.2	(62.5-71.7)	25.6	(21.5-30.1)	1.2	(0.6-2.4)*	4.3	(2.7-6.7)	1.7	(0.9-3.2)*
	\$50,000-\$74,999	809	65.1	(61.1-68.9)	28.7	(25.0-32.8)	0.7	(0.2-2.2)*	4.2	(2.7-6.5)	1.2	(0.6-2.5)*
	\$75,000 and more	1,494	70.5	(67.4-73.4)	25.4	(22.6-28.5)	0.6	(0.3-1.2)*	2.9	(1.8-4.6)	0.6	(0.3-1.3)*
Employment Status	Employed	2,736	65.6	(63.3-67.9)	28.1	(25.9-30.4)	0.9	(0.6-1.4)	4.5	(3.4-6.0)	0.9	(0.6-1.4)
	Out of work	181	64.9	(55.6-73.1)	26.2	(19.0-35.0)	0.3	(0.0-2.4)*	5.9	(3.1-10.9)*	2.7	(1.0-7.3)*
	Homemaker	446	68.6	(63.0-73.8)	22.4	(17.7-27.8)	0.6	(0.2-1.6)*	3.7	(1.6-8.5)*	4.7	(2.2-9.4)*
	Student	116	68.0	(57.6-76.8)	30.4	(21.7-40.6)	0.0	(0.0-0.0)	0.7	(0.1-4.8)*	1.0	(0.2-4.6)*
	Retired	1,217	61.8	(58.5-65.0)	30.0	(27.0-33.3)	1.7	(1.1-2.7)	4.1	(2.9-5.7)	2.4	(1.5-3.7)
	Unable to work	249	66.6	(58.2-74.2)	25.9	(19.2-34.0)	1.1	(0.5-2.6)*	4.8	(2.7-8.3)	1.6	(0.7-3.7)*
Veteran Status	Yes	577	64.4	(59.9-68.7)	26.7	(22.7-31.1)	1.3	(0.6-2.7)*	5.3	(3.5-8.0)	2.3	(1.3-4.0)
	No	4,369	65.6	(63.6-67.4)	28.0	(26.2-29.9)	0.9	(0.6-1.2)	4.1	(3.2-5.1)	1.5	(1.1-2.0)
Diabetes	Yes	540	70.2	(65.0-74.9)	22.0	(17.6-27.1)	0.7	(0.3-1.6)*	5.1	(3.2-7.9)	2.0	(1.1-3.8)*
	No	4,406	64.9	(63.0-66.8)	28.5	(26.7-30.3)	1.0	(0.7-1.4)	4.1	(3.2-5.1)	1.5	(1.1-2.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	509	65.7	(60.6-70.5)	26.5	(22.0-31.4)	1.1	(0.5-2.6)*	4.4	(2.6-7.4)	2.2	(1.2-4.2)*
	No	4,417	65.4	(63.5-67.2)	28.1	(26.3-30.0)	0.9	(0.6-1.3)	4.2	(3.3-5.2)	1.5	(1.1-2.0)
Asthma	Yes	589	65.4	(60.0-70.5)	28.7	(23.6-34.4)	1.1	(0.5-2.4)*	3.6	(2.0-6.4)	1.2	(0.6-2.4)*
	No	4,351	65.3	(63.4-67.2)	27.8	(26.0-29.6)	1.0	(0.7-1.3)	4.3	(3.4-5.3)	1.7	(1.2-2.2)
Arthritis	Yes	1,794	67.3	(64.5-70.1)	26.3	(23.8-29.1)	1.0	(0.7-1.6)	3.4	(2.5-4.7)	1.9	(1.2-2.8)
	No	3,146	64.7	(62.5-66.9)	28.5	(26.4-30.7)	0.9	(0.6-1.3)	4.4	(3.5-5.7)	1.5	(1.0-2.1)
Frequent Mental Distress	Yes	415	67.7	(61.7-73.2)	24.4	(19.4-30.2)	0.8	(0.3-1.9)*	4.6	(2.6-8.0)	2.5	(1.3-4.5)*
	No	4,502	65.5	(63.6-67.4)	28.1	(26.3-30.0)	0.9	(0.7-1.3)	4.1	(3.3-5.1)	1.3	(1.0-1.9)
Serious Psychological Distress	Yes	149	68.9	(59.4-77.0)	17.5	(11.4-25.7)	1.3	(0.4-4.2)*	4.8	(1.8-12.0)*	7.6	(4.1-13.7)*
	No	4,763	65.5	(63.6-67.3)	28.0	(26.3-29.8)	1.0	(0.7-1.3)	4.2	(3.4-5.2)	1.4	(1.0-1.9)
Receiving Treatment for Mental Health Condition	Yes	521	75.7	(70.6-80.1)	17.8	(13.8-22.6)	0.6	(0.2-1.7)*	4.3	(2.5-7.4)	1.6	(0.7-3.5)*
	No	4,421	64.4	(62.5-66.3)	28.9	(27.1-30.8)	1.0	(0.7-1.3)	4.1	(3.3-5.2)	1.5	(1.1-2.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,940	23.5	(22.0-25.1)	38.6	(36.9-40.5)	1.1	(0.8-1.6)	26.6	(25.1-28.1)	10.1	(9.1-11.3)
Age (years)												
	18-24	196	15.7	(11.2-21.5)	48.1	(40.2-56.0)	1.4	(0.5-3.9)*	28.1	(21.1-36.3)	6.8	(3.7-12.1)*
	25-34	527	20.9	(17.0-25.4)	40.1	(35.3-45.1)	0.7	(0.3-2.0)*	28.2	(24.3-32.4)	10.1	(7.3-13.9)
	35-54	1,846	22.6	(20.3-25.1)	38.3	(35.7-41.0)	0.9	(0.4-1.8)*	27.8	(25.6-30.1)	10.4	(8.9-12.2)
	55+	2,367	29.6	(27.5-31.8)	34.4	(32.2-36.7)	1.6	(1.2-2.3)	23.5	(21.7-25.5)	10.9	(9.5-12.5)
Sex												
	Men	1,911	25.2	(22.8-27.8)	41.6	(38.7-44.6)	1.4	(0.9-2.2)	23.3	(21.0-25.8)	8.4	(6.8-10.3)
	Women	3,029	21.9	(20.1-23.8)	35.8	(33.7-38.0)	0.9	(0.6-1.4)	29.7	(27.8-31.7)	11.7	(10.4-13.2)
Race/Ethnicity												
	Hispanic or Latino	273	34.1	(27.9-40.9)	36.2	(29.2-43.8)	1.0	(0.3-3.0)*	15.5	(11.1-21.1)	13.2	(8.7-19.6)
	Not Hispanic or Latino:	4,027	19.8	(18.4-21.2)	40.7	(38.9-42.6)	1.3	(0.9-1.9)	28.7	(27.1-30.5)	9.4	(8.4-10.5)
	White											
	Black or African American	449	22.7	(18.5-27.6)	33.7	(28.2-39.6)	0.9	(0.4-1.8)*	29.1	(23.8-35.0)	13.7	(10.5-17.6)
	All Other Races	174	39.2	(30.7-48.3)	32.6	(24.9-41.3)	0.5	(0.1-2.1)*	21.9	(15.5-30.1)	5.9	(3.4-10.1)
Education												
	<High School	319	36.3	(29.4-43.7)	34.5	(27.5-42.3)	1.8	(0.4-6.9)*	17.4	(12.8-23.1)	10.0	(5.6-17.3)
	High School Graduate (or GED)	1,374	30.3	(27.1-33.6)	36.4	(32.8-40.1)	1.0	(0.5-1.8)*	23.8	(21.0-26.9)	8.6	(6.9-10.5)
	Some College	1,389	19.6	(17.0-22.5)	39.1	(35.7-42.5)	1.5	(0.9-2.5)	28.1	(25.1-31.3)	11.8	(9.8-14.1)
	College Graduate	1,855	16.3	(14.3-18.6)	41.8	(39.0-44.6)	0.9	(0.6-1.5)	30.4	(27.9-33.1)	10.6	(8.9-12.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	660	25.9	(21.7-30.7)	31.7	(26.5-37.4)	2.7	(1.3-5.6)*	26.4	(21.7-31.6)	13.3	(9.9-17.8)
	\$20,000-\$34,999	804	27.6	(23.7-31.9)	39.2	(34.4-44.2)	1.1	(0.5-2.4)*	21.7	(18.4-25.6)	10.3	(7.8-13.6)
	\$35,000-\$49,999	710	23.9	(20.0-28.3)	36.9	(32.4-41.6)	0.6	(0.2-1.5)*	28.4	(24.5-32.5)	10.2	(7.6-13.6)
	\$50,000-\$74,999	812	19.2	(16.1-22.7)	40.7	(36.7-44.8)	1.1	(0.5-2.4)*	31.0	(27.4-34.8)	8.0	(6.2-10.3)
	\$75,000 and more	1,493	22.4	(19.7-25.3)	40.6	(37.6-43.7)	0.7	(0.4-1.2)*	26.2	(23.8-28.9)	10.1	(8.4-12.2)
Employment Status	Employed	2,734	22.9	(20.8-25.1)	40.5	(38.1-42.8)	1.1	(0.8-1.6)	26.2	(24.3-28.2)	9.4	(8.1-10.7)
	Out of work	182	21.0	(14.7-28.9)	30.7	(22.8-40.0)	3.1	(0.8-11.2)*	28.7	(21.4-37.4)	16.5	(9.4-27.3)
	Homemaker	444	31.9	(26.8-37.5)	31.9	(27.0-37.2)	1.0	(0.5-1.9)*	24.9	(20.6-29.7)	10.3	(7.2-14.7)
	Student	116	17.9	(11.1-27.5)	42.6	(32.5-53.3)	0.4	(0.3-0.6)	29.1	(19.9-40.6)	10.0	(5.1-18.5)*
	Retired	1,214	25.6	(23.0-28.4)	37.6	(34.4-41.0)	1.6	(1.0-2.5)	26.4	(23.5-29.5)	8.8	(7.1-10.7)
	Unable to work	248	19.3	(14.1-26.0)	31.1	(23.0-40.7)	1.5	(0.5-4.0)*	30.9	(23.8-39.1)	17.1	(11.8-24.2)
Veteran Status	Yes	577	23.4	(19.9-27.3)	34.1	(29.8-38.7)	1.5	(0.8-2.9)*	27.6	(23.4-32.2)	13.3	(9.8-17.9)
	No	4,362	23.5	(21.8-25.2)	39.1	(37.1-41.0)	1.2	(0.9-1.7)	26.4	(24.8-28.1)	9.8	(8.8-11.0)
Diabetes	Yes	533	25.1	(20.8-30.0)	36.0	(31.1-41.1)	1.4	(0.7-2.9)*	26.2	(22.0-30.9)	11.3	(8.6-14.8)
	No	4,406	23.3	(21.7-25.0)	38.9	(37.0-40.8)	1.2	(0.9-1.7)	26.6	(25.0-28.2)	10.0	(8.9-11.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Illinois, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics		Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	510	26.0	(21.8-30.7)	36.1	(31.0-41.5)	1.1	(0.4-2.5)*	24.4	(20.1-29.2)	12.5	(9.2-16.6)
	No	4,409	23.3	(21.7-25.0)	39.0	(37.1-40.9)	1.1	(0.8-1.6)	26.7	(25.1-28.4)	9.9	(8.8-11.1)
Asthma	Yes	585	22.0	(18.0-26.5)	40.8	(35.5-46.3)	1.3	(0.5-3.3)*	24.6	(20.6-29.1)	11.2	(8.5-14.7)
	No	4,349	23.7	(22.0-25.4)	38.4	(36.5-40.3)	1.1	(0.8-1.6)	26.9	(25.2-28.6)	10.0	(8.8-11.2)
Arthritis	Yes	1,790	22.9	(20.7-25.2)	34.1	(31.4-36.9)	1.3	(0.8-2.1)	30.1	(27.5-32.8)	11.7	(9.9-13.6)
	No	3,143	23.7	(21.8-25.7)	40.3	(38.0-42.5)	1.1	(0.7-1.6)	25.4	(23.5-27.3)	9.6	(8.3-11.0)
Frequent Mental Distress	Yes	411	15.7	(12.0-20.2)	34.2	(28.4-40.5)	0.7	(0.3-1.9)*	28.5	(23.6-33.9)	21.0	(16.4-26.3)
	No	4,504	24.2	(22.6-25.9)	39.0	(37.2-40.9)	1.2	(0.9-1.7)	26.4	(24.8-28.1)	9.1	(8.0-10.2)
Serious Psychological Distress	Yes	150	11.5	(7.2-17.9)	21.3	(14.1-30.9)	0.5	(0.1-1.9)*	35.5	(26.1-46.3)	31.1	(22.0-42.0)
	No	4,756	23.8	(22.3-25.5)	39.1	(37.3-41.0)	1.2	(0.9-1.6)	26.4	(24.9-28.0)	9.5	(8.4-10.6)
Receiving Treatment for Mental Health Condition	Yes	517	20.2	(16.3-24.9)	32.2	(27.3-37.5)	1.2	(0.5-2.8)*	29.6	(25.1-34.5)	16.8	(13.1-21.4)
	No	4,416	23.9	(22.3-25.6)	39.3	(37.4-41.3)	1.1	(0.8-1.6)	26.2	(24.6-27.9)	9.4	(8.4-10.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,254	66.7	(64.9-68.5)	26.8	(25.1-28.6)	1.3	(1.0-1.7)	3.6	(2.9-4.4)	1.6	(1.1-2.3)
Age (years)	18-24	209	56.1	(47.7-64.2)	35.9	(28.4-44.1)	0.6	(0.2-2.2)*	5.1	(2.6-9.8)*	2.4	(0.8-6.4)*
	25-34	601	60.3	(55.5-65.0)	32.3	(27.7-37.2)	1.1	(0.5-2.3)*	4.3	(2.7-6.7)	2.0	(1.0-4.1)*
	35-54	2,009	70.8	(68.2-73.2)	23.5	(21.2-26.0)	1.3	(0.8-2.1)	3.1	(2.2-4.3)	1.4	(0.8-2.4)
	55+	2,398	69.8	(67.5-72.0)	24.0	(21.9-26.2)	1.7	(1.1-2.4)	3.2	(2.4-4.3)	1.3	(0.8-2.1)
Sex	Men	1,923	60.7	(57.7-63.7)	31.5	(28.7-34.5)	1.3	(0.9-1.9)	4.5	(3.4-6.0)	2.0	(1.2-3.3)
	Women	3,331	72.3	(70.1-74.3)	22.5	(20.6-24.5)	1.2	(0.8-1.8)	2.7	(2.0-3.7)	1.3	(0.8-2.0)
Race/Ethnicity	Hispanic or Latino	154	61.0	(47.4-73.0)	33.2	(22.3-46.2)	1.2	(0.3-4.8)*	3.7	(1.6-8.5)*	0.9	(0.2-4.6)*
	Not Hispanic or Latino:	4,346	67.9	(66.0-69.7)	25.9	(24.2-27.7)	1.2	(0.9-1.6)	3.4	(2.7-4.3)	1.5	(1.0-2.2)
	White											
	Black or African American	579	62.8	(55.2-69.8)	32.0	(25.0-39.9)	0.7	(0.2-1.8)*	3.0	(1.6-5.7)*	1.5	(0.7-3.2)*
	All Other Races	133	64.8	(53.6-74.5)	27.8	(18.9-38.8)	3.8	(1.4-9.5)*	3.6	(1.5-8.4)*	0.1	(0.0-0.5)*
Education	<High School	497	56.9	(50.3-63.3)	32.5	(26.2-39.4)	1.7	(0.7-3.7)*	4.1	(2.5-6.6)	4.8	(3.1-7.6)
	High School Graduate (or GED)	1,929	62.4	(59.0-65.7)	29.7	(26.7-32.9)	1.7	(1.1-2.5)	4.8	(3.5-6.5)	1.5	(0.7-2.9)*
	Some College	1,308	67.8	(64.1-71.2)	26.1	(22.9-29.6)	1.0	(0.5-1.7)	3.9	(2.4-6.2)	1.3	(0.7-2.5)*
	College Graduate	1,512	72.5	(69.5-75.4)	23.5	(20.7-26.4)	1.0	(0.6-1.7)	1.8	(1.2-2.8)	1.2	(0.5-2.8)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	890	64.7 (58.7-70.2)	28.8 (23.7-34.4)	0.6 (0.3-1.2)*	3.4 (2.2-5.2)	2.6 (1.4-4.8)*				
	\$20,000-\$34,999	1,039	64.6 (60.4-68.7)	27.8 (24.0-31.9)	0.9 (0.4-1.8)*	4.7 (3.0-7.2)	2.0 (1.0-4.1)*				
	\$35,000-\$49,999	780	62.9 (58.4-67.2)	31.6 (27.4-36.1)	1.4 (0.7-2.9)*	3.0 (1.7-5.4)	1.1 (0.5-2.4)*				
	\$50,000-\$74,999	862	68.9 (64.6-72.9)	24.3 (20.7-28.4)	1.3 (0.7-2.4)*	3.4 (2.0-5.8)	2.1 (0.8-5.7)*				
	\$75,000 and more	1,040	72.4 (68.6-76.0)	22.5 (19.1-26.2)	1.2 (0.6-2.3)*	3.2 (1.8-5.7)	0.7 (0.3-1.7)*				
Employment Status	Employed	2,851	67.1 (64.6-69.5)	26.6 (24.4-29.0)	1.2 (0.8-1.7)	3.6 (2.7-4.9)	1.5 (0.9-2.5)				
	Out of work	257	67.2 (58.0-75.2)	28.4 (20.7-37.5)	0.2 (0.0-1.4)*	2.3 (1.1-4.6)*	2.0 (0.6-6.8)*				
	Homemaker	441	64.3 (58.2-70.0)	28.0 (22.6-34.1)	3.0 (1.5-5.7)*	2.7 (1.3-5.5)*	2.0 (0.7-5.7)*				
	Student	100	72.7 (61.3-81.8)	22.4 (14.2-33.5)	0.6 (0.1-4.5)*	3.9 (1.3-11.2)*	0.3 (0.0-2.2)*				
	Retired	1,233	66.4 (63.0-69.6)	27.3 (24.2-30.6)	1.7 (1.0-2.8)	3.6 (2.5-5.3)	1.0 (0.5-2.0)*				
	Unable to work	354	61.6 (54.3-68.4)	29.3 (22.8-36.7)	0.0 (0.0-0.2)*	4.7 (2.5-8.7)*	4.4 (2.2-8.6)*				
Veteran Status	Yes	627	64.7 (59.8-69.3)	28.1 (23.7-32.9)	1.7 (0.9-3.0)*	4.0 (2.4-6.7)	1.6 (0.7-3.6)*				
	No	4,625	67.0 (65.0-68.9)	26.7 (24.8-28.6)	1.2 (0.9-1.6)	3.5 (2.8-4.5)	1.6 (1.1-2.4)				
Diabetes	Yes	598	68.4 (63.4-72.9)	26.8 (22.4-31.7)	1.1 (0.5-2.2)*	2.7 (1.5-4.9)*	1.0 (0.4-2.3)*				
	No	4,648	66.5 (64.5-68.4)	26.8 (25.0-28.7)	1.3 (1.0-1.7)	3.7 (2.9-4.6)	1.7 (1.2-2.5)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	605	69.9 (63.8-75.4)	22.8 (18.4-28.0)	1.5 (0.7-3.2)*	3.9 (2.1-7.1)*	1.9 (1.0-3.5)*				
	No	4,590	66.3 (64.4-68.2)	27.3 (25.4-29.2)	1.2 (0.9-1.7)	3.6 (2.8-4.5)	1.6 (1.1-2.4)				
Asthma	Yes	686	66.9 (61.9-71.6)	26.4 (21.9-31.4)	0.7 (0.3-1.3)*	2.5 (1.5-4.3)	3.5 (1.6-7.3)*				
	No	4,555	66.7 (64.7-68.6)	26.7 (24.9-28.6)	1.4 (1.1-1.8)	3.8 (3.0-4.7)	1.4 (1.0-2.1)				
Arthritis	Yes	2,029	68.3 (65.4-71.0)	24.9 (22.4-27.6)	1.2 (0.8-2.0)	4.0 (2.8-5.9)	1.5 (0.9-2.4)				
	No	3,207	66.1 (63.8-68.4)	27.5 (25.4-29.8)	1.3 (0.9-1.7)	3.4 (2.6-4.4)	1.7 (1.1-2.6)				
Frequent Mental Distress	Yes	598	64.8 (58.9-70.3)	27.0 (22.1-32.5)	0.6 (0.1-2.6)*	4.6 (2.7-7.8)	3.0 (1.6-5.9)*				
	No	4,575	67.2 (65.3-69.1)	26.4 (24.6-28.3)	1.4 (1.0-1.8)	3.5 (2.8-4.4)	1.5 (1.0-2.2)				
Serious Psychological Distress	Yes	232	51.8 (43.4-60.1)	28.7 (21.8-36.8)	0.1 (0.1-0.2)	6.7 (3.1-14.0)*	12.6 (8.3-18.7)				
	No	4,929	67.3 (65.4-69.2)	26.6 (24.8-28.4)	1.3 (1.0-1.7)	3.4 (2.7-4.3)	1.4 (0.9-2.0)				
Receiving Treatment for Mental Health Condition	Yes	722	73.2 (68.9-77.1)	21.0 (17.5-25.0)	0.1 (0.0-0.3)*	3.4 (2.0-5.9)	2.2 (1.0-5.0)*				
	No	4,518	66.0 (63.9-67.9)	27.5 (25.7-29.5)	1.4 (1.1-1.9)	3.6 (2.8-4.5)	1.5 (1.0-2.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,262	23.9	(22.3-25.6)	38.1	(36.3-40.0)	2.0	(1.6-2.6)	25.5	(23.9-27.1)	10.4	(9.4-11.5)
Age (years)	18-24	209	24.9	(18.8-32.2)	46.7	(38.5-55.2)	0.7	(0.2-2.4)*	21.2	(15.3-28.6)	6.5	(3.7-11.3)
	25-34	609	18.9	(15.0-23.5)	41.1	(36.4-46.0)	1.2	(0.6-2.5)*	30.2	(25.9-34.9)	8.6	(6.4-11.6)
	35-54	2,006	22.4	(20.2-24.9)	36.1	(33.6-38.8)	2.7	(1.8-4.0)	26.7	(24.4-29.1)	12.0	(10.4-13.9)
	55+	2,401	28.1	(25.9-30.4)	35.8	(33.4-38.2)	2.3	(1.7-3.1)	22.9	(20.9-25.0)	10.9	(9.5-12.6)
Sex	Men	1,942	25.9	(23.3-28.7)	40.8	(37.8-43.8)	2.2	(1.5-3.1)	23.0	(20.6-25.5)	8.2	(6.8-9.9)
	Women	3,320	22.1	(20.3-24.1)	35.6	(33.4-37.9)	1.9	(1.4-2.6)	27.9	(25.9-30.0)	12.5	(11.1-14.0)
Race/ Ethnicity	Hispanic or Latino	153	34.9	(24.7-46.7)	33.2	(21.6-47.3)	3.5	(0.9-13.0)*	18.6	(11.2-29.4)	9.8	(5.6-16.4)
	Not Hispanic or Latino:	4,350	20.7	(19.2-22.4)	40.4	(38.4-42.3)	2.0	(1.6-2.6)	27.1	(25.3-28.9)	9.8	(8.8-10.9)
	White											
	Black or African American	587	27.4	(20.7-35.4)	25.0	(19.1-32.0)	1.5	(0.6-3.8)*	28.5	(22.4-35.6)	17.5	(12.4-24.2)
	All Other Races	132	24.9	(16.4-35.9)	31.5	(22.4-42.3)	1.8	(0.3-10.7)*	21.5	(13.8-32.1)	20.3	(12.9-30.6)
Education	<High School	503	37.1	(30.5-44.2)	28.3	(22.7-34.8)	2.8	(1.2-6.5)*	19.9	(15.1-25.7)	11.9	(8.6-16.3)
	High School Graduate (or GED)	1,946	28.2	(25.4-31.2)	37.2	(34.0-40.5)	2.2	(1.6-3.1)	23.3	(20.7-26.0)	9.1	(7.6-10.8)
	Some College	1,309	22.8	(19.7-26.2)	37.9	(34.3-41.6)	1.7	(1.1-2.7)	26.6	(23.5-30.0)	11.0	(9.2-13.2)
	College Graduate	1,496	15.9	(13.5-18.6)	43.1	(40.0-46.4)	1.8	(1.1-3.0)	28.0	(25.3-30.9)	11.2	(9.2-13.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	889	25.5 (21.1-30.5)	31.8 (26.4-37.8)	1.2 (0.7-2.2)	24.1 (19.9-28.8)	17.4 (14.0-21.4)				
	\$20,000-\$34,999	1,046	25.5 (21.9-29.6)	33.1 (29.1-37.3)	2.3 (1.2-4.4)*	25.4 (21.9-29.2)	13.7 (11.0-16.8)				
	\$35,000-\$49,999	784	24.2 (20.4-28.4)	39.2 (34.7-43.8)	2.5 (1.5-4.2)	24.0 (20.6-27.9)	10.1 (7.7-13.2)				
	\$50,000-\$74,999	866	23.9 (20.2-28.1)	39.9 (35.7-44.4)	1.5 (0.8-2.7)	27.9 (24.1-32.0)	6.7 (5.1-8.9)				
	\$75,000 and more	1,036	19.4 (16.2-23.1)	43.7 (39.8-47.7)	1.6 (0.9-2.6)	26.7 (23.4-30.2)	8.6 (6.9-10.8)				
Employment Status	Employed	2,864	23.8 (21.6-26.1)	39.1 (36.6-41.6)	1.9 (1.4-2.6)	25.6 (23.6-27.8)	9.6 (8.3-11.1)				
	Out of work	256	25.7 (19.5-33.1)	35.3 (27.3-44.2)	1.3 (0.5-3.7)*	26.8 (19.0-36.3)	10.9 (7.4-15.6)				
	Homemaker	435	23.9 (19.0-29.7)	38.5 (32.7-44.6)	3.9 (1.7-8.7)*	23.2 (19.0-27.9)	10.5 (7.7-14.1)				
	Student	100	20.5 (11.5-33.8)	45.1 (32.8-58.0)	1.2 (0.3-4.8)*	21.7 (13.5-32.9)	11.5 (5.5-22.5)*				
	Retired	1,229	25.3 (22.5-28.5)	37.2 (33.8-40.7)	2.7 (1.8-4.0)	25.7 (22.7-29.1)	9.1 (7.3-11.3)				
	Unable to work	360	23.2 (17.0-30.8)	26.2 (20.1-33.3)	0.8 (0.3-2.1)*	28.6 (22.6-35.6)	21.2 (16.4-26.9)				
Veteran Status	Yes	622	21.1 (17.5-25.1)	40.0 (35.0-45.1)	1.3 (0.7-2.2)	24.5 (20.2-29.5)	13.2 (9.8-17.4)				
	No	4,638	24.4 (22.6-26.2)	37.8 (35.8-39.8)	2.2 (1.7-2.8)	25.5 (23.9-27.3)	10.1 (9.1-11.3)				
Diabetes	Yes	599	20.6 (16.8-25.1)	33.2 (28.4-38.4)	2.2 (1.1-4.3)*	28.2 (23.6-33.2)	15.8 (12.5-19.8)				
	No	4,655	24.3 (22.6-26.1)	38.5 (36.6-40.5)	2.0 (1.6-2.6)	25.2 (23.6-26.9)	9.9 (8.9-11.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Indiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics		Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	599	23.2	(18.8-28.2)	34.6	(28.4-41.5)	2.7	(1.5-4.8)	23.6	(19.2-28.5)	16.0	(12.6-20.0)
	No	4,605	24.0	(22.3-25.9)	38.5	(36.6-40.5)	2.0	(1.5-2.6)	25.6	(23.9-27.3)	9.9	(8.8-11.0)
Asthma	Yes	692	22.9	(18.5-28.0)	32.3	(27.6-37.5)	0.9	(0.4-2.0)*	27.5	(23.2-32.2)	16.4	(13.2-20.2)
	No	4,557	24.2	(22.4-26.0)	39.1	(37.1-41.1)	2.2	(1.7-2.8)	25.0	(23.4-26.8)	9.5	(8.5-10.7)
Arthritis	Yes	2,046	21.5	(19.2-24.0)	33.7	(31.0-36.6)	1.6	(1.1-2.5)	29.7	(27.0-32.7)	13.4	(11.6-15.4)
	No	3,196	25.1	(23.0-27.3)	39.9	(37.6-42.3)	2.2	(1.6-2.9)	23.8	(21.9-25.8)	9.0	(7.9-10.4)
Frequent Mental Distress	Yes	605	18.2	(13.8-23.5)	30.8	(25.7-36.5)	3.0	(1.4-6.5)*	26.9	(22.4-32.0)	21.1	(17.2-25.5)
	No	4,581	24.8	(23.0-26.6)	38.9	(36.9-40.9)	2.0	(1.5-2.5)	25.2	(23.5-26.9)	9.2	(8.2-10.3)
Serious Psychological Distress	Yes	238	25.9	(19.0-34.4)	14.3	(10.0-20.2)	0.8	(0.2-3.1)*	34.3	(26.2-43.5)	24.7	(18.9-31.5)
	No	4,929	23.9	(22.3-25.7)	38.9	(37.0-40.9)	2.1	(1.6-2.7)	25.3	(23.7-26.9)	9.8	(8.8-10.9)
Receiving Treatment for Mental Health Condition	Yes	727	19.4	(15.9-23.4)	34.2	(29.8-38.9)	1.4	(0.7-2.6)*	27.4	(23.5-31.8)	17.5	(14.6-21.0)
	No	4,522	24.6	(22.8-26.4)	38.7	(36.7-40.8)	2.1	(1.6-2.7)	25.2	(23.5-26.9)	9.4	(8.4-10.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,804	67.9	(66.2-69.6)	26.3	(24.7-28.0)	0.6	(0.4-1.0)	3.5	(2.8-4.4)	1.7	(1.2-2.4)
Age (years)	18-24	174	56.1	(48.2-63.8)	37.7	(30.2-45.8)	0.2	(0.0-1.1)*	4.0	(1.7-9.1)*	2.0	(0.6-6.4)*
	25-34	517	60.7	(56.0-65.2)	30.9	(26.6-35.5)	1.3	(0.5-3.7)*	6.4	(4.1-9.8)	0.7	(0.2-2.6)*
	35-54	1,872	71.6	(69.2-73.9)	22.6	(20.5-24.9)	0.6	(0.3-1.1)*	3.3	(2.4-4.5)	1.9	(1.2-2.9)
	55+	2,228	72.0	(69.9-74.0)	23.4	(21.5-25.4)	0.4	(0.2-1.1)*	2.4	(1.8-3.2)	1.8	(1.2-2.7)
Sex	Men	1,852	57.4	(54.6-60.2)	34.3	(31.6-37.2)	1.0	(0.6-1.8)	5.0	(3.7-6.7)	2.2	(1.4-3.4)
	Women	2,952	77.7	(75.7-79.5)	18.8	(17.1-20.6)	0.2	(0.1-0.4)*	2.2	(1.6-3.0)	1.2	(0.7-2.1)
Race/ Ethnicity	Hispanic or Latino	106	68.4	(58.2-77.1)	15.1	(9.2-23.6)	0.2	(0.0-1.4)*	10.3	(4.9-20.2)*	6.1	(2.3-15.0)*
	Not Hispanic or Latino:	4,527	68.9	(67.1-70.6)	26.9	(25.2-28.6)	0.6	(0.4-1.0)	2.7	(2.1-3.4)	0.9	(0.7-1.3)
	White											
	Black or African American	66	71.6	(53.7-84.5)	16.7	(7.6-32.7)*	0.0	(0.0-0.0)	4.3	(1.0-16.7)*	7.4	(1.6-28.3)*
	All Other Races	90	60.6	(47.5-72.2)	25.3	(15.0-39.4)	0.1	(0.1-0.2)	12.5	(4.7-29.2)*	1.5	(0.3-7.0)*
Education	<High School	331	61.7	(54.0-68.9)	25.4	(18.9-33.2)	0.3	(0.1-0.8)*	5.1	(2.4-10.5)*	7.5	(5.0-11.2)
	High School Graduate (or GED)	1,733	62.5	(59.6-65.3)	31.5	(28.7-34.4)	0.4	(0.2-0.9)*	4.0	(3.0-5.4)	1.6	(0.9-2.8)*
	Some College	1,316	67.7	(64.3-71.0)	25.4	(22.4-28.8)	1.1	(0.5-2.3)*	4.5	(2.9-6.7)	1.2	(0.7-2.3)*
	College Graduate	1,417	74.3	(71.5-77.0)	22.3	(19.8-25.1)	0.7	(0.3-1.6)*	1.7	(1.0-2.8)	1.0	(0.5-2.1)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	621	62.6 (57.4-67.6)	27.4 (22.8-32.6)	0.2 (0.1-0.7)*	4.7 (2.8-7.9)	5.0 (2.6-9.4)*				
	\$20,000-\$34,999	936	65.3 (61.3-69.1)	29.0 (25.2-33.0)	0.1 (0.0-0.8)*	3.4 (2.2-5.3)	2.2 (1.3-3.7)				
	\$35,000-\$49,999	750	65.3 (60.8-69.5)	28.9 (24.8-33.3)	0.7 (0.1-3.4)*	4.4 (2.7-7.0)	0.9 (0.3-2.4)*				
	\$50,000-\$74,999	881	70.7 (66.9-74.2)	24.7 (21.3-28.4)	0.7 (0.2-2.0)*	2.7 (1.6-4.5)	1.3 (0.6-2.7)*				
	\$75,000 and more	1,060	74.5 (71.1-77.6)	22.1 (19.1-25.3)	0.8 (0.3-1.9)*	2.3 (1.2-4.1)*	0.4 (0.1-1.2)*				
Employment Status	Employed	2,907	66.6 (64.5-68.7)	27.7 (25.6-29.8)	0.7 (0.4-1.3)*	3.6 (2.7-4.8)	1.4 (0.8-2.3)				
	Out of work	114	66.5 (53.9-77.2)	21.6 (11.8-36.2)	1.9 (0.3-12.8)*	5.1 (2.1-11.8)*	4.8 (1.0-19.9)*				
	Homemaker	261	72.2 (65.3-78.1)	25.4 (19.6-32.2)	0.5 (0.1-3.3)*	0.9 (0.2-3.5)*	1.1 (0.3-4.5)*				
	Student	89	79.0 (69.1-86.4)	17.6 (10.9-27.0)	0.0 (0.0-0.0)	3.2 (1.0-10.1)*	0.1 (0.0-1.0)*				
	Retired	1,248	67.9 (64.9-70.8)	26.1 (23.4-29.0)	0.3 (0.1-0.9)*	3.3 (2.3-4.7)	2.4 (1.5-4.0)				
	Unable to work	184	68.2 (59.7-75.6)	22.8 (16.2-31.0)	0.0 (0.0-0.0)	5.1 (2.0-12.6)*	3.9 (1.9-7.7)*				
Veteran Status	Yes	576	61.9 (57.3-66.2)	29.1 (24.9-33.7)	1.5 (0.5-4.2)*	4.7 (2.6-8.2)	2.9 (1.7-4.8)				
	No	4,227	68.8 (66.9-70.6)	25.9 (24.2-27.8)	0.4 (0.2-0.7)	3.4 (2.7-4.3)	1.5 (1.0-2.3)				
Diabetes	Yes	427	72.2 (67.2-76.7)	23.1 (18.9-28.0)	0.5 (0.1-1.8)*	2.3 (1.1-4.7)*	1.9 (0.7-4.8)*				
	No	4,373	67.5 (65.7-69.3)	26.5 (24.8-28.3)	0.7 (0.4-1.1)	3.6 (2.9-4.6)	1.7 (1.2-2.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	490	65.7	(61.0-70.2)	28.6	(24.3-33.3)	0.1	(0.0-0.9)*	3.3	(2.0-5.7)	2.1	(1.1-4.3)*
	No	4,276	68.1	(66.2-69.8)	26.1	(24.4-27.9)	0.6	(0.4-1.1)	3.6	(2.8-4.6)	1.6	(1.1-2.4)
Asthma	Yes	505	74.1	(68.8-78.7)	20.1	(15.7-25.4)	0.4	(0.2-0.9)*	4.1	(2.2-7.5)*	1.3	(0.6-2.8)*
	No	4,292	67.1	(65.3-68.9)	27.0	(25.2-28.8)	0.7	(0.5-1.1)	3.4	(2.7-4.4)	1.8	(1.2-2.5)
Arthritis	Yes	1,660	71.4	(68.6-74.0)	23.4	(20.9-26.2)	0.4	(0.1-1.1)*	2.8	(2.0-3.9)	2.0	(1.3-3.1)
	No	3,133	66.7	(64.6-68.8)	27.3	(25.3-29.3)	0.7	(0.4-1.2)	3.8	(2.9-5.0)	1.6	(1.0-2.5)
Frequent Mental Distress	Yes	356	71.3	(64.5-77.2)	21.7	(15.9-28.8)	0.3	(0.0-2.1)*	4.0	(2.0-7.6)*	2.8	(1.5-5.3)*
	No	4,373	67.7	(65.9-69.5)	26.6	(24.9-28.3)	0.6	(0.4-1.0)	3.5	(2.7-4.4)	1.6	(1.1-2.3)
Serious Psychological Distress	Yes	119	54.8	(44.3-64.9)	30.2	(20.7-41.7)	0.3	(0.2-0.3)	5.0	(2.3-10.5)*	9.7	(5.8-15.9)
	No	4,605	68.3	(66.6-70.0)	26.0	(24.3-27.7)	0.7	(0.5-1.1)	3.4	(2.7-4.4)	1.6	(1.1-2.3)
Receiving Treatment for Mental Health Condition	Yes	621	80.9	(76.2-84.8)	13.4	(9.9-17.9)	0.1	(0.1-0.1)	3.4	(1.9-6.0)	2.3	(1.1-4.5)*
	No	4,162	66.3	(64.4-68.1)	27.9	(26.1-29.7)	0.7	(0.4-1.1)	3.6	(2.8-4.5)	1.6	(1.1-2.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,779	22.7	(21.2-24.3)	40.0	(38.2-41.8)	1.5	(1.1-2.0)	24.9	(23.4-26.4)	10.9	(9.9-12.1)
Age (years)	18-24	176	17.8	(12.6-24.6)	49.8	(41.8-57.9)	0.8	(0.1-5.5)*	22.0	(16.3-28.9)	9.5	(5.7-15.7)
	25-34	520	23.8	(19.8-28.3)	42.0	(37.4-46.7)	1.3	(0.6-2.6)*	25.3	(21.6-29.5)	7.7	(5.6-10.3)
	35-54	1,861	22.7	(20.6-25.0)	36.5	(34.1-38.9)	2.2	(1.6-3.1)	26.7	(24.6-28.9)	11.9	(10.3-13.6)
	55+	2,209	24.9	(23.0-27.0)	39.2	(37.0-41.5)	1.1	(0.7-1.7)	22.9	(21.1-24.9)	11.8	(10.4-13.3)
Sex	Men	1,846	24.9	(22.4-27.4)	43.0	(40.2-45.9)	1.6	(1.1-2.3)	21.9	(19.8-24.1)	8.7	(7.1-10.5)
	Women	2,933	20.7	(19.0-22.5)	37.1	(35.0-39.3)	1.4	(0.9-2.2)	27.7	(25.8-29.8)	13.0	(11.6-14.5)
Race/ Ethnicity	Hispanic or Latino	109	42.7	(32.1-54.1)	19.9	(12.9-29.4)	2.4	(0.8-7.4)*	20.6	(13.5-30.3)	14.3	(6.7-28.0)*
	Not Hispanic or Latino:	4,504	19.2	(17.8-20.6)	42.0	(40.2-43.9)	1.6	(1.2-2.1)	26.7	(25.1-28.3)	10.6	(9.5-11.7)
	White											
	Black or African American	64	25.1	(12.2-44.5)*	34.7	(19.3-54.1)	1.9	(0.3-12.6)*	19.1	(10.4-32.3)	19.2	(10.1-33.5)*
	All Other Races	88	34.0	(21.3-49.5)	32.0	(21.1-45.3)	0.2	(0.2-0.3)	18.4	(10.0-31.4)	15.4	(9.6-23.7)
Education	<High School	317	33.8	(27.4-40.7)	34.0	(26.7-42.2)	2.2	(0.6-7.8)*	17.2	(11.8-24.3)	12.9	(8.6-18.9)
	High School Graduate (or GED)	1,747	27.6	(25.0-30.5)	39.3	(36.4-42.3)	1.1	(0.6-1.7)	21.5	(19.3-23.9)	10.5	(8.7-12.5)
	Some College	1,301	21.8	(19.0-24.9)	39.5	(36.1-43.0)	1.3	(0.8-2.1)	25.9	(23.0-28.9)	11.6	(9.7-13.8)
	College Graduate	1,406	14.4	(12.3-16.8)	43.3	(40.3-46.3)	2.1	(1.4-3.2)	29.6	(26.9-32.4)	10.6	(8.9-12.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	618	27.9	(23.7-32.4)	33.5	(28.3-39.0)	1.5	(0.3-6.5)*	21.9	(18.0-26.4)	15.2	(12.1-19.0)
	\$20,000-\$34,999	931	24.2	(20.7-28.0)	36.7	(32.7-40.8)	0.7	(0.4-1.3)*	24.4	(21.2-27.9)	14.1	(11.1-17.7)
	\$35,000-\$49,999	750	21.8	(18.3-25.8)	41.7	(37.3-46.2)	2.4	(1.4-3.9)	23.4	(20.1-27.1)	10.7	(8.5-13.4)
	\$50,000-\$74,999	876	19.5	(16.6-22.7)	44.3	(40.5-48.2)	1.9	(1.1-3.2)	24.8	(21.8-28.2)	9.5	(7.6-11.7)
	\$75,000 and more	1,057	20.6	(17.7-23.8)	41.5	(38.0-45.0)	1.3	(0.7-2.2)	28.1	(25.1-31.3)	8.6	(6.8-10.7)
Employment Status	Employed	2,893	23.0	(21.1-25.1)	40.1	(37.9-42.3)	1.8	(1.3-2.4)	25.0	(23.2-26.8)	10.2	(8.9-11.6)
	Out of work	115	18.1	(10.1-30.3)	43.4	(30.9-56.8)	0.8	(0.2-4.0)*	18.2	(11.8-26.9)	19.6	(13.0-28.3)
	Homemaker	268	25.7	(20.4-31.8)	36.4	(30.3-43.0)	1.2	(0.4-3.3)*	27.2	(21.9-33.2)	9.5	(6.5-13.7)
	Student	89	20.7	(12.3-32.8)	44.7	(32.9-57.1)	2.3	(0.3-14.4)*	22.9	(14.7-33.9)	9.4	(3.9-21.0)*
	Retired	1,234	23.2	(20.8-25.8)	41.1	(37.9-44.3)	1.0	(0.6-1.6)	24.3	(21.6-27.2)	10.4	(8.6-12.5)
	Unable to work	179	19.6	(13.8-27.2)	25.9	(18.9-34.4)	0.5	(0.1-2.3)*	28.2	(20.7-37.3)	25.7	(19.6-32.9)
Veteran Status	Yes	578	20.0	(16.7-23.8)	35.0	(30.5-39.8)	1.5	(0.8-3.1)*	31.4	(26.7-36.5)	12.1	(8.7-16.6)
	No	4,201	23.1	(21.5-24.8)	40.6	(38.7-42.5)	1.5	(1.1-2.0)	24.0	(22.5-25.6)	10.8	(9.7-12.0)
Diabetes	Yes	420	21.5	(17.5-26.1)	38.9	(33.8-44.3)	0.6	(0.3-1.4)*	26.9	(22.3-32.0)	12.1	(9.3-15.4)
	No	4,355	22.8	(21.2-24.5)	40.0	(38.2-41.9)	1.6	(1.2-2.1)	24.7	(23.2-26.3)	10.8	(9.7-12.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Iowa, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics		Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	484	24.0	(20.4-28.0)	36.1	(31.3-41.2)	0.4	(0.1-1.9)*	25.2	(20.9-30.0)	14.4	(11.2-18.3)
	No	4,256	22.5	(20.9-24.2)	40.3	(38.4-42.2)	1.6	(1.2-2.1)	24.9	(23.4-26.5)	10.7	(9.6-11.9)
Asthma	Yes	498	25.0	(20.5-30.2)	35.9	(30.5-41.6)	0.5	(0.1-1.7)*	23.3	(19.1-28.2)	15.3	(12.1-19.2)
	No	4,271	22.5	(20.9-24.1)	40.4	(38.5-42.3)	1.6	(1.2-2.1)	25.1	(23.6-26.7)	10.5	(9.4-11.7)
Arthritis	Yes	1,646	19.5	(17.5-21.7)	36.3	(33.5-39.2)	1.1	(0.6-1.8)	27.7	(25.2-30.3)	15.5	(13.4-17.7)
	No	3,122	24.0	(22.1-26.0)	41.2	(39.1-43.5)	1.7	(1.2-2.3)	23.9	(22.1-25.7)	9.3	(8.1-10.6)
Frequent Mental Distress	Yes	355	15.6	(11.5-20.8)	32.3	(25.7-39.6)	2.9	(0.9-9.1)*	29.5	(23.7-36.0)	19.7	(15.5-24.8)
	No	4,353	23.4	(21.8-25.0)	40.6	(38.7-42.5)	1.4	(1.1-1.8)	24.4	(22.9-26.0)	10.2	(9.1-11.4)
Serious Psychological Distress	Yes	119	12.6	(7.8-19.9)	39.7	(29.1-51.2)	3.1	(0.8-11.1)*	24.4	(16.0-35.5)	20.2	(14.1-28.0)
	No	4,580	23.1	(21.5-24.7)	40.0	(38.1-41.8)	1.5	(1.1-2.0)	24.9	(23.4-26.4)	10.6	(9.6-11.8)
Receiving Treatment for Mental Health Condition	Yes	616	19.0	(15.2-23.3)	32.1	(27.5-37.0)	2.2	(0.8-5.7)*	27.7	(23.3-32.5)	19.1	(15.8-23.0)
	No	4,143	23.2	(21.6-24.9)	41.0	(39.1-42.9)	1.4	(1.1-1.8)	24.5	(23.0-26.1)	9.9	(8.8-11.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		8,410	65.0	(63.6-66.5)	27.1	(25.8-28.5)	3.8	(3.1-4.7)	3.1	(2.6-3.7)	0.9	(0.7-1.3)
Age (years)	18-24	209	56.4	(49.0-63.5)	33.6	(27.0-40.8)	6.0	(3.6-9.8)	3.6	(1.8-7.2)*	0.4	(0.1-2.4)*
	25-34	715	62.3	(58.3-66.2)	28.4	(24.8-32.3)	4.0	(2.6-6.0)	3.8	(2.4-5.7)	1.5	(0.7-3.2)*
	35-54	2,939	68.5	(66.4-70.5)	25.1	(23.2-27.1)	3.1	(2.4-4.1)	2.5	(1.9-3.3)	0.7	(0.5-1.2)
	55+	4,503	65.0	(63.3-66.6)	26.1	(24.6-27.7)	4.5	(3.7-5.5)	3.2	(2.6-3.9)	1.3	(0.9-1.7)
Sex	Men	3,188	60.4	(58.1-62.7)	29.7	(27.6-31.8)	4.6	(3.5-6.0)	4.3	(3.4-5.3)	1.1	(0.6-1.7)
	Women	5,222	69.3	(67.6-71.0)	24.8	(23.2-26.4)	3.1	(2.4-3.9)	2.1	(1.6-2.7)	0.8	(0.5-1.2)
Race/ Ethnicity	Hispanic or Latino	313	64.1	(57.3-70.4)	26.2	(20.6-32.9)	6.2	(3.8-9.8)	2.4	(1.3-4.3)*	1.1	(0.4-2.9)*
	Not Hispanic or Latino:	7,467	64.4	(62.9-65.8)	26.5	(25.1-27.9)	5.0	(4.4-5.6)	3.1	(2.7-3.7)	1.0	(0.8-1.3)
	White											
	Black or African American	318	58.7	(50.9-66.1)	27.3	(20.7-35.0)	6.7	(4.5-9.8)	5.1	(2.2-11.3)*	2.2	(1.0-4.7)*
	All Other Races	284	53.8	(46.2-61.2)	25.7	(19.2-33.6)	13.5	(10.0-17.8)	5.0	(2.7-9.1)*	2.1	(0.5-7.6)*
Education	<High School	483	55.4	(49.3-61.4)	37.4	(31.6-43.6)	2.7	(1.6-4.5)	3.5	(1.7-7.1)*	1.0	(0.5-1.8)*
	High School Graduate (or GED)	2,444	61.4	(58.6-64.0)	29.4	(26.8-32.0)	4.7	(3.6-6.1)	3.7	(2.8-4.8)	0.9	(0.5-1.6)
	Some College	2,435	66.3	(63.6-69.0)	25.7	(23.3-28.2)	4.1	(2.9-5.8)	3.0	(2.2-4.0)	0.9	(0.5-1.6)
	College Graduate	3,041	69.0	(66.6-71.3)	24.3	(22.2-26.5)	3.3	(1.9-5.4)	2.6	(1.8-3.7)	0.9	(0.5-1.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,044	62.9	(58.4-67.2)	25.3	(21.7-29.4)	5.1	(3.2-8.1)	5.2	(3.3-7.9)	1.5	(0.8-2.6)
	\$20,000-\$34,999	1,703	62.7	(59.4-65.9)	30.2	(27.2-33.4)	3.0	(2.1-4.3)	3.0	(2.1-4.4)	1.1	(0.6-1.9)
	\$35,000-\$49,999	1,271	64.6	(61.1-68.0)	27.6	(24.4-31.0)	3.2	(2.1-4.7)	3.4	(2.4-4.9)	1.2	(0.5-2.9)*
	\$50,000-\$74,999	1,371	71.4	(68.3-74.4)	22.5	(19.7-25.4)	3.3	(2.3-4.8)	2.3	(1.5-3.7)	0.5	(0.2-1.0)*
	\$75,000 and more	2,109	68.4	(65.7-71.1)	26.4	(23.9-29.0)	2.3	(1.6-3.3)	2.3	(1.5-3.5)	0.6	(0.3-1.4)*
Employment Status	Employed	4,676	66.5	(64.6-68.4)	26.7	(25.0-28.5)	3.3	(2.6-4.3)	2.6	(2.1-3.4)	0.7	(0.4-1.3)
	Out of work	350	67.0	(60.5-72.9)	24.0	(18.8-30.0)	4.8	(2.4-9.1)*	2.8	(1.1-6.7)*	1.5	(0.4-4.9)*
	Homemaker	501	62.3	(57.1-67.3)	28.7	(23.9-33.9)	4.1	(2.2-7.5)*	3.8	(2.0-7.1)*	1.0	(0.3-3.3)*
	Student	112	66.4	(55.3-75.9)	25.7	(17.5-36.1)	6.2	(2.4-14.9)*	1.3	(0.3-4.9)*	0.4	(0.1-3.1)*
	Retired	2,332	59.1	(56.4-61.7)	31.5	(28.9-34.2)	4.7	(3.4-6.3)	3.7	(2.7-4.9)	1.1	(0.7-1.6)
	Unable to work	425	65.1	(59.6-70.3)	22.5	(18.1-27.6)	2.8	(1.2-6.3)*	8.3	(5.2-13.0)	1.3	(0.6-2.7)*
Veteran Status	Yes	1,080	64.0	(60.3-67.6)	27.3	(24.0-30.8)	4.9	(3.5-6.9)	2.7	(1.7-4.2)	1.1	(0.5-2.4)*
	No	7,328	65.1	(63.5-66.7)	27.2	(25.7-28.7)	3.6	(2.9-4.5)	3.2	(2.6-3.9)	0.9	(0.6-1.2)
Diabetes	Yes	989	65.5	(61.9-69.0)	26.8	(23.5-30.4)	4.0	(2.2-6.9)	3.0	(2.1-4.4)	0.7	(0.3-1.3)*
	No	7,419	65.0	(63.4-66.5)	27.2	(25.7-28.6)	3.8	(3.1-4.7)	3.1	(2.6-3.8)	0.9	(0.7-1.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	930	62.3	(58.4-66.0)	28.7	(25.3-32.5)	4.9	(3.4-7.1)	3.2	(2.0-5.0)	0.9	(0.5-1.6)
	No	7,416	65.3	(63.7-66.8)	27.0	(25.5-28.4)	3.7	(3.0-4.6)	3.1	(2.6-3.7)	0.9	(0.7-1.3)
Asthma	Yes	921	68.6	(64.4-72.5)	23.6	(20.0-27.6)	3.3	(2.1-5.2)	3.8	(2.2-6.6)	0.7	(0.3-1.8)*
	No	7,462	64.5	(63.0-66.1)	27.6	(26.2-29.1)	3.9	(3.1-4.8)	3.1	(2.5-3.7)	0.9	(0.7-1.3)
Arthritis	Yes	2,845	66.4	(64.0-68.7)	26.0	(23.8-28.2)	3.6	(2.6-4.9)	3.3	(2.5-4.3)	0.7	(0.5-1.1)
	No	5,532	64.7	(62.9-66.4)	27.5	(25.9-29.2)	3.9	(3.1-4.8)	3.0	(2.4-3.6)	1.0	(0.7-1.5)
Frequent Mental Distress	Yes	729	68.1	(63.5-72.4)	23.0	(19.2-27.4)	3.0	(1.9-4.6)	4.7	(2.7-7.9)	1.2	(0.6-2.3)*
	No	7,592	64.8	(63.2-66.3)	27.5	(26.1-29.0)	3.9	(3.1-4.8)	2.9	(2.4-3.5)	0.9	(0.6-1.3)
Serious Psychological Distress	Yes	205	60.3	(51.2-68.8)	25.2	(18.2-33.9)	5.2	(2.6-10.1)*	7.6	(3.5-15.8)*	1.7	(0.7-3.9)*
	No	8,068	65.5	(64.0-67.0)	27.3	(25.9-28.7)	3.5	(2.9-4.2)	2.9	(2.4-3.5)	0.8	(0.6-1.2)
Receiving Treatment for Mental Health Condition	Yes	1,082	76.0	(72.5-79.2)	17.1	(14.3-20.3)	1.8	(1.1-3.0)	4.0	(2.3-6.7)	1.0	(0.6-1.9)*
	No	7,305	63.7	(62.1-65.3)	28.3	(26.9-29.8)	4.1	(3.3-5.0)	3.0	(2.5-3.6)	0.9	(0.6-1.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		8,414	22.2	(21.0-23.5)	38.2	(36.8-39.6)	6.1	(5.3-6.9)	24.3	(23.2-25.5)	9.2	(8.4-10.0)
Age (years)	18-24	209	21.3	(16.0-27.7)	46.9	(39.6-54.3)	5.9	(3.4-10.1)	21.8	(16.3-28.6)	4.1	(2.1-7.7)*
	25-34	714	16.5	(13.7-19.8)	41.3	(37.3-45.4)	5.6	(4.0-7.7)	25.3	(22.1-28.7)	11.3	(8.9-14.2)
	35-54	2,950	22.0	(20.2-23.9)	36.9	(34.8-39.0)	6.2	(5.2-7.5)	25.0	(23.3-26.8)	9.9	(8.7-11.2)
	55+	4,499	26.4	(24.9-28.0)	35.3	(33.7-36.9)	6.1	(5.3-7.1)	23.3	(22.0-24.7)	8.9	(8.0-9.8)
Sex	Men	3,207	25.0	(23.1-27.0)	40.8	(38.6-43.1)	6.5	(5.4-7.8)	20.1	(18.5-21.9)	7.6	(6.5-8.9)
	Women	5,207	19.6	(18.2-21.1)	35.7	(33.9-37.4)	5.7	(4.8-6.6)	28.4	(26.8-30.0)	10.7	(9.7-11.8)
Race/ Ethnicity	Hispanic or Latino	316	29.5	(23.5-36.2)	39.4	(32.6-46.7)	3.6	(1.6-7.7)*	19.8	(15.0-25.6)	7.7	(5.1-11.6)
	Not Hispanic or Latino:	7,458	20.1	(18.9-21.3)	36.6	(35.1-38.1)	9.6	(8.9-10.3)	24.8	(23.6-26.1)	8.9	(8.1-9.8)
	White											
	Black or African American	323	21.5	(15.5-29.0)	33.5	(26.2-41.7)	15.5	(11.8-20.2)	19.7	(14.9-25.5)	9.8	(6.6-14.5)
	All Other Races	288	25.6	(19.6-32.6)	36.4	(29.4-44.0)	7.3	(4.7-11.3)	19.4	(14.6-25.3)	11.3	(7.4-16.7)
Education	<High School	498	27.9	(23.0-33.3)	41.5	(35.5-47.7)	2.9	(1.8-4.5)	16.8	(12.7-21.7)	11.0	(7.7-15.4)
	High School Graduate (or GED)	2,437	26.2	(23.8-28.7)	39.0	(36.3-41.8)	5.8	(4.7-7.2)	20.7	(18.6-22.9)	8.3	(7.0-9.9)
	Some College	2,439	21.2	(19.1-23.4)	34.3	(31.8-36.9)	6.9	(5.6-8.5)	26.0	(23.8-28.3)	11.6	(9.9-13.4)
	College Graduate	3,032	17.5	(15.8-19.4)	40.1	(37.8-42.5)	6.6	(5.3-8.1)	27.8	(25.8-29.9)	8.0	(6.9-9.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,067	23.2	(19.7-27.0)	33.4	(29.2-37.8)	5.2	(3.5-7.5)	24.7	(21.1-28.6)	13.6	(11.0-16.8)
	\$20,000-\$34,999	1,695	24.8	(22.1-27.6)	39.7	(36.4-43.0)	4.0	(3.0-5.2)	22.8	(20.3-25.6)	8.7	(7.0-10.8)
	\$35,000-\$49,999	1,273	21.8	(18.9-25.1)	36.8	(33.4-40.4)	8.0	(6.2-10.2)	22.6	(19.9-25.4)	10.8	(8.7-13.4)
	\$50,000-\$74,999	1,369	19.7	(17.3-22.4)	40.4	(37.1-43.8)	5.0	(3.8-6.7)	26.1	(23.3-29.1)	8.7	(7.1-10.7)
	\$75,000 and more	2,111	20.7	(18.4-23.1)	39.8	(37.1-42.5)	5.6	(4.5-7.1)	26.5	(24.3-28.8)	7.4	(6.2-8.7)
Employment Status	Employed	4,688	22.8	(21.2-24.5)	38.7	(36.9-40.6)	6.2	(5.3-7.3)	23.6	(22.2-25.1)	8.7	(7.7-9.8)
	Out of work	346	20.3	(15.1-26.7)	38.3	(31.6-45.4)	3.9	(2.2-7.1)*	25.3	(20.0-31.3)	12.2	(8.3-17.6)
	Homemaker	504	19.2	(15.3-23.8)	40.6	(35.6-45.9)	3.6	(2.3-5.6)	25.3	(21.4-29.8)	11.2	(8.3-15.0)
	Student	110	16.0	(9.3-26.0)	36.6	(26.4-48.1)	11.0	(5.5-21.0)*	29.1	(19.7-40.8)	7.3	(3.3-15.5)*
	Retired	2,325	24.5	(22.4-26.8)	38.3	(35.7-41.0)	6.3	(5.0-7.8)	24.1	(21.9-26.3)	6.8	(5.7-8.1)
	Unable to work	427	18.4	(14.4-23.2)	26.8	(21.8-32.5)	6.3	(3.8-10.3)	29.1	(23.6-35.4)	19.3	(15.4-23.9)
Veteran Status	Yes	1,084	20.6	(17.9-23.6)	35.8	(32.3-39.6)	6.9	(5.1-9.1)	26.3	(22.9-29.9)	10.4	(8.0-13.5)
	No	7,328	22.5	(21.1-23.8)	38.4	(36.9-40.0)	5.9	(5.2-6.8)	24.1	(22.9-25.4)	9.1	(8.3-9.9)
Diabetes	Yes	986	22.0	(19.2-25.2)	37.3	(33.5-41.3)	6.7	(4.6-9.6)	22.4	(19.4-25.8)	11.5	(9.4-14.1)
	No	7,427	22.3	(21.0-23.6)	38.3	(36.8-39.8)	6.0	(5.2-6.8)	24.5	(23.3-25.8)	9.0	(8.2-9.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kansas, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	932	22.2 (19.2-25.5)	36.9 (33.1-40.9)	5.3 (4.0-7.0)	25.1 (21.8-28.8)	10.5 (8.4-13.0)				
	No	7,417	22.3 (21.0-23.6)	38.3 (36.8-39.8)	6.1 (5.4-7.0)	24.2 (23.0-25.5)	9.1 (8.3-10.0)				
Asthma	Yes	912	20.6 (17.1-24.5)	34.8 (30.7-39.1)	5.0 (3.6-6.9)	28.1 (24.4-32.1)	11.6 (9.3-14.3)				
	No	7,475	22.4 (21.1-23.7)	38.7 (37.2-40.2)	6.2 (5.4-7.1)	23.9 (22.7-25.1)	8.9 (8.0-9.7)				
Arthritis	Yes	2,835	21.1 (19.3-23.1)	33.9 (31.7-36.3)	5.6 (4.6-6.9)	27.2 (25.1-29.4)	12.2 (10.7-13.9)				
	No	5,547	22.5 (21.1-24.1)	39.5 (37.8-41.2)	6.2 (5.4-7.2)	23.5 (22.2-25.0)	8.2 (7.3-9.2)				
Frequent Mental Distress	Yes	718	20.1 (16.0-24.9)	29.7 (25.3-34.6)	4.0 (2.6-6.1)	27.9 (23.9-32.3)	18.3 (15.0-22.1)				
	No	7,608	22.5 (21.3-23.9)	39.0 (37.5-40.5)	6.3 (5.5-7.1)	23.9 (22.7-25.2)	8.3 (7.5-9.1)				
Serious Psychological Distress	Yes	204	16.9 (10.4-26.1)	21.5 (14.8-30.3)	5.4 (2.6-10.9)*	30.0 (21.9-39.5)	26.2 (19.2-34.8)				
	No	8,076	22.3 (21.1-23.6)	38.7 (37.3-40.2)	6.1 (5.3-6.9)	24.3 (23.1-25.5)	8.6 (7.9-9.5)				
Receiving Treatment for Mental Health Condition	Yes	1,075	16.7 (14.1-19.8)	32.0 (28.3-36.0)	8.4 (6.4-10.9)	28.4 (25.1-32.1)	14.4 (12.0-17.2)				
	No	7,318	22.8 (21.5-24.1)	37.7 (36.2-39.3)	7.7 (6.9-8.5)	23.3 (22.1-24.6)	8.5 (7.7-9.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,916	66.1	(64.1-68.0)	25.1	(23.3-27.0)	5.3	(4.3-6.4)	1.5	(1.2-2.0)	2.0	(1.2-3.2)
Age (years)	18-24	221	74.8	(66.6-81.5)	13.9	(9.6-19.6)	5.4	(2.8-10.3)*	0.9	(0.3-2.5)*	5.0	(1.4-16.4)*
	25-34	614	63.5	(57.8-68.8)	24.8	(20.1-30.1)	8.5	(5.3-13.3)	1.0	(0.5-2.1)*	2.2	(1.1-4.3)*
	35-54	2,111	67.6	(64.6-70.5)	25.5	(22.7-28.4)	3.8	(2.8-5.2)	1.5	(1.0-2.4)	1.6	(0.7-3.4)*
	55+	2,952	63.8	(61.4-66.2)	27.5	(25.3-29.9)	5.3	(4.5-6.4)	1.9	(1.3-2.8)	1.4	(0.9-2.1)
Sex	Men	1,927	60.3	(56.8-63.6)	29.3	(26.2-32.6)	6.3	(4.7-8.5)	1.7	(1.1-2.6)	2.4	(1.1-5.0)*
	Women	3,989	71.5	(69.4-73.5)	21.1	(19.3-23.1)	4.3	(3.5-5.3)	1.4	(1.1-1.9)	1.7	(1.0-2.6)
Race/ Ethnicity	Hispanic or Latino	71	53.3	(32.0-73.5)	34.5	(18.3-55.2)	8.3	(2.6-23.3)*	3.7	(1.1-11.7)*	0.2	(0.0-1.7)*
	Not Hispanic or Latino:	5,554	66.7	(64.7-68.7)	25.1	(23.3-27.0)	5.4	(4.5-6.5)	1.5	(1.1-1.9)	1.3	(0.9-2.0)
	White											
	Black or African American	187	66.9	(57.4-75.2)	21.3	(14.4-30.4)	8.8	(4.2-17.4)*	1.3	(0.4-3.6)*	1.7	(0.4-6.9)*
	All Other Races	84	0.0	(0.0-3.6)*	0.0	(0.0-3.6)*	0.0	(0.0-3.6)*	0.0	(0.0-3.6)*	100	(96.4-100)*
Education	<High School	1,085	49.2	(44.3-54.1)	31.0	(26.1-36.3)	11.9	(8.9-15.7)	1.6	(0.9-3.0)*	6.4	(2.8-13.8)*
	High School Graduate (or GED)	2,324	58.9	(55.7-62.1)	28.0	(25.1-31.0)	9.6	(7.8-11.8)	2.0	(1.3-3.0)	1.5	(1.0-2.5)
	Some College	1,284	71.4	(67.2-75.3)	21.1	(17.7-25.0)	4.7	(3.2-6.8)	1.4	(0.8-2.5)	1.3	(0.5-3.4)*
	College Graduate	1,207	72.4	(68.4-76.2)	22.5	(19.0-26.5)	2.9	(1.8-4.8)	1.3	(0.8-2.1)	0.8	(0.3-2.3)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,143	59.0	(54.3-63.5)	29.3	(25.2-33.7)	6.8	(4.5-10.3)	2.1	(1.2-3.6)	2.8	(1.8-4.3)
	\$20,000-\$34,999	1,149	58.5	(53.8-63.0)	28.8	(24.6-33.5)	6.4	(4.3-9.6)	1.3	(0.8-2.2)	4.9	(2.1-11.0)*
	\$35,000-\$49,999	717	66.7	(61.5-71.5)	27.8	(23.2-32.9)	3.6	(2.0-6.4)	1.4	(0.5-4.1)*	0.5	(0.2-1.6)*
	\$50,000-\$74,999	647	67.0	(61.0-72.6)	26.4	(21.2-32.5)	4.4	(2.4-7.9)*	0.6	(0.2-1.7)*	1.5	(0.5-4.0)*
	\$75,000 and more	626	73.4	(67.8-78.3)	19.0	(14.9-24.0)	6.1	(3.2-11.2)*	1.1	(0.5-2.3)*	0.4	(0.1-1.1)*
Employment Status	Employed	2,499	68.3	(65.3-71.2)	23.1	(20.6-25.9)	5.1	(3.7-6.9)	1.3	(0.8-1.9)	2.2	(1.0-4.7)*
	Out of work	175	63.9	(52.6-73.8)	19.7	(12.5-29.7)	15.2	(7.5-28.5)*	1.1	(0.3-3.8)*	0.0	(0.0-0.0)
	Homemaker	573	66.1	(60.6-71.2)	27.4	(22.5-32.9)	3.8	(2.5-6.0)	1.3	(0.6-2.6)*	1.3	(0.5-3.2)*
	Student	80	83.0	(71.0-90.7)	11.5	(6.0-21.0)*	2.3	(0.4-11.7)*	0.2	(0.0-1.2)*	3.0	(0.4-18.2)*
	Retired	1,695	58.0	(54.7-61.1)	32.6	(29.5-35.9)	5.5	(4.4-6.9)	2.7	(1.7-4.3)	1.3	(0.7-2.2)
	Unable to work	886	60.9	(55.8-65.7)	29.0	(24.6-33.9)	4.7	(3.3-6.7)	2.1	(1.2-3.8)	3.2	(2.1-5.1)
Veteran Status	Yes	662	64.0	(58.9-68.9)	27.9	(23.4-32.8)	3.9	(2.5-6.1)	3.3	(1.8-6.2)*	0.8	(0.3-2.0)*
	No	5,254	66.3	(64.2-68.5)	24.7	(22.7-26.7)	5.5	(4.5-6.8)	1.3	(1.0-1.7)	2.2	(1.3-3.6)
Diabetes	Yes	803	65.5	(60.5-70.1)	27.1	(22.7-32.0)	4.2	(3.0-5.7)	2.0	(1.2-3.1)	1.3	(0.6-2.8)*
	No	5,110	65.5	(63.3-67.6)	24.6	(22.7-26.6)	6.3	(5.3-7.5)	1.5	(1.1-2.0)	2.1	(1.3-3.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	924	58.6	(54.2-62.9)	32.9	(28.8-37.4)	5.1	(3.7-6.9)	2.2	(1.2-4.0)*	1.2	(0.5-2.5)*
	No	4,941	67.1	(64.9-69.2)	24.1	(22.2-26.1)	5.2	(4.2-6.5)	1.5	(1.1-1.9)	2.1	(1.3-3.5)
Asthma	Yes	777	58.2	(52.5-63.7)	27.3	(22.1-33.2)	11.9	(9.1-15.3)	1.7	(0.8-3.3)*	0.9	(0.4-2.0)*
	No	5,131	66.2	(64.0-68.2)	24.4	(22.5-26.3)	5.7	(4.7-6.9)	1.6	(1.2-2.1)	2.2	(1.3-3.6)
Arthritis	Yes	2,474	64.2	(61.3-67.1)	26.3	(23.6-29.1)	5.8	(4.4-7.5)	2.5	(1.7-3.5)	1.3	(0.9-1.9)
	No	3,427	67.0	(64.4-69.5)	24.5	(22.3-26.9)	5.0	(3.8-6.4)	1.1	(0.8-1.7)	2.4	(1.3-4.2)
Frequent Mental Distress	Yes	859	60.8	(55.5-65.8)	26.1	(21.5-31.4)	7.6	(4.2-13.4)	2.6	(1.6-4.3)	2.9	(1.8-4.5)
	No	4,920	66.8	(64.6-68.9)	25.2	(23.2-27.2)	4.8	(4.0-5.9)	1.4	(1.0-1.9)	1.8	(1.0-3.3)*
Serious Psychological Distress	Yes	449	56.4	(49.4-63.1)	23.0	(17.5-29.7)	12.9	(8.5-19.1)	4.8	(2.8-8.2)	2.9	(1.5-5.5)*
	No	5,327	65.9	(63.8-68.0)	25.1	(23.2-27.1)	5.8	(4.8-6.9)	1.3	(1.0-1.8)	1.9	(1.1-3.3)
Receiving Treatment for Mental Health Condition	Yes	981	71.3	(66.9-75.3)	19.2	(15.8-23.1)	4.6	(2.5-8.0)	2.6	(1.5-4.5)	2.4	(1.4-3.9)
	No	4,929	65.4	(63.2-67.5)	25.9	(24.0-28.0)	5.4	(4.4-6.5)	1.4	(1.0-1.8)	2.0	(1.1-3.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,899	25.2	(23.3-27.1)	28.5	(26.6-30.4)	10.6	(9.2-12.1)	19.3	(17.7-21.0)	16.4	(14.9-18.1)
Age (years)	18-24	225	22.6	(15.9-31.1)	28.2	(20.9-36.8)	14.5	(9.0-22.4)	18.8	(13.0-26.3)	16.0	(9.4-25.8)
	25-34	613	28.0	(22.9-33.7)	23.2	(19.0-27.9)	10.8	(7.3-15.7)	21.8	(17.2-27.2)	16.2	(12.7-20.6)
	35-54	2,114	24.5	(21.7-27.5)	28.3	(25.5-31.3)	9.5	(7.7-11.6)	19.0	(16.7-21.4)	18.7	(16.4-21.2)
	55+	2,928	25.6	(23.5-27.8)	31.6	(29.2-34.1)	10.6	(9.2-12.3)	18.2	(16.4-20.2)	14.0	(12.3-15.9)
Sex	Men	1,937	27.3	(24.3-30.5)	30.4	(27.4-33.6)	12.1	(9.9-14.6)	18.2	(15.6-21.1)	12.0	(9.8-14.6)
	Women	3,962	23.2	(21.1-25.4)	26.6	(24.5-28.8)	9.3	(7.8-11.0)	20.4	(18.6-22.4)	20.5	(18.6-22.6)
Race/ Ethnicity	Hispanic or Latino	70	7.3	(2.7-18.1)*	36.5	(20.2-56.7)	12.1	(5.8-23.4)*	22.1	(10.3-41.0)*	21.9	(10.7-39.7)*
	Not Hispanic or Latino:	5,540	22.2	(20.4-24.0)	29.4	(27.4-31.3)	11.7	(10.4-13.2)	20.5	(18.7-22.3)	16.3	(14.8-17.8)
	White											
	Black or African American	188	30.5	(22.7-39.6)	22.2	(14.9-31.7)	15.3	(8.1-27.2)*	15.4	(9.2-24.8)	16.5	(10.3-25.4)
	All Other Races	81	21.5	(8.2-45.9)*	24.3	(10.4-47.2)*	4.8	(1.7-12.9)*	15.9	(8.1-28.9)*	33.5	(9.5-70.7)*
Education	<High School	1,073	26.4	(22.2-31.0)	30.4	(26.0-35.3)	12.7	(9.4-16.9)	15.6	(11.8-20.4)	14.9	(10.4-21.0)
	High School Graduate (or GED)	2,325	28.9	(25.8-32.2)	27.8	(25.0-30.7)	10.7	(8.6-13.2)	16.8	(14.7-19.1)	15.8	(13.7-18.2)
	Some College	1,281	24.0	(20.2-28.2)	28.1	(24.2-32.3)	12.1	(9.3-15.6)	18.6	(15.5-22.2)	17.3	(14.4-20.5)
	College Graduate	1,205	25.9	(22.2-29.9)	27.7	(24.0-31.7)	6.8	(5.1-9.2)	22.9	(19.5-26.8)	16.6	(13.7-20.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,125	21.9	(18.4-25.9)	27.6	(23.5-32.1)	13.0	(9.7-17.2)	18.2	(15.0-21.9)	19.3	(16.0-22.9)
	\$20,000-\$34,999	1,135	21.7	(18.1-25.8)	26.8	(23.0-31.0)	10.5	(7.7-14.0)	21.6	(17.4-26.5)	19.4	(15.1-24.6)
	\$35,000-\$49,999	720	23.2	(19.0-28.0)	31.5	(26.7-36.8)	10.9	(7.9-14.8)	20.6	(16.4-25.5)	13.8	(10.8-17.5)
	\$50,000-\$74,999	641	26.6	(21.5-32.5)	30.4	(25.4-36.0)	10.1	(7.1-14.0)	15.3	(11.8-19.6)	17.6	(13.6-22.4)
	\$75,000 and more	629	32.3	(26.8-38.4)	26.8	(22.2-32.0)	10.0	(6.8-14.5)	16.5	(13.2-20.6)	14.3	(10.9-18.6)
Employment Status	Employed	2,506	27.5	(24.6-30.5)	24.6	(22.1-27.2)	12.4	(10.5-14.5)	19.2	(16.9-21.7)	16.4	(14.1-19.0)
	Out of work	174	17.3	(10.0-28.3)	30.3	(20.7-42.0)	18.0	(9.1-32.7)*	22.1	(14.4-32.3)	12.2	(7.1-20.3)
	Homemaker	572	22.6	(17.7-28.3)	36.1	(30.4-42.3)	6.9	(4.6-10.2)	15.4	(12.0-19.5)	19.0	(14.4-24.5)
	Student	81	0.0	(0.0-0.0)	79.8	(60.1-91.2)	20.2	(8.8-39.9)*	0.0	(0.0-0.0)	0.0	(0.0-0.0)
	Retired	1,670	22.5	(20.0-25.3)	33.8	(30.5-37.3)	13.7	(11.7-15.9)	18.4	(15.8-21.2)	11.6	(9.6-13.9)
	Unable to work	887	17.2	(13.8-21.2)	30.3	(25.7-35.3)	12.5	(9.5-16.3)	21.0	(17.1-25.6)	19.0	(15.6-23.0)
Veteran Status	Yes	661	26.1	(21.8-31.0)	28.5	(24.1-33.4)	9.9	(7.5-13.0)	20.4	(16.2-25.3)	15.0	(11.1-19.9)
	No	5,238	24.1	(22.1-26.2)	27.4	(25.5-29.5)	14.4	(12.8-16.1)	18.4	(16.7-20.2)	15.7	(14.1-17.5)
Diabetes	Yes	801	23.4	(19.5-27.6)	29.8	(25.0-35.1)	12.2	(9.2-16.0)	20.4	(16.4-25.2)	14.3	(11.3-17.9)
	No	5,095	24.5	(22.5-26.6)	27.3	(25.4-29.4)	14.0	(12.5-15.7)	18.4	(16.7-20.2)	15.8	(14.1-17.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Kentucky, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	922	20.0	(17.0-23.3)	34.5	(30.1-39.1)	10.2	(7.9-13.1)	18.5	(15.4-22.1)	16.9	(13.8-20.4)
	No	4,927	26.0	(23.9-28.1)	27.7	(25.7-29.7)	10.6	(9.1-12.2)	19.4	(17.6-21.2)	16.4	(14.8-18.2)
Asthma	Yes	777	22.4	(17.6-27.9)	29.3	(24.3-34.9)	11.7	(8.3-16.3)	19.3	(14.6-25.1)	17.3	(13.9-21.4)
	No	5,114	25.5	(23.5-27.5)	28.3	(26.4-30.4)	10.4	(9.0-12.0)	19.4	(17.7-21.2)	16.4	(14.7-18.2)
Arthritis	Yes	2,474	21.2	(18.9-23.6)	27.7	(25.0-30.6)	10.5	(8.7-12.7)	22.7	(20.2-25.4)	17.9	(15.8-20.2)
	No	3,409	27.0	(24.5-29.5)	28.6	(26.3-31.1)	10.5	(8.8-12.4)	17.9	(16.0-20.1)	16.0	(14.0-18.2)
Frequent Mental Distress	Yes	851	19.2	(15.4-23.6)	19.4	(15.5-24.0)	12.8	(8.5-18.8)	23.4	(18.5-29.1)	25.2	(21.1-29.8)
	No	4,910	25.3	(23.3-27.4)	28.9	(26.9-31.1)	12.7	(11.3-14.2)	18.4	(16.7-20.2)	14.7	(13.0-16.5)
Serious Psychological Distress	Yes	442	13.6	(9.9-18.4)	18.2	(13.2-24.7)	13.4	(7.7-22.2)	24.1	(18.7-30.6)	30.7	(24.7-37.4)
	No	5,316	26.0	(24.1-28.1)	29.0	(27.1-31.0)	10.2	(8.9-11.8)	18.9	(17.3-20.7)	15.7	(14.1-17.5)
Receiving Treatment for Mental Health Condition	Yes	968	19.3	(15.6-23.6)	19.4	(16.1-23.3)	11.8	(8.4-16.3)	21.9	(18.3-25.9)	27.7	(23.3-32.5)
	No	4,927	26.0	(24.0-28.1)	29.7	(27.7-31.8)	10.4	(9.0-12.0)	19.0	(17.3-20.9)	14.9	(13.3-16.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,837	75.6	(74.0-77.2)	17.1	(15.8-18.5)	1.8	(1.4-2.4)	3.4	(2.6-4.3)	2.1	(1.6-2.8)
Age (years)	18-24	221	64.8	(56.8-72.0)	22.4	(16.3-30.1)	2.2	(0.5-9.6)*	7.2	(4.2-11.9)	3.4	(1.6-7.3)*
	25-34	661	70.0	(65.1-74.6)	20.1	(16.7-24.1)	1.3	(0.6-2.8)*	4.9	(2.4-9.7)*	3.7	(2.2-6.0)
	35-54	2,295	77.2	(74.8-79.4)	16.8	(14.8-19.0)	1.6	(1.0-2.5)	2.5	(1.8-3.4)	1.9	(1.3-3.0)
	55+	2,613	81.2	(79.3-82.9)	13.5	(12.0-15.1)	2.3	(1.7-3.1)	2.1	(1.5-2.9)	1.0	(0.7-1.5)
Sex	Men	1,931	70.3	(67.4-73.0)	20.6	(18.3-23.1)	2.3	(1.4-3.6)	4.2	(2.9-6.0)	2.7	(1.8-4.0)
	Women	3,906	80.6	(78.9-82.1)	13.8	(12.5-15.3)	1.4	(1.0-1.8)	2.6	(2.0-3.4)	1.6	(1.2-2.2)
Race/ Ethnicity	Hispanic or Latino	153	68.5	(56.8-78.2)	23.9	(15.1-35.9)	0.3	(0.1-1.4)*	4.2	(1.9-9.1)*	3.1	(0.6-13.8)*
	Not Hispanic or Latino:	4,299	76.7	(75.0-78.3)	17.7	(16.2-19.2)	1.5	(1.2-2.0)	2.6	(2.0-3.4)	1.5	(1.1-2.1)
	White											
	Black or African American	1,092	74.1	(70.5-77.4)	14.9	(12.3-17.9)	2.3	(1.1-4.5)*	5.4	(3.8-7.7)	3.3	(2.2-5.0)
	All Other Races	245	66.9	(55.1-76.9)	17.1	(10.8-26.1)	5.0	(2.0-11.8)*	8.2	(2.4-24.4)*	2.7	(1.0-7.0)*
Education	<High School	653	69.4	(64.0-74.3)	18.0	(14.1-22.8)	3.4	(2.1-5.5)	5.1	(3.2-7.9)	4.1	(2.5-6.5)
	High School Graduate (or GED)	1,869	73.9	(71.0-76.6)	17.6	(15.4-20.1)	2.6	(1.5-4.4)	3.8	(2.7-5.4)	2.1	(1.4-3.3)
	Some College	1,523	75.9	(72.7-78.8)	18.4	(15.8-21.4)	1.0	(0.5-1.8)*	2.8	(1.8-4.3)	1.9	(1.0-3.5)*
	College Graduate	1,781	78.2	(75.1-81.0)	15.8	(13.6-18.4)	1.1	(0.7-1.7)	2.9	(1.4-6.0)*	1.9	(1.1-3.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	980	73.7 (69.4-77.6)	16.6 (13.4-20.3)	1.9 (1.0-3.3)	5.1 (3.4-7.6)	2.8 (1.6-4.7)				
	\$20,000-\$34,999	1,042	75.5 (71.8-78.9)	17.3 (14.5-20.6)	1.5 (0.8-2.6)	3.3 (2.0-5.3)	2.4 (1.4-4.3)				
	\$35,000-\$49,999	728	75.7 (70.8-80.0)	17.4 (13.6-21.9)	0.8 (0.4-1.8)*	1.8 (0.9-3.5)*	4.3 (2.3-7.8)*				
	\$50,000-\$74,999	762	71.0 (66.7-75.0)	20.0 (16.4-24.1)	2.9 (1.5-5.9)*	4.3 (2.6-6.9)	1.8 (0.9-3.9)*				
	\$75,000 and more	1,202	79.6 (75.8-82.9)	15.4 (12.8-18.4)	0.6 (0.3-1.3)*	3.3 (1.3-8.3)*	1.1 (0.6-2.1)*				
Employment Status	Employed	3,124	74.6 (72.4-76.7)	17.4 (15.7-19.3)	1.8 (1.1-2.9)	3.7 (2.7-5.3)	2.4 (1.7-3.5)				
	Out of work	195	76.9 (67.7-84.2)	17.1 (10.7-26.3)	0.4 (0.1-1.3)*	3.5 (1.4-8.2)*	2.1 (0.7-6.1)*				
	Homemaker	575	76.6 (72.2-80.6)	15.2 (12.0-19.1)	1.2 (0.6-2.5)*	4.7 (2.6-8.4)	2.2 (1.1-4.5)*				
	Student	145	78.0 (69.2-84.9)	17.3 (11.3-25.5)	0.6 (0.1-4.4)*	1.7 (0.5-5.2)*	2.4 (0.6-9.1)*				
	Retired	1,331	77.1 (73.4-80.4)	16.6 (13.6-20.1)	2.4 (1.6-3.5)	2.5 (1.5-4.2)	1.4 (0.8-2.4)				
	Unable to work	436	75.7 (70.2-80.4)	16.8 (12.7-21.8)	3.3 (1.8-5.9)*	3.0 (1.5-5.9)*	1.3 (0.7-2.3)*				
Veteran Status	Yes	726	75.7 (71.2-79.8)	18.5 (14.8-22.9)	2.0 (1.1-3.5)	2.7 (1.5-4.6)	1.1 (0.6-2.1)*				
	No	5,106	75.5 (73.8-77.2)	16.9 (15.5-18.4)	1.8 (1.3-2.5)	3.5 (2.7-4.6)	2.3 (1.7-3.1)				
Diabetes	Yes	732	78.2 (74.3-81.6)	15.3 (12.3-18.9)	2.1 (1.2-3.5)	2.1 (1.2-3.7)	2.3 (1.2-4.3)*				
	No	5,098	75.2 (73.5-76.9)	17.3 (15.8-18.8)	1.8 (1.3-2.5)	3.5 (2.7-4.6)	2.2 (1.6-2.9)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	662	78.0	(72.7-82.6)	16.6	(12.4-22.0)	1.8	(0.9-3.4)*	2.5	(1.5-4.3)	1.1	(0.5-2.3)*
	No	5,117	75.3	(73.6-77.0)	17.2	(15.8-18.7)	1.8	(1.3-2.5)	3.4	(2.6-4.5)	2.3	(1.7-3.0)
Asthma	Yes	574	77.1	(71.1-82.1)	17.8	(12.9-23.9)	1.4	(0.7-2.7)*	1.7	(0.8-3.6)*	2.0	(0.8-4.8)*
	No	5,248	75.3	(73.6-77.0)	17.0	(15.7-18.5)	1.8	(1.3-2.5)	3.6	(2.8-4.6)	2.2	(1.6-2.9)
Arthritis	Yes	1,955	78.9	(76.4-81.2)	15.4	(13.4-17.7)	1.4	(1.0-2.1)	2.6	(1.7-3.8)	1.6	(1.1-2.5)
	No	3,836	74.6	(72.6-76.6)	17.5	(15.9-19.3)	2.0	(1.4-2.8)	3.6	(2.7-4.9)	2.2	(1.6-3.1)
Frequent Mental Distress	Yes	566	73.2	(66.9-78.6)	17.1	(12.5-22.9)	1.0	(0.5-2.0)*	6.8	(4.2-10.9)	2.0	(1.1-3.6)*
	No	5,121	75.8	(74.1-77.5)	17.2	(15.8-18.7)	1.9	(1.4-2.6)	2.9	(2.2-3.9)	2.2	(1.6-2.9)
Serious Psychological Distress	Yes	304	70.1	(62.5-76.7)	20.3	(14.6-27.6)	1.2	(0.5-2.9)*	4.5	(2.4-8.3)*	3.8	(2.3-6.3)
	No	5,230	75.8	(74.1-77.4)	17.0	(15.6-18.5)	1.8	(1.3-2.5)	3.3	(2.5-4.3)	2.1	(1.5-2.8)
Receiving Treatment for Mental Health Condition	Yes	780	80.6	(76.7-83.9)	12.3	(9.6-15.6)	0.7	(0.4-1.5)*	4.4	(2.6-7.5)	2.0	(1.0-3.9)*
	No	5,039	75.2	(73.4-76.8)	17.6	(16.2-19.2)	1.9	(1.4-2.6)	3.2	(2.5-4.2)	2.0	(1.5-2.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,712	34.8	(33.1-36.6)	27.8	(26.2-29.5)	3.7	(3.2-4.4)	20.1	(18.6-21.6)	13.5	(12.4-14.8)
Age (years)												
	18-24	222	27.9	(21.1-35.9)	35.6	(27.7-44.4)	1.2	(0.5-2.9)*	24.0	(17.9-31.3)	11.4	(7.1-17.6)
	25-34	665	31.8	(27.5-36.5)	28.6	(24.2-33.4)	2.5	(1.6-4.1)	23.1	(19.1-27.6)	14.0	(11.1-17.4)
	35-54	2,268	34.0	(31.4-36.6)	27.1	(24.8-29.5)	3.7	(2.8-4.9)	20.8	(18.6-23.3)	14.4	(12.6-16.4)
	55+	2,508	40.1	(37.8-42.4)	25.4	(23.3-27.6)	5.2	(4.3-6.4)	16.1	(14.6-17.9)	13.2	(11.7-14.8)
Sex												
	Men	1,920	35.5	(32.7-38.4)	30.6	(27.8-33.4)	3.4	(2.6-4.5)	18.8	(16.4-21.5)	11.7	(9.9-13.8)
	Women	3,792	34.2	(32.3-36.2)	25.2	(23.5-27.1)	4.0	(3.4-4.8)	21.3	(19.7-23.0)	15.2	(13.9-16.6)
Race/Ethnicity												
	Hispanic or Latino	150	32.2	(23.5-42.4)	25.2	(16.8-35.8)	3.8	(1.6-8.8)*	19.7	(10.9-33.0)	19.1	(10.8-31.4)
	Not Hispanic or Latino:	4,201	33.0	(31.2-34.8)	28.4	(26.7-30.1)	3.5	(2.9-4.2)	21.3	(19.8-22.8)	13.9	(12.7-15.2)
	White											
	Black or African American	1,076	37.5	(33.7-41.6)	26.2	(22.5-30.3)	4.4	(3.0-6.3)	17.3	(14.2-20.9)	14.6	(11.8-17.9)
	All Other Races	235	29.0	(21.6-37.8)	29.0	(19.8-40.4)	4.6	(2.3-9.1)*	22.9	(13.2-36.8)	14.4	(9.7-20.9)
Education												
	<High School	626	41.7	(35.9-47.8)	27.9	(22.6-33.8)	3.7	(2.4-5.7)	13.8	(9.3-19.8)	12.9	(9.5-17.3)
	High School Graduate (or GED)	1,839	39.4	(36.2-42.6)	28.0	(25.2-31.1)	3.8	(2.9-5.0)	17.6	(15.0-20.6)	11.2	(9.5-13.1)
	Some College	1,501	34.6	(31.5-37.9)	26.4	(23.5-29.6)	4.2	(3.0-5.8)	21.1	(18.4-24.0)	13.6	(11.6-15.9)
	College Graduate	1,735	29.1	(26.3-32.0)	28.5	(25.8-31.5)	3.2	(2.4-4.3)	23.0	(20.7-25.5)	16.2	(13.9-18.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	949	34.5 (30.1-39.2)	27.5 (23.2-32.2)	4.3 (3.1-6.0)	20.2 (15.2-26.4)	13.5 (10.6-17.0)				
	\$20,000-\$34,999	1,024	39.4 (35.4-43.6)	26.9 (23.3-30.8)	3.5 (2.4-5.1)	16.9 (14.0-20.3)	13.2 (10.7-16.2)				
	\$35,000-\$49,999	722	36.6 (32.1-41.5)	23.6 (19.7-27.9)	3.6 (2.2-5.8)	20.1 (16.9-23.8)	16.1 (12.4-20.5)				
	\$50,000-\$74,999	752	33.0 (29.0-37.4)	30.3 (26.4-34.6)	3.3 (2.0-5.2)	20.3 (17.0-24.0)	13.1 (10.5-16.3)				
	\$75,000 and more	1,181	32.8 (29.4-36.4)	29.6 (26.3-33.2)	2.6 (1.7-4.0)	21.7 (19.1-24.6)	13.3 (11.3-15.5)				
Employment Status	Employed	3,078	34.8 (32.5-37.1)	27.3 (25.2-29.5)	3.4 (2.7-4.3)	20.7 (18.9-22.6)	13.8 (12.3-15.5)				
	Out of work	193	23.9 (16.9-32.6)	28.8 (19.4-40.6)	2.3 (0.9-5.7)*	24.5 (15.8-35.9)	20.5 (13.7-29.4)				
	Homemaker	560	36.5 (31.9-41.2)	27.8 (23.3-32.7)	3.3 (2.0-5.3)	20.5 (16.8-24.8)	11.9 (9.5-15.0)				
	Student	145	38.6 (28.3-50.2)	35.6 (26.5-45.8)	2.9 (1.2-7.1)*	14.2 (9.2-21.2)	8.6 (4.2-17.1)*				
	Retired	1,286	36.5 (32.7-40.4)	27.0 (23.8-30.5)	5.8 (4.5-7.6)	18.0 (14.0-22.9)	12.6 (10.1-15.8)				
	Unable to work	417	33.1 (27.6-39.0)	26.4 (21.3-32.1)	4.3 (2.5-7.4)	19.4 (14.3-25.9)	16.8 (12.7-21.9)				
Veteran Status	Yes	713	34.0 (29.8-38.4)	26.5 (22.5-30.9)	5.1 (3.3-7.7)	16.1 (13.1-19.8)	18.3 (14.5-22.9)				
	No	4,994	34.9 (33.1-36.8)	28.0 (26.3-29.9)	3.5 (3.0-4.2)	20.6 (19.0-22.3)	12.9 (11.7-14.1)				
Diabetes	Yes	719	34.5 (30.5-38.7)	26.6 (22.7-30.9)	6.0 (3.8-9.3)	17.7 (14.3-21.6)	15.3 (12.4-18.8)				
	No	4,984	34.9 (33.1-36.8)	27.9 (26.2-29.7)	3.6 (3.0-4.2)	20.3 (18.7-21.9)	13.4 (12.1-14.7)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Louisiana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	638	32.6	(27.8-37.8)	26.6	(21.9-31.8)	4.2	(2.8-6.5)	20.2	(15.0-26.8)	16.3	(12.0-21.9)
	No	5,016	35.0	(33.2-36.9)	27.9	(26.2-29.7)	3.6	(3.1-4.3)	20.1	(18.6-21.7)	13.3	(12.1-14.6)
Asthma	Yes	557	28.4	(23.5-33.8)	28.1	(22.8-34.2)	2.8	(1.8-4.3)	22.4	(18.0-27.5)	18.3	(13.8-23.8)
	No	5,140	35.5	(33.7-37.4)	27.8	(26.1-29.5)	3.8	(3.2-4.5)	19.8	(18.3-21.5)	13.0	(11.9-14.3)
Arthritis	Yes	1,890	34.1	(31.4-36.8)	28.1	(25.5-30.8)	4.3	(3.4-5.5)	18.8	(16.6-21.2)	14.7	(12.9-16.8)
	No	3,783	35.2	(33.1-37.3)	27.6	(25.6-29.6)	3.5	(2.9-4.3)	20.6	(18.8-22.4)	13.2	(11.8-14.7)
Frequent Mental Distress	Yes	544	32.5	(27.1-38.4)	23.1	(18.3-28.6)	3.9	(2.4-6.4)	17.7	(13.9-22.3)	22.8	(17.9-28.4)
	No	5,026	35.2	(33.4-37.0)	28.2	(26.5-30.0)	3.8	(3.2-4.5)	20.3	(18.7-22.0)	12.5	(11.4-13.7)
Serious Psychological Distress	Yes	301	32.0	(25.7-38.9)	28.3	(21.8-35.8)	3.6	(1.9-6.5)*	15.0	(10.5-21.0)	21.2	(15.5-28.2)
	No	5,132	34.9	(33.1-36.7)	27.5	(25.8-29.3)	3.7	(3.1-4.4)	20.6	(19.0-22.2)	13.3	(12.1-14.6)
Receiving Treatment for Mental Health Condition	Yes	758	32.8	(28.8-37.1)	27.2	(23.2-31.6)	3.1	(2.1-4.6)	18.8	(15.7-22.4)	18.1	(14.7-22.0)
	No	4,938	35.1	(33.3-37.0)	27.9	(26.1-29.7)	3.8	(3.2-4.5)	20.3	(18.7-22.0)	12.9	(11.7-14.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,734	70.0	(68.0-72.0)	23.8	(22.0-25.7)	1.0	(0.7-1.5)	3.4	(2.6-4.3)	1.8	(1.0-3.0)
Age (years)	18-24	118	64.8	(54.8-73.7)	26.9	(18.9-36.6)	1.0	(0.1-6.7)*	4.2	(1.6-10.8)*	3.1	(0.6-13.7)*
	25-34	348	60.9	(55.2-66.2)	34.1	(28.9-39.7)	1.2	(0.4-3.3)*	1.7	(0.8-3.7)*	2.2	(0.9-5.2)*
	35-54	1,398	71.3	(68.5-73.9)	22.3	(19.9-24.9)	0.8	(0.5-1.6)*	4.1	(2.9-5.6)	1.5	(0.8-2.7)*
	55+	1,848	75.1	(72.7-77.4)	19.5	(17.5-21.7)	1.2	(0.8-1.9)	3.0	(2.2-4.2)	1.1	(0.4-3.2)*
Sex	Men	1,392	64.6	(61.4-67.6)	27.3	(24.5-30.3)	1.2	(0.7-1.9)	4.5	(3.3-6.2)	2.4	(1.4-4.1)
	Women	2,342	75.1	(72.5-77.5)	20.5	(18.3-22.9)	0.9	(0.5-1.7)*	2.3	(1.5-3.5)	1.2	(0.4-3.6)*
Race/ Ethnicity	Hispanic or Latino	30	--	--	--	--	--	--	--	--	--	--
	Not Hispanic or Latino:	3,547	71.2	(69.2-73.1)	24.0	(22.1-25.9)	0.9	(0.6-1.2)	2.8	(2.2-3.7)	1.2	(0.6-2.1)
	White											
	Black or African American	15	--	--	--	--	--	--	--	--	--	--
	All Other Races	113	60.1	(48.7-70.6)	21.2	(13.7-31.3)	4.5	(1.0-17.5)*	7.6	(2.8-19.3)*	6.6	(2.0-19.5)*
Education	<High School	230	53.8	(44.9-62.5)	29.1	(21.3-38.4)	1.3	(0.3-4.8)*	5.0	(1.6-14.0)*	10.8	(4.9-22.0)*
	High School Graduate (or GED)	1,123	65.4	(61.6-69.0)	26.9	(23.6-30.4)	1.4	(0.7-2.9)*	4.8	(3.3-7.0)	1.5	(0.6-3.6)*
	Some College	973	68.5	(64.3-72.3)	26.3	(22.6-30.3)	0.8	(0.4-1.5)*	2.9	(1.9-4.4)	1.6	(0.7-3.2)*
	College Graduate	1,405	78.5	(75.8-81.1)	17.8	(15.4-20.4)	0.8	(0.5-1.3)	1.9	(1.2-2.9)	1.1	(0.5-2.4)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	645	62.7 (57.5-67.5)	28.3 (23.8-33.3)	1.0 (0.3-2.8)*	5.0 (3.2-7.5)	3.1 (1.3-7.5)*				
	\$20,000-\$34,999	753	67.0 (62.1-71.7)	25.4 (21.5-29.8)	1.2 (0.6-2.4)*	3.1 (1.8-5.3)	3.2 (1.0-9.7)*				
	\$35,000-\$49,999	597	67.0 (61.9-71.7)	26.5 (22.0-31.6)	0.9 (0.4-1.9)*	3.9 (2.3-6.4)	1.7 (0.8-3.7)*				
	\$50,000-\$74,999	594	74.3 (69.6-78.5)	20.3 (16.7-24.5)	0.4 (0.1-1.2)*	4.4 (2.3-8.3)*	0.5 (0.1-2.0)*				
	\$75,000 and more	736	79.7 (75.7-83.2)	17.8 (14.5-21.7)	0.4 (0.2-1.1)*	1.3 (0.6-3.1)*	0.8 (0.2-2.5)*				
Employment Status	Employed	2,153	71.0 (68.5-73.4)	23.5 (21.3-26.0)	1.2 (0.7-2.1)	2.9 (2.1-4.0)	1.4 (0.8-2.3)				
	Out of work	147	70.3 (58.3-80.1)	17.0 (10.9-25.6)	0.7 (0.1-4.9)*	4.4 (1.2-14.3)*	7.6 (1.9-25.9)*				
	Homemaker	251	66.3 (59.3-72.6)	30.4 (24.2-37.3)	1.3 (0.4-4.4)*	2.1 (0.8-5.3)*	0.0 (0.0-0.0)				
	Student	55	66.4 (51.0-79.0)	30.4 (18.7-45.3)	0.0 (0.0-0.0)	3.2 (0.4-19.6)*	0.0 (0.0-0.0)				
	Retired	864	67.8 (63.9-71.6)	25.5 (22.0-29.3)	0.9 (0.5-1.8)*	3.8 (2.4-5.9)	2.0 (0.5-6.8)*				
	Unable to work	260	72.3 (65.2-78.5)	18.1 (13.2-24.5)	0.4 (0.1-1.8)*	6.9 (4.0-11.8)	2.1 (0.8-5.6)*				
Veteran Status	Yes	543	72.8 (68.2-77.0)	21.8 (18.0-26.1)	0.9 (0.4-1.9)*	3.5 (2.1-6.0)	1.0 (0.5-2.1)*				
	No	3,189	69.5 (67.2-71.6)	24.1 (22.2-26.3)	1.0 (0.7-1.6)	3.4 (2.5-4.5)	1.9 (1.1-3.4)				
Diabetes	Yes	332	72.9 (66.8-78.3)	20.8 (16.0-26.7)	0.7 (0.3-2.0)*	4.3 (2.2-8.4)*	1.2 (0.3-4.5)*				
	No	3,401	69.7 (67.6-71.8)	24.0 (22.1-26.0)	1.1 (0.7-1.6)	3.3 (2.5-4.3)	1.9 (1.1-3.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	371	71.5	(65.9-76.5)	21.7	(17.1-27.0)	0.9	(0.3-2.6)*	4.6	(2.5-8.1)	1.3	(0.4-3.9)*
	No	3,325	69.9	(67.8-72.0)	24.0	(22.1-26.0)	1.0	(0.7-1.6)	3.3	(2.5-4.4)	1.7	(0.9-3.1)*
Asthma	Yes	530	73.5	(67.9-78.4)	23.2	(18.4-28.7)	0.8	(0.4-1.6)*	1.7	(0.8-3.2)*	0.9	(0.3-2.6)*
	No	3,189	69.3	(67.1-71.3)	23.9	(21.9-25.9)	1.1	(0.7-1.7)	3.8	(2.9-4.9)	2.0	(1.2-3.5)
Arthritis	Yes	1,458	71.9	(69.0-74.6)	22.2	(19.6-24.9)	1.0	(0.5-1.7)	3.5	(2.5-5.0)	1.5	(0.8-2.6)
	No	2,268	69.2	(66.6-71.7)	24.4	(22.1-26.9)	1.0	(0.6-1.8)	3.3	(2.4-4.6)	1.9	(1.0-3.8)*
Frequent Mental Distress	Yes	371	65.1	(59.1-70.6)	26.7	(21.7-32.5)	1.7	(0.3-7.9)*	4.4	(2.4-7.8)	2.1	(0.8-5.1)*
	No	3,316	70.4	(68.3-72.5)	23.6	(21.6-25.6)	1.0	(0.7-1.4)	3.3	(2.5-4.3)	1.8	(1.0-3.2)*
Serious Psychological Distress	Yes	133	58.1	(46.3-69.1)	31.3	(21.9-42.5)	1.8	(0.4-8.4)*	2.9	(1.3-6.3)*	5.9	(2.8-12.2)*
	No	3,540	70.8	(68.7-72.8)	23.2	(21.4-25.1)	1.0	(0.7-1.6)	3.5	(2.7-4.5)	1.5	(0.8-2.8)*
Receiving Treatment for Mental Health Condition	Yes	628	76.1	(71.6-80.1)	17.6	(14.1-21.7)	1.5	(0.5-4.9)*	3.4	(1.9-6.0)	1.3	(0.5-3.2)*
	No	3,097	69.2	(66.9-71.3)	24.7	(22.7-26.8)	0.9	(0.6-1.3)	3.4	(2.5-4.5)	1.9	(1.0-3.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,734	23.9	(22.2-25.7)	38.1	(36.1-40.2)	2.1	(1.6-2.6)	26.7	(24.9-28.5)	9.2	(8.2-10.4)
Age (years)	18-24	117	24.2	(17.0-33.1)	43.4	(33.7-53.7)	0.7	(0.1-4.8)*	28.1	(19.8-38.3)	3.6	(1.3-9.6)*
	25-34	351	14.3	(10.5-19.1)	46.4	(40.8-52.1)	2.4	(1.2-4.5)*	28.6	(23.9-33.8)	8.4	(5.8-12.0)
	35-54	1,407	20.5	(18.0-23.3)	39.3	(36.3-42.2)	2.1	(1.4-3.0)	27.3	(24.9-29.9)	10.8	(9.1-12.8)
	55+	1,837	32.1	(29.6-34.7)	31.6	(29.2-34.1)	2.5	(1.8-3.4)	24.1	(22.0-26.4)	9.6	(8.1-11.4)
Sex	Men	1,402	27.0	(24.2-30.0)	41.8	(38.6-45.0)	1.8	(1.2-2.6)	22.2	(19.7-24.9)	7.2	(5.8-9.0)
	Women	2,332	21.1	(19.0-23.3)	34.7	(32.1-37.3)	2.3	(1.7-3.1)	30.9	(28.5-33.5)	11.0	(9.6-12.6)
Race/ Ethnicity	Hispanic or Latino	32	--	--	--	--	--	--	--	--	--	--
	Not Hispanic or Latino:	3,533	21.6	(19.9-23.3)	39.1	(37.1-41.2)	2.1	(1.7-2.7)	28.4	(26.5-30.3)	8.8	(7.8-10.0)
	White											
	Black or African American	16	--	--	--	--	--	--	--	--	--	--
	All Other Races	121	13.4	(7.2-23.5)*	36.2	(25.6-48.4)	3.1	(1.1-7.9)*	29.5	(21.1-39.5)	17.8	(10.0-29.8)
Education	<High School	237	34.1	(26.9-42.1)	38.1	(29.1-48.1)	2.2	(0.9-5.2)*	19.3	(12.5-28.6)	6.3	(3.8-10.3)
	High School Graduate (or GED)	1,136	27.5	(24.2-31.0)	38.3	(34.6-42.1)	2.1	(1.4-3.1)	23.0	(19.9-26.4)	9.2	(7.3-11.6)
	Some College	975	23.4	(20.0-27.1)	37.6	(33.6-41.9)	1.8	(1.0-3.2)	26.4	(23.0-30.1)	10.8	(8.7-13.3)
	College Graduate	1,383	17.5	(15.1-20.1)	38.9	(35.9-42.1)	2.3	(1.6-3.3)	32.4	(29.5-35.4)	8.9	(7.4-10.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	648	27.5	(23.2-32.3)	29.4	(24.9-34.4)	0.8	(0.4-1.8)*	26.6	(22.0-31.7)	15.7	(12.2-19.9)
	\$20,000-\$34,999	752	26.0	(22.1-30.2)	37.1	(32.4-42.0)	2.3	(1.4-3.7)	24.5	(20.7-28.7)	10.2	(7.9-13.1)
	\$35,000-\$49,999	595	18.7	(15.0-23.2)	40.7	(35.6-45.9)	2.5	(1.3-4.5)*	30.0	(25.5-34.8)	8.2	(6.0-11.0)
	\$50,000-\$74,999	597	20.1	(16.5-24.3)	43.2	(38.5-48.1)	2.2	(1.3-3.8)	27.2	(23.3-31.5)	7.3	(5.3-9.9)
	\$75,000 and more	732	23.9	(20.2-28.1)	41.0	(36.8-45.3)	2.0	(1.2-3.2)	26.0	(22.6-29.7)	7.2	(5.2-9.8)
Employment Status	Employed	2,155	23.8	(21.5-26.3)	38.8	(36.2-41.4)	2.1	(1.6-2.9)	27.1	(24.9-29.5)	8.2	(7.0-9.6)
	Out of work	147	19.1	(12.2-28.5)	40.7	(29.4-53.1)	1.6	(0.5-5.0)*	31.2	(22.1-41.9)	7.5	(4.0-13.7)*
	Homemaker	250	23.0	(17.8-29.2)	41.0	(34.3-48.0)	2.3	(1.0-5.2)*	24.1	(19.0-30.1)	9.6	(6.1-14.8)
	Student	56	17.6	(7.6-35.7)*	40.3	(24.9-57.8)	3.3	(0.9-10.6)*	30.0	(17.5-46.5)	8.8	(2.3-28.0)*
	Retired	860	29.4	(26.0-33.1)	37.0	(33.3-40.9)	2.3	(1.5-3.7)	23.5	(20.4-26.9)	7.7	(5.6-10.6)
	Unable to work	264	19.2	(14.2-25.5)	28.1	(21.1-36.3)	0.8	(0.2-3.1)*	27.4	(20.6-35.5)	24.5	(18.9-31.0)
Veteran Status	Yes	547	26.2	(22.4-30.4)	38.8	(34.0-43.9)	1.5	(0.7-2.8)*	24.4	(20.4-28.8)	9.2	(6.4-13.1)
	No	3,185	23.4	(21.5-25.5)	38.0	(35.8-40.3)	2.2	(1.7-2.8)	27.1	(25.1-29.1)	9.3	(8.1-10.5)
Diabetes	Yes	338	28.5	(23.3-34.3)	28.1	(22.7-34.3)	1.9	(0.9-4.1)*	29.8	(24.1-36.1)	11.7	(8.6-15.7)
	No	3,395	23.5	(21.7-25.4)	38.9	(36.7-41.1)	2.1	(1.6-2.7)	26.5	(24.7-28.4)	9.0	(7.9-10.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Maine, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	380	28.8	(24.0-34.1)	31.3	(26.2-36.9)	1.5	(0.6-3.6)*	24.2	(19.3-29.9)	14.2	(10.3-19.2)
	No	3,314	23.3	(21.5-25.3)	38.8	(36.6-41.0)	2.1	(1.7-2.7)	26.9	(25.1-28.9)	8.8	(7.7-10.0)
Asthma	Yes	527	23.2	(18.8-28.3)	32.9	(27.7-38.5)	1.2	(0.6-2.5)*	29.6	(24.6-35.2)	13.1	(10.2-16.8)
	No	3,191	24.1	(22.2-26.0)	38.9	(36.8-41.2)	2.2	(1.7-2.8)	26.3	(24.4-28.2)	8.5	(7.4-9.8)
Arthritis	Yes	1,456	25.7	(23.2-28.4)	35.5	(32.6-38.6)	1.7	(1.1-2.6)	26.1	(23.4-29.0)	11.0	(9.3-12.8)
	No	2,269	22.7	(20.5-25.1)	39.2	(36.6-41.9)	2.3	(1.7-3.0)	27.2	(25.0-29.6)	8.6	(7.3-10.1)
Frequent Mental Distress	Yes	366	15.7	(11.7-20.9)	29.5	(24.0-35.6)	1.4	(0.5-4.1)*	33.8	(28.1-40.1)	19.6	(15.5-24.4)
	No	3,321	24.8	(22.9-26.8)	38.9	(36.8-41.2)	2.1	(1.7-2.7)	26.0	(24.1-27.9)	8.2	(7.1-9.4)
Serious Psychological Distress	Yes	130	14.4	(9.1-22.1)	21.3	(13.8-31.5)	0.6	(0.1-4.0)*	34.3	(23.8-46.5)	29.4	(20.9-39.6)
	No	3,538	24.2	(22.4-26.1)	38.7	(36.7-40.9)	2.2	(1.7-2.7)	26.6	(24.8-28.5)	8.3	(7.3-9.4)
Receiving Treatment for Mental Health Condition	Yes	622	19.1	(15.4-23.4)	31.5	(26.9-36.4)	1.9	(1.0-3.7)*	32.6	(28.2-37.2)	15.0	(12.2-18.3)
	No	3,103	24.7	(22.8-26.8)	39.1	(36.9-41.4)	2.1	(1.6-2.7)	25.8	(23.8-27.8)	8.3	(7.1-9.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,552	68.9	(66.6-71.0)	23.7	(21.7-25.8)	2.8	(2.0-3.9)	3.4	(2.5-4.6)	1.2	(0.8-1.7)
Age (years)	18-24	146	57.9	(45.3-69.5)	39.1	(27.9-51.6)	0.2	(0.0-0.6)*	2.3	(0.8-6.7)*	0.5	(0.2-1.5)*
	25-34	393	65.2	(58.2-71.6)	24.1	(18.5-30.6)	5.3	(2.7-10.4)*	5.1	(2.1-11.6)*	0.3	(0.2-0.7)*
	35-54	1,734	70.3	(67.0-73.5)	22.0	(19.2-25.1)	3.0	(1.9-4.7)	3.4	(2.3-5.1)	1.2	(0.7-2.0)
	55+	2,222	72.3	(69.5-75.0)	20.3	(17.9-23.0)	2.3	(1.5-3.4)	2.7	(1.9-4.0)	2.3	(1.4-3.8)
Sex	Men	1,706	62.5	(58.8-66.0)	29.2	(25.9-32.7)	3.5	(2.2-5.5)	3.8	(2.4-5.9)	1.1	(0.6-1.8)
	Women	2,846	74.9	(72.3-77.3)	18.6	(16.5-21.0)	2.2	(1.5-3.4)	3.0	(2.1-4.3)	1.3	(0.8-2.1)
Race/ Ethnicity	Hispanic or Latino	356	54.3	(44.1-64.1)	20.7	(14.1-29.3)	7.0	(3.6-13.0)*	11.7	(6.0-21.6)*	6.3	(4.4-9.0)
	Not Hispanic or Latino:	3,735	69.3	(66.9-71.6)	24.0	(21.8-26.3)	2.3	(1.6-3.2)	3.2	(2.5-4.1)	1.2	(0.9-1.7)
	White											
	Black or African American	237	50.9	(40.0-61.6)	18.7	(11.7-28.6)	25.0	(18.1-33.5)	4.7	(1.7-12.5)*	0.7	(0.3-1.8)*
	All Other Races	171	68.9	(56.2-79.3)	24.3	(15.4-36.2)	2.4	(0.7-7.7)*	2.6	(0.8-7.9)*	1.8	(0.4-7.2)*
Education	<High School	424	59.2	(49.9-67.8)	29.8	(22.0-39.0)	4.4	(1.9-9.9)*	4.5	(1.9-10.1)*	2.1	(1.2-3.8)
	High School Graduate (or GED)	1,094	66.4	(61.4-71.0)	25.1	(20.9-29.8)	3.5	(1.9-6.6)*	3.2	(2.0-4.9)	1.8	(1.0-3.4)*
	Some College	1,121	70.3	(65.8-74.5)	22.8	(19.0-27.0)	2.4	(1.2-4.6)*	3.5	(1.8-6.8)*	1.0	(0.5-1.9)*
	College Graduate	1,904	72.7	(69.5-75.7)	21.3	(18.6-24.3)	2.6	(1.4-4.8)*	2.8	(1.8-4.4)	0.6	(0.3-1.3)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	701	64.0 (57.3-70.1)	24.0 (19.2-29.6)	3.6 (1.5-8.6)*	5.0 (2.5-9.6)*	3.4 (2.0-5.8)				
	\$20,000-\$34,999	734	63.9 (57.8-69.6)	25.1 (19.9-31.0)	3.6 (1.7-7.5)*	6.9 (3.6-12.6)*	0.6 (0.3-1.2)*				
	\$35,000-\$49,999	489	66.8 (60.0-73.1)	26.0 (20.2-32.8)	2.2 (0.8-6.2)*	3.8 (1.9-7.5)*	1.1 (0.3-4.0)*				
	\$50,000-\$74,999	632	72.0 (66.4-76.9)	23.0 (18.4-28.5)	2.4 (0.9-6.2)*	1.9 (0.8-4.4)*	0.7 (0.2-2.4)*				
	\$75,000 and more	1,337	72.6 (68.9-76.0)	22.0 (18.8-25.6)	2.5 (1.4-4.2)	2.3 (1.4-3.9)	0.6 (0.2-1.6)*				
Employment Status	Employed	2,513	69.3 (66.4-72.1)	23.1 (20.6-25.9)	3.0 (1.9-4.6)	3.6 (2.4-5.4)	1.0 (0.5-1.7)				
	Out of work	322	67.5 (59.0-75.1)	24.5 (18.0-32.5)	2.8 (0.9-8.4)*	3.6 (1.3-9.1)*	1.6 (0.7-3.9)*				
	Homemaker	195	69.7 (59.0-78.6)	23.7 (15.8-34.0)	3.3 (0.9-11.8)*	1.7 (0.3-8.0)*	1.7 (0.6-4.6)*				
	Student	70	72.9 (56.4-84.8)	26.2 (14.5-42.7)	0.0 (0.0-0.0)	0.2 (0.0-1.2)*	0.7 (0.1-5.2)*				
	Retired	1,089	64.6 (60.0-68.8)	28.3 (24.2-32.8)	2.9 (1.7-5.0)	3.3 (1.9-5.5)	0.9 (0.3-2.5)*				
	Unable to work	352	75.8 (67.5-82.6)	15.3 (9.7-23.2)	2.1 (0.6-6.7)*	3.9 (1.7-8.7)*	2.8 (1.2-6.4)*				
Veteran Status	Yes	512	65.3 (59.2-71.0)	28.6 (23.2-34.7)	2.6 (1.1-5.9)*	2.8 (1.5-5.1)*	0.8 (0.3-1.7)*				
	No	4,036	69.2 (66.8-71.6)	23.2 (21.1-25.4)	2.8 (2.0-4.0)	3.5 (2.5-4.8)	1.3 (0.9-1.9)				
Diabetes	Yes	504	71.1 (64.6-76.9)	19.6 (14.7-25.6)	2.6 (1.2-5.4)*	4.2 (2.0-8.4)*	2.6 (1.3-5.2)*				
	No	4,045	68.7 (66.4-71.0)	24.1 (22.0-26.4)	2.8 (1.9-4.0)	3.3 (2.4-4.6)	1.0 (0.7-1.6)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	448	69.1 (62.5-75.0)	22.8 (17.6-28.9)	3.6 (1.6-7.9)*	3.1 (1.5-6.1)*	1.4 (0.7-3.0)*				
	No	4,068	68.8 (66.4-71.1)	23.8 (21.7-26.0)	2.8 (1.9-3.9)	3.4 (2.5-4.7)	1.2 (0.8-1.8)				
Asthma	Yes	736	72.0 (66.4-76.9)	20.3 (15.9-25.7)	2.5 (1.2-5.1)*	3.7 (2.2-6.3)	1.5 (0.7-3.2)*				
	No	3,800	68.2 (65.7-70.6)	24.4 (22.3-26.8)	2.9 (2.0-4.2)	3.4 (2.4-4.7)	1.2 (0.8-1.7)				
Arthritis	Yes	1,589	67.9 (64.1-71.4)	23.4 (20.1-26.9)	3.7 (2.2-6.1)	3.8 (2.4-5.9)	1.4 (0.7-2.4)*				
	No	2,952	69.1 (66.4-71.7)	23.9 (21.5-26.5)	2.5 (1.7-3.8)	3.3 (2.2-4.7)	1.2 (0.7-1.8)				
Frequent Mental Distress	Yes	503	65.1 (58.4-71.3)	23.3 (17.7-30.0)	2.9 (1.1-7.3)*	5.1 (2.5-10.1)*	3.6 (1.9-6.8)*				
	No	3,976	69.1 (66.7-71.4)	24.0 (21.9-26.2)	2.8 (2.0-4.0)	3.2 (2.3-4.5)	0.9 (0.6-1.3)				
Serious Psychological Distress	Yes	215	54.0 (43.5-64.2)	32.6 (22.8-44.2)	3.8 (0.7-18.8)*	5.0 (2.3-10.7)*	4.6 (2.1-9.8)*				
	No	4,257	69.6 (67.3-71.9)	23.6 (21.5-25.7)	2.6 (1.9-3.6)	3.3 (2.4-4.6)	0.9 (0.6-1.4)				
Receiving Treatment for Mental Health Condition	Yes	726	76.3 (71.3-80.6)	18.3 (14.4-23.0)	1.4 (0.5-3.7)*	2.2 (1.1-4.4)*	1.9 (0.9-3.7)*				
	No	3,807	67.8 (65.4-70.2)	24.5 (22.3-26.8)	3.0 (2.1-4.2)	3.6 (2.6-4.9)	1.1 (0.7-1.7)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,535	25.1	(23.1-27.2)	36.4	(34.2-38.7)	4.3	(3.5-5.4)	24.1	(22.3-26.1)	10.0	(8.8-11.4)
Age (years)	18-24	146	23.8	(15.0-35.6)	41.7	(30.1-54.4)	1.9	(0.5-6.9)*	26.9	(17.4-39.3)	5.7	(2.2-14.0)*
	25-34	392	21.4	(15.8-28.3)	40.9	(33.8-48.4)	5.7	(3.0-10.4)*	20.1	(15.3-25.9)	12.0	(8.2-17.2)
	35-54	1,736	21.6	(18.8-24.8)	37.7	(34.4-41.2)	4.4	(3.1-6.1)	25.5	(22.7-28.5)	10.7	(8.8-13.0)
	55+	2,207	33.1	(30.4-35.9)	31.1	(28.3-33.9)	4.4	(3.3-5.8)	21.9	(19.6-24.4)	9.6	(8.0-11.4)
Sex	Men	1,701	26.6	(23.5-29.9)	41.5	(38.0-45.1)	4.8	(3.5-6.5)	19.9	(17.2-22.8)	7.3	(5.7-9.4)
	Women	2,834	23.7	(21.3-26.2)	31.7	(28.9-34.6)	3.9	(2.9-5.3)	28.2	(25.7-30.8)	12.5	(10.8-14.4)
Race/ Ethnicity	Hispanic or Latino	351	26.6	(19.3-35.5)	27.6	(17.6-40.5)	22.3	(16.1-30.0)	15.1	(9.4-23.4)	8.4	(3.3-19.6)*
	Not Hispanic or Latino:	3,719	23.3	(21.3-25.5)	37.0	(34.6-39.5)	4.2	(3.4-5.2)	25.8	(23.7-28.0)	9.7	(8.5-11.2)
	White											
	Black or African American	240	31.8	(22.8-42.3)	31.2	(22.1-42.1)	2.7	(0.8-8.8)*	20.2	(13.5-29.1)	14.1	(8.1-23.3)
	All Other Races	171	31.5	(21.5-43.6)	31.6	(21.7-43.4)	4.1	(1.3-12.4)*	20.3	(12.7-30.9)	12.4	(7.1-20.8)
Education	<High School	412	40.4	(31.2-50.3)	27.7	(19.9-37.2)	4.7	(2.1-10.3)*	18.8	(12.3-27.7)	8.4	(5.2-13.3)
	High School Graduate (or GED)	1,109	30.6	(26.6-35.0)	35.2	(30.6-40.2)	4.2	(2.7-6.6)	21.6	(18.0-25.8)	8.3	(6.2-10.9)
	Some College	1,124	24.3	(20.3-28.7)	35.1	(30.5-40.0)	3.9	(2.6-5.9)	22.6	(19.0-26.6)	14.1	(11.2-17.7)
	College Graduate	1,881	18.1	(15.6-20.9)	39.1	(35.9-42.5)	4.9	(3.5-6.9)	28.6	(25.6-31.7)	9.2	(7.6-11.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	692	36.1 (30.1-42.5)	30.1 (24.4-36.3)	5.0 (2.3-10.7)*	16.5 (12.5-21.6)	12.3 (9.2-16.3)				
	\$20,000-\$34,999	737	28.3 (23.5-33.6)	34.3 (28.3-40.9)	5.7 (3.2-10.0)	21.2 (16.7-26.4)	10.5 (7.5-14.7)				
	\$35,000-\$49,999	482	22.8 (18.0-28.5)	30.6 (24.0-38.1)	7.8 (4.7-12.7)	28.7 (22.6-35.6)	10.1 (6.8-14.8)				
	\$50,000-\$74,999	631	24.1 (19.6-29.2)	41.4 (35.8-47.2)	3.4 (2.0-5.8)	23.6 (19.4-28.4)	7.5 (5.3-10.6)				
	\$75,000 and more	1,334	19.7 (16.5-23.4)	40.8 (37.0-44.7)	3.1 (2.1-4.5)	26.7 (23.5-30.1)	9.8 (7.8-12.2)				
Employment Status	Employed	2,506	22.8 (20.3-25.5)	39.8 (36.8-42.8)	3.9 (2.9-5.2)	23.4 (21.1-25.9)	10.2 (8.6-12.0)				
	Out of work	323	22.9 (16.4-30.9)	29.7 (22.8-37.8)	7.6 (3.6-15.2)*	29.2 (21.7-37.9)	10.7 (6.9-16.1)				
	Homemaker	196	21.4 (14.6-30.2)	34.6 (26.0-44.4)	4.6 (1.9-11.2)*	32.5 (24.5-41.8)	6.8 (3.7-12.3)*				
	Student	70	35.4 (19.7-55.0)	24.6 (12.9-41.7)*	3.4 (0.7-15.7)*	21.9 (11.6-37.5)*	14.7 (5.3-34.7)*				
	Retired	1,081	34.9 (30.9-39.1)	31.9 (27.8-36.3)	5.1 (3.5-7.5)	21.2 (17.8-25.0)	6.8 (5.2-8.9)				
	Unable to work	349	24.3 (17.0-33.3)	24.4 (17.8-32.6)	3.4 (1.5-7.7)*	27.1 (18.8-37.2)	20.8 (14.4-29.1)				
Veteran Status	Yes	503	27.8 (22.8-33.4)	32.7 (27.0-38.9)	4.7 (2.7-8.1)	24.1 (18.7-30.6)	10.7 (7.1-15.8)				
	No	4,027	24.8 (22.7-27.0)	36.8 (34.4-39.3)	4.3 (3.4-5.5)	24.2 (22.2-26.2)	10.0 (8.7-11.4)				
Diabetes	Yes	504	29.0 (22.8-36.0)	35.7 (29.1-42.9)	2.5 (1.3-4.7)*	22.1 (16.8-28.6)	10.7 (7.6-14.9)				
	No	4,028	24.7 (22.7-26.9)	36.5 (34.2-39.0)	4.5 (3.6-5.7)	24.3 (22.3-26.3)	10.0 (8.7-11.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Massachusetts, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	451	28.1	(22.9-33.9)	27.2	(21.5-33.9)	2.4	(1.1-5.3)*	27.2	(21.0-34.6)	15.0	(10.2-21.6)
	No	4,047	24.8	(22.7-27.0)	37.0	(34.6-39.5)	4.5	(3.6-5.7)	24.0	(22.0-26.0)	9.7	(8.4-11.1)
Asthma	Yes	728	23.9	(18.9-29.7)	34.7	(29.1-40.6)	3.6	(2.1-6.1)	26.2	(21.2-31.8)	11.7	(8.5-15.9)
	No	3,792	25.3	(23.2-27.6)	36.8	(34.3-39.3)	4.5	(3.5-5.7)	23.7	(21.7-25.8)	9.7	(8.4-11.1)
Arthritis	Yes	1,582	23.8	(21.0-26.9)	33.3	(29.5-37.3)	4.2	(2.8-6.2)	26.4	(23.1-30.0)	12.3	(10.0-15.0)
	No	2,943	25.6	(23.2-28.2)	37.5	(34.8-40.3)	4.4	(3.4-5.7)	23.3	(21.1-25.6)	9.2	(7.8-10.9)
Frequent Mental Distress	Yes	498	16.9	(12.7-22.2)	27.1	(21.2-34.0)	4.5	(2.3-8.7)*	31.9	(25.3-39.3)	19.6	(14.8-25.4)
	No	3,966	25.9	(23.8-28.2)	37.4	(35.0-39.8)	4.3	(3.4-5.5)	23.4	(21.5-25.5)	8.9	(7.7-10.3)
Serious Psychological Distress	Yes	212	17.1	(10.3-26.9)	27.9	(18.7-39.4)	9.4	(3.6-22.6)*	27.1	(16.4-41.5)	18.5	(11.8-27.8)
	No	4,247	25.4	(23.3-27.5)	37.1	(34.7-39.5)	4.1	(3.3-5.2)	24.1	(22.2-26.1)	9.4	(8.2-10.8)
Receiving Treatment for Mental Health Condition	Yes	714	21.6	(17.2-26.9)	33.4	(27.8-39.5)	2.5	(1.2-5.5)*	28.2	(23.5-33.5)	14.2	(10.9-18.3)
	No	3,806	25.7	(23.6-28.0)	36.3	(33.9-38.8)	5.1	(4.1-6.2)	23.4	(21.4-25.5)	9.5	(8.2-11.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,722	69.5	(67.1-71.8)	23.5	(21.4-25.9)	1.3	(0.8-2.2)	3.8	(2.9-5.1)	1.8	(1.3-2.5)
Age (years)	18-24	109	54.2	(43.9-64.1)	42.9	(33.0-53.3)	0.5	(0.1-3.5)*	2.4	(0.7-8.1)*	0.1	(0.1-0.2)
	25-34	184	67.8	(59.7-75.1)	25.0	(18.4-32.8)	1.2	(0.3-5.1)*	4.6	(1.7-11.9)*	1.4	(0.3-6.0)*
	35-54	926	72.1	(68.3-75.5)	20.5	(17.5-23.9)	1.8	(0.8-3.9)*	4.1	(2.7-6.1)	1.6	(0.9-3.1)*
	55+	1,476	72.4	(69.7-74.9)	19.5	(17.2-21.9)	1.2	(0.7-2.1)	3.6	(2.6-4.9)	3.3	(2.3-4.8)
Sex	Men	1,007	61.8	(57.9-65.6)	29.5	(25.9-33.4)	1.3	(0.6-3.0)*	4.5	(3.0-6.9)	2.8	(1.8-4.2)
	Women	1,715	76.7	(73.9-79.2)	17.9	(15.6-20.4)	1.3	(0.7-2.4)*	3.2	(2.2-4.5)	1.0	(0.5-1.7)
Race/ Ethnicity	Hispanic or Latino	47	--	--	--	--	--	--	--	--	--	--
	Not Hispanic or Latino:	2,082	70.3	(67.7-72.7)	23.1	(20.8-25.6)	1.3	(0.8-2.2)	3.7	(2.8-4.8)	1.6	(1.2-2.2)
	White											
	Black or African American	443	66.2	(58.3-73.2)	22.9	(16.5-30.8)	0.6	(0.3-1.0)	4.9	(2.5-9.3)*	5.5	(3.2-9.2)
	All Other Races	121	57.4	(45.8-68.3)	28.3	(18.7-40.3)	4.1	(1.5-11.0)*	8.0	(2.8-20.5)*	2.2	(1.0-5.0)*
Education	<High School	161	62.1	(50.1-72.7)	28.1	(19.0-39.4)	5.1	(1.0-22.1)*	1.9	(0.7-4.7)*	2.8	(1.3-6.0)*
	High School Graduate (or GED)	838	63.2	(58.7-67.5)	28.6	(24.5-33.1)	0.7	(0.3-1.4)*	5.4	(3.7-8.0)	2.1	(1.2-3.7)
	Some College	791	69.6	(65.1-73.7)	22.9	(19.1-27.1)	1.8	(0.9-3.4)*	3.4	(1.8-6.4)*	2.4	(1.4-4.2)
	College Graduate	926	76.4	(72.4-80.0)	18.9	(15.6-22.6)	1.1	(0.4-3.5)*	3.0	(1.7-5.0)	0.6	(0.3-1.5)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	433	66.8 (59.9-73.1)	25.9 (20.1-32.7)	0.5 (0.2-1.8)*	4.7 (2.5-8.4)*	2.1 (1.1-3.8)*				
	\$20,000-\$34,999	558	67.3 (61.6-72.5)	24.3 (19.6-29.7)	2.2 (0.9-5.4)*	4.1 (2.2-7.4)*	2.1 (1.0-4.3)*				
	\$35,000-\$49,999	376	70.5 (64.1-76.2)	21.5 (16.4-27.7)	2.6 (0.8-7.8)*	4.4 (2.0-9.2)*	1.0 (0.3-3.1)*				
	\$50,000-\$74,999	426	71.4 (65.7-76.4)	25.0 (20.1-30.6)	0.0 (0.0-0.2)*	2.1 (0.9-4.6)*	1.5 (0.7-3.4)*				
	\$75,000 and more	553	73.6 (68.6-78.1)	20.4 (16.3-25.3)	1.2 (0.5-3.2)*	3.3 (1.8-6.1)*	1.4 (0.6-3.2)*				
Employment Status	Employed	1,230	68.5 (65.1-71.8)	23.7 (20.7-26.9)	1.6 (0.8-3.3)*	4.4 (2.9-6.6)	1.8 (1.0-3.2)				
	Out of work	222	72.5 (64.7-79.2)	23.5 (17.2-31.3)	0.5 (0.1-3.1)*	1.9 (0.9-4.2)*	1.6 (0.5-4.7)*				
	Homemaker	192	67.3 (58.3-75.1)	21.9 (15.4-30.2)	3.2 (0.7-12.7)*	6.0 (2.4-14.4)*	1.6 (0.5-5.0)*				
	Student	55	66.8 (52.2-78.8)	33.2 (21.2-47.8)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)				
	Retired	825	69.8 (65.8-73.6)	23.6 (20.0-27.5)	1.1 (0.5-2.4)*	3.5 (2.3-5.4)	2.0 (1.1-3.6)*				
	Unable to work	193	79.0 (70.8-85.4)	12.8 (8.1-19.7)	0.9 (0.3-3.1)*	5.0 (2.4-10.1)*	2.2 (1.0-4.7)*				
Veteran Status	Yes	310	72.2 (65.9-77.7)	22.9 (17.8-29.0)	1.4 (0.5-3.4)*	1.1 (0.4-2.9)*	2.4 (1.2-5.0)*				
	No	2,400	69.1 (66.5-71.6)	23.6 (21.3-26.1)	1.3 (0.7-2.3)	4.2 (3.2-5.7)	1.7 (1.2-2.5)				
Diabetes	Yes	376	74.8 (68.5-80.2)	15.5 (11.2-21.1)	0.6 (0.2-2.0)*	5.7 (2.9-10.9)*	3.4 (1.7-6.4)*				
	No	2,345	69.0 (66.4-71.5)	24.4 (22.1-26.9)	1.4 (0.8-2.4)	3.6 (2.6-4.9)	1.5 (1.0-2.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	339	72.8	(67.1-77.8)	19.5	(15.2-24.7)	2.7	(1.0-7.0)*	2.9	(1.6-5.2)	2.1	(1.0-4.4)*
	No	2,346	69.3	(66.7-71.8)	24.0	(21.7-26.5)	1.0	(0.6-1.7)	4.0	(2.9-5.4)	1.7	(1.2-2.5)
Asthma	Yes	389	71.2	(64.5-77.1)	18.7	(13.9-24.7)	0.4	(0.1-3.0)*	7.2	(3.9-13.0)*	2.3	(1.1-5.1)*
	No	2,324	69.2	(66.7-71.7)	24.3	(22.0-26.8)	1.5	(0.9-2.5)	3.3	(2.4-4.5)	1.6	(1.1-2.4)
Arthritis	Yes	1,171	70.5	(66.9-73.9)	21.8	(18.6-25.3)	0.7	(0.4-1.5)*	4.3	(2.8-6.5)	2.7	(1.7-4.3)
	No	1,534	69.2	(66.1-72.2)	24.3	(21.6-27.3)	1.6	(0.9-2.9)*	3.6	(2.5-5.2)	1.2	(0.7-2.1)
Frequent Mental Distress	Yes	283	69.9	(62.1-76.8)	22.5	(16.4-30.0)	0.6	(0.2-2.1)*	5.9	(2.9-11.5)*	1.1	(0.4-3.4)*
	No	2,403	69.5	(67.0-72.0)	23.8	(21.5-26.2)	1.5	(0.9-2.5)	3.5	(2.6-4.8)	1.7	(1.2-2.5)
Serious Psychological Distress	Yes	92	59.7	(45.3-72.6)	29.3	(18.1-43.7)	1.4	(0.2-9.6)*	6.7	(3.0-14.2)*	2.9	(1.1-7.2)*
	No	2,560	70.4	(67.9-72.8)	22.9	(20.7-25.2)	1.3	(0.7-2.1)	3.7	(2.7-5.0)	1.7	(1.2-2.5)
Receiving Treatment for Mental Health Condition	Yes	392	74.9	(68.5-80.4)	14.8	(10.6-20.5)	2.0	(0.9-4.6)*	7.0	(3.8-12.3)	1.3	(0.6-2.8)*
	No	2,323	68.9	(66.3-71.4)	24.6	(22.3-27.1)	1.2	(0.7-2.2)*	3.4	(2.5-4.7)	1.9	(1.3-2.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,736	23.0	(20.9-25.3)	36.9	(34.5-39.5)	1.4	(1.0-2.1)	27.7	(25.5-30.0)	10.9	(9.5-12.5)
Age (years)	18-24	110	22.8	(15.5-32.4)	45.5	(35.6-55.8)	0.4	(0.1-2.8)*	22.4	(15.1-32.0)	8.8	(4.8-15.8)*
	25-34	185	16.0	(10.5-23.7)	41.0	(32.9-49.7)	0.6	(0.1-2.2)*	31.1	(23.9-39.4)	11.3	(6.8-18.2)
	35-54	926	21.7	(18.5-25.4)	38.5	(34.8-42.5)	1.5	(0.7-2.9)*	28.2	(25.0-31.7)	10.1	(8.1-12.4)
	55+	1,488	28.2	(25.6-31.0)	30.3	(27.5-33.1)	2.0	(1.3-3.1)	26.8	(24.2-29.5)	12.8	(10.9-14.9)
Sex	Men	1,015	25.0	(21.7-28.7)	41.1	(37.2-45.1)	1.2	(0.7-2.3)*	24.0	(20.9-27.5)	8.6	(6.7-11.1)
	Women	1,721	21.2	(18.7-23.9)	33.0	(30.0-36.1)	1.6	(1.0-2.5)	31.2	(28.4-34.2)	13.0	(11.1-15.1)
Race/ Ethnicity	Hispanic or Latino	48	--	--	--	--	--	--	--	--	--	--
	Not Hispanic or Latino:	2,095	21.6	(19.5-23.9)	36.7	(34.1-39.5)	1.6	(1.1-2.4)	29.7	(27.3-32.3)	10.3	(8.8-12.0)
	White											
	Black or African American	442	26.6	(19.1-35.7)	32.7	(24.6-42.0)	1.8	(0.8-4.0)*	20.3	(15.0-26.8)	18.6	(13.9-24.4)
	All Other Races	120	27.8	(18.2-40.0)	38.2	(26.4-51.6)	0.8	(0.6-1.1)	20.0	(12.7-30.2)	13.1	(7.4-22.2)
Education	<High School	166	40.6	(29.5-52.8)	30.6	(20.5-43.1)	2.0	(0.5-7.4)*	20.5	(11.0-34.8)	6.4	(3.2-12.1)*
	High School Graduate (or GED)	851	31.2	(27.1-35.6)	36.5	(32.2-41.1)	1.3	(0.6-2.9)*	19.9	(16.8-23.5)	11.1	(8.6-14.1)
	Some College	792	20.4	(16.5-24.8)	37.7	(33.1-42.5)	1.7	(1.0-2.8)	29.1	(25.0-33.5)	11.2	(8.9-14.0)
	College Graduate	921	14.3	(11.7-17.3)	37.6	(33.6-41.7)	1.4	(0.7-2.9)*	34.9	(31.1-38.9)	11.9	(9.4-14.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	439	33.9 (27.5-40.8)	26.6 (20.8-33.2)	1.5 (0.6-3.6)*	21.0 (15.7-27.6)	17.0 (12.6-22.7)				
	\$20,000-\$34,999	556	25.9 (21.3-31.2)	33.0 (27.7-38.7)	2.1 (0.9-5.0)*	27.9 (23.3-33.1)	11.1 (7.9-15.2)				
	\$35,000-\$49,999	381	21.9 (16.6-28.2)	38.1 (31.5-45.1)	0.3 (0.1-1.2)*	30.2 (24.7-36.3)	9.6 (6.7-13.5)				
	\$50,000-\$74,999	421	16.9 (12.7-22.0)	43.3 (37.4-49.3)	2.4 (1.0-5.7)*	29.3 (24.4-34.6)	8.3 (6.0-11.3)				
	\$75,000 and more	555	18.2 (14.3-22.9)	38.6 (33.7-43.7)	1.0 (0.5-2.0)*	31.2 (26.7-36.1)	11.1 (8.4-14.4)				
Employment Status	Employed	1,221	21.0 (18.0-24.4)	39.9 (36.4-43.5)	0.9 (0.5-1.6)	27.6 (24.7-30.8)	10.5 (8.5-12.9)				
	Out of work	226	24.6 (18.0-32.7)	37.0 (29.1-45.7)	2.0 (0.5-7.5)*	25.3 (19.2-32.5)	11.1 (6.9-17.4)				
	Homemaker	189	26.2 (19.2-34.8)	33.9 (26.2-42.6)	4.1 (1.5-10.8)*	27.2 (20.4-35.2)	8.5 (5.1-13.9)				
	Student	56	14.0 (6.0-29.4)*	40.8 (27.2-56.0)	1.3 (0.2-6.9)*	35.0 (21.8-50.9)	8.9 (3.1-23.1)*				
	Retired	838	27.0 (23.6-30.6)	31.8 (27.8-36.0)	1.8 (1.0-3.3)*	27.7 (24.1-31.7)	11.7 (9.4-14.5)				
	Unable to work	200	27.8 (20.0-37.3)	22.9 (15.6-32.3)	0.6 (0.2-2.6)*	30.1 (19.5-43.3)	18.5 (12.7-26.2)				
Veteran Status	Yes	315	27.1 (22.0-32.9)	32.5 (26.1-39.7)	2.1 (0.9-4.6)*	26.7 (20.9-33.5)	11.5 (7.7-17.0)				
	No	2,408	22.6 (20.3-25.0)	37.4 (34.8-40.2)	1.3 (0.9-2.1)	27.9 (25.6-30.3)	10.7 (9.3-12.4)				
Diabetes	Yes	373	24.0 (19.5-29.2)	27.4 (21.8-33.8)	1.3 (0.5-3.2)*	32.7 (26.2-39.8)	14.7 (10.9-19.5)				
	No	2,362	22.9 (20.7-25.4)	37.8 (35.2-40.5)	1.4 (0.9-2.1)	27.3 (25.0-29.7)	10.5 (9.0-12.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Michigan, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	336	23.2	(18.7-28.3)	31.4	(25.3-38.1)	1.8	(0.7-4.4)*	32.7	(26.5-39.6)	10.9	(7.5-15.7)
	No	2,362	22.8	(20.6-25.2)	37.5	(34.9-40.2)	1.4	(0.9-2.1)	27.4	(25.2-29.8)	10.8	(9.4-12.5)
Asthma	Yes	392	19.4	(14.5-25.4)	34.3	(28.1-41.1)	1.2	(0.4-3.2)*	30.7	(25.1-36.9)	14.5	(10.7-19.3)
	No	2,334	23.6	(21.3-26.0)	37.4	(34.7-40.1)	1.5	(1.0-2.2)	27.3	(25.0-29.7)	10.3	(8.8-12.0)
Arthritis	Yes	1,174	23.4	(20.4-26.8)	32.2	(28.5-36.1)	0.7	(0.4-1.4)*	31.3	(27.8-35.1)	12.3	(10.2-14.7)
	No	1,546	22.7	(20.0-25.7)	39.0	(35.9-42.3)	1.8	(1.1-2.8)	26.1	(23.5-28.9)	10.4	(8.6-12.5)
Frequent Mental Distress	Yes	278	22.7	(16.4-30.7)	31.1	(23.7-39.5)	0.0	(0.0-0.0)	28.7	(21.6-37.0)	17.6	(12.4-24.4)
	No	2,422	23.2	(21.0-25.5)	37.6	(35.0-40.3)	1.6	(1.1-2.4)	27.8	(25.5-30.1)	9.8	(8.5-11.4)
Serious Psychological Distress	Yes	96	19.3	(10.3-33.3)*	32.1	(19.7-47.7)	0.4	(0.1-2.6)*	25.8	(13.1-44.4)*	22.4	(13.4-35.0)
	No	2,571	23.2	(21.1-25.6)	37.1	(34.5-39.7)	1.4	(0.9-2.1)	28.1	(25.9-30.4)	10.3	(8.9-11.8)
Receiving Treatment for Mental Health Condition	Yes	386	22.0	(16.5-28.6)	32.4	(26.3-39.2)	1.2	(0.5-2.8)*	29.9	(23.8-36.8)	14.5	(11.1-18.9)
	No	2,343	23.6	(21.3-25.9)	37.2	(34.6-39.9)	1.6	(1.1-2.3)	27.4	(25.2-29.8)	10.2	(8.8-11.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,669	63.1	(61.1-65.0)	28.7	(26.9-30.6)	3.3	(2.7-4.0)	3.5	(2.9-4.4)	1.3	(0.8-2.1)
Age (years)												
18-24		141	54.9	(45.5-63.9)	38.0	(29.6-47.3)	1.6	(0.5-5.1)*	4.2	(1.9-8.8)*	1.3	(0.2-8.5)*
25-34		461	57.4	(52.1-62.5)	33.9	(28.8-39.4)	3.9	(2.2-7.0)	2.9	(1.5-5.5)*	1.9	(0.7-5.1)*
35-54		1,832	67.6	(65.1-70.0)	25.3	(23.1-27.7)	2.6	(1.9-3.7)	3.2	(2.3-4.3)	1.3	(0.8-2.3)
55+		2,233	63.9	(61.6-66.0)	26.3	(24.3-28.3)	4.7	(3.8-5.8)	4.2	(3.3-5.3)	1.0	(0.6-1.7)
Sex												
Men		1,729	55.9	(52.7-59.0)	34.2	(31.2-37.3)	3.6	(2.7-4.8)	4.1	(3.1-5.5)	2.2	(1.3-3.8)
Women		2,940	70.0	(67.7-72.2)	23.5	(21.4-25.8)	3.1	(2.4-4.0)	3.0	(2.1-4.1)	0.4	(0.2-0.9)*
Race/Ethnicity												
Hispanic or Latino		49	75.4	(51.5-89.8)	14.4	(4.7-36.5)*	3.8	(1.0-13.9)*	0.0	(0.0-0.0)*	6.4	(1.8-20.7)*
Not Hispanic or Latino:		4,388	63.9	(62.0-65.8)	28.9	(27.1-30.8)	3.2	(2.6-4.0)	3.1	(2.5-3.9)	0.8	(0.4-1.5)*
White												
Black or African American		115	54.6	(43.0-65.8)	34.4	(24.3-46.3)	5.6	(2.2-13.4)*	3.2	(1.3-7.7)*	2.2	(0.7-6.1)*
All Other Races		109	57.4	(44.5-69.4)	27.9	(17.5-41.5)	2.6	(0.9-7.3)*	11.2	(5.8-20.7)*	0.8	(0.2-3.4)*
Education												
<High School		247	49.6	(37.8-61.4)	28.5	(20.0-38.9)	15.1	(10.6-21.1)	5.6	(2.3-12.8)*	1.2	(0.3-5.1)*
High School Graduate (or GED)		1,271	56.8	(52.7-60.8)	32.9	(29.2-37.0)	3.2	(2.3-4.6)	4.7	(3.2-6.6)	2.4	(1.1-5.0)*
Some College		1,396	63.5	(59.9-67.0)	28.8	(25.5-32.4)	3.9	(2.7-5.4)	2.7	(1.8-4.1)	1.1	(0.6-2.2)*
College Graduate		1,753	69.9	(67.0-72.6)	25.4	(22.8-28.1)	1.6	(1.1-2.5)	2.5	(1.7-3.7)	0.6	(0.3-1.1)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	529	57.8 (52.0-63.4)	30.2 (24.8-36.1)	5.3 (3.3-8.5)	5.2 (3.5-7.8)	1.5 (0.5-4.0)*				
	\$20,000-\$34,999	894	60.3 (55.3-65.2)	32.5 (28.0-37.4)	2.8 (2.0-4.0)	2.9 (2.0-4.3)	1.4 (0.6-3.7)*				
	\$35,000-\$49,999	776	60.9 (56.6-65.2)	30.0 (25.9-34.5)	4.0 (2.7-6.0)	3.5 (2.0-5.9)	1.5 (0.7-3.4)*				
	\$50,000-\$74,999	796	68.1 (63.8-72.0)	25.3 (21.6-29.6)	1.9 (1.0-3.8)*	3.9 (2.3-6.5)	0.8 (0.3-1.8)*				
	\$75,000 and more	1,382	66.7 (63.3-69.9)	27.9 (24.8-31.2)	1.8 (1.1-3.0)	3.1 (2.0-4.9)	0.5 (0.1-1.5)*				
Employment Status	Employed	2,762	64.9 (62.5-67.3)	28.0 (25.8-30.4)	2.9 (2.1-3.9)	3.3 (2.5-4.4)	0.9 (0.5-1.6)*				
	Out of work	101	64.0 (51.1-75.1)	23.1 (13.1-37.4)	1.9 (0.7-4.7)*	4.1 (1.3-12.0)*	6.9 (2.7-16.4)*				
	Homemaker	223	62.2 (54.5-69.4)	30.3 (23.3-38.3)	5.6 (2.0-14.6)*	1.4 (0.4-4.6)*	0.5 (0.1-3.5)*				
	Student	73	62.6 (48.2-75.2)	31.7 (20.2-45.9)	0.4 (0.1-2.8)*	2.4 (0.6-9.5)*	2.9 (0.4-17.3)*				
	Retired	1,380	57.6 (54.6-60.5)	31.8 (28.9-34.7)	5.3 (4.1-6.8)	4.1 (3.0-5.6)	1.2 (0.6-2.4)*				
	Unable to work	130	61.7 (48.7-73.1)	26.2 (17.3-37.5)	3.9 (1.4-10.1)*	7.8 (4.1-14.3)*	0.4 (0.1-3.1)*				
Veteran Status	Yes	593	60.2 (55.5-64.7)	29.6 (25.5-34.2)	2.6 (1.7-4.0)	6.0 (4.1-8.6)	1.6 (0.8-3.2)*				
	No	4,076	63.4 (61.3-65.5)	28.6 (26.6-30.7)	3.4 (2.8-4.3)	3.2 (2.5-4.2)	1.3 (0.7-2.2)				
Diabetes	Yes	351	55.1 (49.2-60.9)	35.6 (29.8-41.9)	3.7 (2.2-6.3)	5.0 (3.0-8.1)	0.6 (0.2-2.6)*				
	No	4,313	63.5 (61.5-65.6)	28.3 (26.4-30.3)	3.3 (2.7-4.1)	3.5 (2.8-4.3)	1.4 (0.9-2.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	437	58.0	(52.8-63.1)	28.1	(23.6-33.1)	4.1	(2.5-6.5)	7.7	(5.1-11.5)	2.1	(1.0-4.3)*
	No	4,218	63.5	(61.5-65.6)	28.8	(26.8-30.8)	3.2	(2.6-4.0)	3.2	(2.5-4.1)	1.2	(0.7-2.1)
Asthma	Yes	467	64.9	(58.6-70.8)	28.2	(22.5-34.6)	3.8	(2.1-6.7)	2.1	(1.2-3.8)*	1.0	(0.2-4.3)*
	No	4,200	62.8	(60.8-64.9)	28.7	(26.8-30.7)	3.2	(2.6-4.0)	3.8	(3.0-4.7)	1.4	(0.9-2.3)
Arthritis	Yes	1,447	63.6	(60.5-66.5)	27.0	(24.2-30.0)	3.6	(2.5-5.1)	4.1	(3.0-5.6)	1.6	(0.9-3.1)*
	No	3,217	62.9	(60.6-65.3)	29.2	(27.0-31.5)	3.2	(2.5-4.1)	3.4	(2.6-4.4)	1.2	(0.7-2.3)*
Frequent Mental Distress	Yes	310	64.8	(57.2-71.7)	26.3	(19.7-34.2)	1.7	(0.7-3.8)*	5.5	(2.8-10.6)*	1.7	(0.5-5.7)*
	No	4,344	63.1	(61.1-65.1)	28.8	(26.9-30.8)	3.4	(2.8-4.2)	3.4	(2.7-4.2)	1.3	(0.8-2.1)
Serious Psychological Distress	Yes	123	58.5	(46.9-69.2)	32.6	(22.0-45.3)	1.9	(0.6-5.9)*	4.0	(1.7-9.3)*	3.0	(0.6-13.4)*
	No	4,518	63.4	(61.4-65.3)	28.6	(26.7-30.5)	3.3	(2.7-4.1)	3.5	(2.8-4.3)	1.2	(0.8-2.0)
Receiving Treatment for Mental Health Condition	Yes	605	80.4	(75.5-84.6)	14.6	(11.1-19.0)	1.2	(0.6-2.4)*	2.7	(1.1-6.1)*	1.1	(0.3-4.4)*
	No	4,062	61.0	(58.9-63.1)	30.5	(28.5-32.5)	3.6	(2.9-4.4)	3.6	(2.9-4.5)	1.3	(0.8-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,669	21.9	(20.2-23.8)	40.8	(38.9-42.8)	5.1	(4.3-6.0)	23.8	(22.3-25.5)	8.3	(7.4-9.3)
Age (years)	18-24	142	25.6	(18.3-34.6)	44.5	(35.6-53.6)	3.5	(1.5-8.0)*	22.4	(15.6-31.1)	4.1	(1.9-8.5)*
	25-34	459	16.0	(12.2-20.6)	45.9	(40.6-51.2)	5.7	(3.5-9.2)	24.1	(20.2-28.6)	8.3	(5.7-12.0)
	35-54	1,835	21.0	(18.8-23.3)	40.7	(38.3-43.2)	4.5	(3.6-5.7)	24.2	(22.2-26.3)	9.6	(8.2-11.2)
	55+	2,231	24.7	(22.7-26.8)	37.0	(34.8-39.3)	6.3	(5.3-7.5)	23.7	(21.9-25.7)	8.3	(7.1-9.6)
Sex	Men	1,736	24.5	(21.8-27.5)	41.4	(38.4-44.5)	6.1	(4.8-7.8)	20.7	(18.4-23.3)	7.2	(5.9-8.8)
	Women	2,933	19.5	(17.5-21.6)	40.3	(38.0-42.7)	4.2	(3.4-5.0)	26.8	(24.9-28.8)	9.3	(8.1-10.7)
Race/ Ethnicity	Hispanic or Latino	49	32.5	(15.5-55.8)*	38.0	(19.1-61.4)*	4.5	(1.4-13.6)*	22.0	(9.1-44.3)*	3.0	(1.0-8.6)*
	Not Hispanic or Latino:	4,390	18.7	(17.1-20.3)	42.6	(40.6-44.5)	5.2	(4.5-6.2)	25.4	(23.8-27.1)	8.1	(7.2-9.1)
	White											
	Black or African American	113	33.6	(21.7-48.2)	26.9	(18.3-37.6)	12.6	(5.6-25.9)*	16.9	(9.6-27.8)	10.1	(5.8-16.9)
	All Other Races	109	26.9	(16.5-40.7)	35.5	(24.4-48.5)	0.8	(0.1-4.9)*	21.0	(12.4-33.1)	15.8	(8.8-26.7)
Education	<High School	251	24.8	(17.2-34.4)	30.4	(21.4-41.3)	7.9	(4.2-14.5)*	27.9	(19.7-37.9)	9.0	(5.1-15.3)
	High School Graduate (or GED)	1,276	24.6	(21.2-28.4)	41.3	(37.4-45.3)	5.4	(4.0-7.4)	21.4	(18.2-25.0)	7.2	(5.6-9.3)
	Some College	1,393	23.3	(20.0-26.8)	39.5	(36.0-43.1)	4.4	(3.3-5.8)	24.5	(21.7-27.5)	8.4	(6.8-10.3)
	College Graduate	1,747	15.7	(13.5-18.1)	43.5	(40.6-46.4)	5.3	(4.0-6.8)	26.1	(23.8-28.6)	9.5	(7.9-11.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	527	18.8 (14.3-24.2)	32.9 (27.4-38.9)	6.0 (4.0-9.0)	30.1 (24.3-36.5)	12.3 (8.8-17.0)				
	\$20,000-\$34,999	895	20.2 (16.6-24.4)	40.7 (36.1-45.5)	5.7 (4.0-8.2)	25.1 (21.2-29.5)	8.3 (6.4-10.6)				
	\$35,000-\$49,999	777	21.5 (17.9-25.6)	40.7 (36.3-45.3)	5.2 (3.6-7.4)	22.6 (19.5-26.0)	10.0 (7.6-13.1)				
	\$50,000-\$74,999	794	21.0 (17.7-24.6)	39.7 (35.4-44.1)	4.4 (3.0-6.6)	26.1 (22.7-29.9)	8.8 (6.5-11.7)				
	\$75,000 and more	1,382	22.1 (18.9-25.7)	45.4 (42.0-48.8)	3.9 (2.8-5.2)	22.0 (19.6-24.5)	6.6 (5.3-8.2)				
Employment Status	Employed	2,764	21.7 (19.6-23.9)	41.7 (39.3-44.1)	4.6 (3.7-5.7)	23.5 (21.7-25.5)	8.5 (7.4-9.9)				
	Out of work	101	25.9 (16.2-38.8)	42.7 (30.3-56.1)	3.8 (1.5-9.3)*	14.3 (7.6-25.3)*	13.3 (6.6-24.9)*				
	Homemaker	225	18.4 (12.0-27.1)	45.3 (37.8-53.0)	8.9 (5.3-14.8)	21.6 (16.8-27.4)	5.8 (3.6-9.2)				
	Student	74	36.9 (24.2-51.7)	34.5 (22.6-48.7)	3.4 (0.6-18.7)*	20.8 (11.6-34.4)	4.4 (1.0-17.0)*				
	Retired	1,376	20.9 (18.7-23.4)	40.9 (37.9-44.0)	7.4 (6.0-9.1)	24.1 (21.6-26.7)	6.7 (5.4-8.4)				
	Unable to work	129	7.8 (3.8-15.5)*	22.2 (14.4-32.4)	4.0 (1.8-8.8)*	50.2 (37.3-63.0)	15.8 (10.1-23.8)				
Veteran Status	Yes	590	22.4 (18.8-26.4)	36.4 (32.0-41.1)	6.5 (4.6-9.2)	24.4 (20.4-28.8)	10.3 (7.5-14.0)				
	No	4,079	21.8 (19.9-23.9)	41.4 (39.3-43.5)	4.9 (4.1-5.9)	23.7 (22.1-25.5)	8.1 (7.1-9.2)				
Diabetes	Yes	351	23.6 (18.2-30.0)	30.9 (25.5-36.8)	4.9 (3.0-8.0)	30.6 (24.8-37.0)	10.1 (7.1-14.1)				
	No	4,313	21.8 (20.0-23.7)	41.4 (39.4-43.5)	5.1 (4.3-6.1)	23.5 (21.9-25.1)	8.2 (7.2-9.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Minnesota, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	440	20.1	(16.5-24.3)	38.9	(33.8-44.1)	5.6	(3.6-8.4)	22.7	(18.6-27.5)	12.7	(9.4-17.1)
	No	4,215	22.1	(20.3-24.1)	41.0	(39.0-43.0)	5.1	(4.3-6.0)	23.9	(22.2-25.6)	8.0	(7.0-9.0)
Asthma	Yes	462	19.5	(14.6-25.5)	39.0	(33.1-45.4)	3.7	(2.2-6.2)	27.2	(22.3-32.7)	10.6	(7.5-14.7)
	No	4,205	22.2	(20.4-24.2)	41.1	(39.0-43.1)	5.3	(4.4-6.2)	23.4	(21.8-25.2)	8.0	(7.0-9.1)
Arthritis	Yes	1,451	20.3	(17.9-22.9)	36.1	(33.2-39.2)	5.5	(4.3-7.1)	26.8	(24.2-29.6)	11.2	(9.4-13.3)
	No	3,213	22.5	(20.3-24.8)	42.1	(39.8-44.5)	5.0	(4.1-6.1)	23.0	(21.2-25.0)	7.4	(6.3-8.6)
Frequent Mental Distress	Yes	312	15.5	(10.1-22.9)	29.7	(23.2-37.2)	1.7	(0.8-3.5)*	32.7	(25.8-40.5)	20.4	(15.1-27.0)
	No	4,343	22.5	(20.7-24.4)	41.6	(39.6-43.6)	5.4	(4.6-6.3)	23.2	(21.6-24.8)	7.3	(6.5-8.3)
Serious Psychological Distress	Yes	125	10.2	(6.1-16.5)	25.4	(16.0-37.8)	1.2	(0.3-5.0)*	30.5	(21.7-41.1)	32.7	(21.2-46.7)
	No	4,516	22.4	(20.7-24.3)	41.4	(39.4-43.3)	5.1	(4.3-6.0)	23.6	(22.0-25.2)	7.5	(6.7-8.5)
Receiving Treatment for Mental Health Condition	Yes	605	15.0	(11.7-19.0)	37.4	(32.3-42.8)	3.1	(2.0-4.9)	30.5	(25.6-35.9)	14.0	(11.1-17.6)
	No	4,062	22.8	(20.9-24.8)	41.2	(39.2-43.3)	5.4	(4.5-6.4)	23.0	(21.4-24.7)	7.6	(6.6-8.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		9,911	70.4	(68.8-72.1)	23.0	(21.6-24.6)	1.3	(1.0-1.7)	3.3	(2.7-4.0)	1.9	(1.5-2.5)
Age (years)	18-24	313	64.8	(58.2-70.9)	28.5	(22.9-34.9)	0.5	(0.2-1.4)*	4.9	(2.9-8.0)	1.4	(0.7-2.7)*
	25-34	761	64.0	(59.6-68.2)	27.2	(23.3-31.5)	1.1	(0.5-2.4)*	3.4	(2.1-5.6)	4.2	(2.8-6.3)
	35-54	3,095	71.9	(69.7-74.0)	21.0	(19.1-23.0)	1.4	(1.0-2.1)	3.5	(2.7-4.6)	2.1	(1.5-3.1)
	55+	5,694	73.7	(72.1-75.3)	20.8	(19.4-22.3)	1.6	(1.1-2.2)	2.5	(2.0-3.1)	1.4	(1.0-1.9)
Sex	Men	3,374	66.9	(64.3-69.4)	25.6	(23.3-28.1)	1.3	(0.9-2.0)	3.8	(2.9-4.9)	2.4	(1.7-3.4)
	Women	6,537	73.8	(72.1-75.6)	20.5	(19.0-22.2)	1.3	(0.9-1.8)	2.8	(2.2-3.7)	1.5	(1.1-2.1)
Race/ Ethnicity	Hispanic or Latino	157	68.8	(53.0-81.2)	23.6	(13.1-38.8)	1.0	(0.3-3.4)*	6.1	(1.8-19.1)*	0.5	(0.3-0.7)
	Not Hispanic or Latino:	6,702	71.0	(69.3-72.7)	22.9	(21.4-24.6)	1.4	(1.1-1.7)	3.1	(2.5-3.8)	1.6	(1.2-2.0)
	White											
	Black or African American	2,796	63.3	(60.5-66.0)	21.9	(19.5-24.4)	1.2	(0.8-1.9)	5.2	(4.2-6.5)	8.4	(7.3-9.6)
	All Other Races	158	66.4	(54.8-76.4)	22.1	(14.1-32.9)	6.2	(3.6-10.6)	2.7	(0.8-8.8)*	2.5	(0.9-6.6)*
Education	<High School	1,548	64.8	(60.3-69.0)	27.1	(23.2-31.4)	1.1	(0.6-1.8)	4.1	(2.6-6.2)	3.0	(1.9-4.6)
	High School Graduate (or GED)	3,185	65.6	(62.7-68.3)	26.2	(23.7-28.9)	1.4	(1.0-2.2)	4.4	(3.3-5.7)	2.4	(1.7-3.6)
	Some College	2,643	70.2	(67.2-73.0)	23.5	(20.9-26.3)	1.4	(0.9-2.1)	3.2	(2.2-4.4)	1.8	(1.1-2.9)
	College Graduate	2,530	76.6	(73.9-79.1)	19.0	(16.8-21.5)	1.4	(0.7-2.8)*	2.0	(1.2-3.3)	1.0	(0.5-1.9)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	2,563	66.3	(62.8-69.7)	25.1	(22.1-28.4)	1.5	(0.9-2.6)	4.4	(3.2-6.0)	2.7	(1.8-4.0)
	\$20,000-\$34,999	2,014	69.1	(65.5-72.5)	22.9	(19.9-26.2)	1.2	(0.7-2.0)	4.4	(3.1-6.4)	2.4	(1.5-3.8)
	\$35,000-\$49,999	1,238	73.4	(69.6-76.9)	21.7	(18.4-25.3)	1.1	(0.6-1.7)	2.4	(1.5-3.8)	1.4	(0.7-3.0)*
	\$50,000-\$74,999	1,164	70.7	(66.6-74.5)	24.2	(20.6-28.2)	0.8	(0.4-1.6)*	1.8	(1.1-3.2)	2.5	(1.4-4.5)*
	\$75,000 and more	1,601	75.7	(72.6-78.6)	19.5	(16.9-22.4)	1.3	(0.7-2.3)	2.7	(1.6-4.5)	0.8	(0.3-1.9)*
Employment Status	Employed	4,159	70.2	(67.9-72.4)	23.2	(21.2-25.3)	1.2	(0.8-1.8)	3.2	(2.4-4.2)	2.2	(1.5-3.3)
	Out of work	618	74.0	(69.0-78.5)	17.5	(13.9-21.8)	1.3	(0.5-3.2)*	4.5	(2.5-7.8)	2.7	(1.4-4.9)*
	Homemaker	788	69.7	(65.0-74.1)	24.0	(19.8-28.7)	1.7	(0.8-3.6)*	3.1	(1.8-5.3)	1.5	(0.8-2.9)*
	Student	146	76.5	(67.7-83.4)	19.0	(12.7-27.5)	0.4	(0.1-3.1)*	2.8	(1.2-6.4)*	1.3	(0.3-4.7)*
	Retired	2,947	68.9	(66.3-71.5)	25.8	(23.4-28.4)	1.5	(1.0-2.3)	2.7	(1.9-3.7)	1.1	(0.7-1.6)
	Unable to work	1,239	67.0	(62.9-70.9)	25.2	(21.5-29.2)	1.3	(0.7-2.7)*	4.4	(3.0-6.4)	2.1	(1.3-3.4)
Veteran Status	Yes	1,289	70.6	(66.8-74.2)	24.5	(21.1-28.4)	1.5	(1.0-2.4)	2.0	(1.3-3.0)	1.3	(0.7-2.4)*
	No	8,622	70.4	(68.6-72.1)	22.9	(21.3-24.5)	1.2	(0.9-1.7)	3.5	(2.8-4.4)	2.0	(1.5-2.7)
Diabetes	Yes	1,635	72.3	(69.1-75.3)	22.9	(20.1-26.0)	1.2	(0.6-2.2)*	2.5	(1.6-3.9)	1.1	(0.6-2.0)*
	No	8,266	70.3	(68.5-72.0)	23.0	(21.4-24.6)	1.3	(1.0-1.8)	3.4	(2.8-4.2)	2.0	(1.5-2.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	1,442	71.4	(68.0-74.4)	22.4	(19.5-25.6)	1.7	(1.0-2.9)	2.8	(1.8-4.2)	1.8	(1.0-3.1)
	No	8,325	70.4	(68.6-72.1)	23.1	(21.6-24.8)	1.2	(0.9-1.7)	3.4	(2.7-4.1)	1.9	(1.4-2.6)
Asthma	Yes	1,222	70.2	(66.0-74.0)	23.1	(19.5-27.1)	1.2	(0.7-2.2)*	3.5	(2.1-5.7)	2.0	(1.1-3.5)
	No	8,665	70.5	(68.7-72.2)	23.0	(21.5-24.7)	1.3	(0.9-1.8)	3.3	(2.6-4.0)	1.9	(1.4-2.5)
Arthritis	Yes	4,266	71.6	(69.4-73.8)	21.9	(19.9-23.9)	1.8	(1.1-2.7)	3.2	(2.4-4.2)	1.6	(1.1-2.4)
	No	5,596	69.9	(67.9-71.9)	23.5	(21.7-25.4)	1.1	(0.8-1.5)	3.3	(2.6-4.2)	2.2	(1.6-3.0)
Frequent Mental Distress	Yes	1,241	68.5	(64.4-72.4)	21.8	(18.3-25.7)	0.8	(0.3-1.7)*	5.4	(3.5-8.1)	3.5	(2.3-5.5)
	No	8,527	70.7	(68.9-72.4)	23.2	(21.7-24.9)	1.4	(1.0-1.9)	3.0	(2.4-3.7)	1.7	(1.2-2.3)
Serious Psychological Distress	Yes	668	67.5	(61.5-73.0)	23.7	(18.9-29.3)	1.0	(0.4-2.6)*	3.7	(2.2-6.2)	4.1	(2.3-7.1)
	No	9,001	70.8	(69.1-72.4)	23.1	(21.6-24.7)	1.2	(0.9-1.6)	3.2	(2.6-3.9)	1.7	(1.3-2.3)
Receiving Treatment for Mental Health Condition	Yes	1,466	74.5	(71.1-77.7)	20.0	(17.1-23.2)	1.3	(0.6-2.5)*	2.5	(1.6-3.6)	1.8	(1.1-2.8)
	No	8,414	69.9	(68.1-71.7)	23.4	(21.8-25.1)	1.3	(1.0-1.8)	3.4	(2.8-4.2)	1.9	(1.4-2.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		9,964	25.5	(24.1-27.1)	29.0	(27.4-30.6)	2.5	(2.1-3.1)	25.2	(23.7-26.7)	17.8	(16.5-19.2)
Age (years)	18-24	316	18.5	(14.1-23.8)	32.7	(26.5-39.5)	1.0	(0.3-2.9)*	32.2	(26.0-39.1)	15.6	(11.2-21.3)
	25-34	761	22.0	(18.5-26.0)	29.1	(25.2-33.4)	1.8	(1.0-3.3)*	28.0	(24.0-32.3)	19.1	(15.8-22.9)
	35-54	3,109	25.5	(23.5-27.7)	28.8	(26.7-31.0)	2.5	(1.8-3.4)	24.0	(22.1-26.0)	19.2	(17.4-21.2)
	55+	5,732	30.1	(28.5-31.8)	27.8	(26.2-29.4)	3.6	(3.0-4.4)	22.2	(20.8-23.7)	16.3	(15.0-17.7)
Sex	Men	3,377	27.4	(25.2-29.7)	30.6	(28.2-33.1)	2.4	(1.8-3.3)	24.2	(21.9-26.6)	15.5	(13.5-17.6)
	Women	6,587	23.8	(22.2-25.5)	27.4	(25.7-29.2)	2.6	(2.1-3.3)	26.2	(24.5-28.0)	20.0	(18.4-21.7)
Race/ Ethnicity	Hispanic or Latino	155	33.1	(21.5-47.3)	23.5	(13.5-37.6)	2.4	(0.6-9.4)*	28.7	(16.5-44.9)	12.3	(6.3-22.7)*
	Not Hispanic or Latino:	6,764	23.4	(22.0-24.9)	28.8	(27.2-30.6)	2.6	(2.1-3.2)	24.3	(22.7-25.9)	20.8	(19.4-22.4)
	White											
	Black or African American	2,787	28.7	(26.2-31.4)	27.1	(24.4-29.9)	1.7	(1.3-2.4)	22.3	(19.7-25.1)	20.2	(18.0-22.5)
	All Other Races	160	16.5	(8.5-29.7)*	15.9	(8.6-27.6)*	1.1	(0.2-5.9)*	12.8	(6.6-23.5)*	53.7	(41.0-65.9)
Education	<High School	1,543	29.8	(26.0-33.8)	30.0	(25.7-34.7)	2.9	(1.9-4.3)	19.7	(16.2-23.8)	17.6	(14.3-21.4)
	High School Graduate (or GED)	3,225	30.4	(27.9-33.1)	26.3	(23.9-28.9)	2.5	(1.8-3.5)	24.7	(22.2-27.4)	16.0	(14.0-18.2)
	Some College	2,659	26.9	(24.3-29.7)	27.4	(24.7-30.3)	2.1	(1.5-2.9)	23.6	(21.1-26.2)	20.1	(17.6-22.9)
	College Graduate	2,531	19.6	(17.5-21.9)	32.5	(29.8-35.3)	2.9	(2.1-4.1)	27.5	(25.0-30.2)	17.5	(15.5-19.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	2,595	26.5 (23.4-29.7)	26.4 (23.3-29.7)	3.0 (2.0-4.5)	25.2 (22.1-28.7)	18.9 (16.1-22.1)				
	\$20,000-\$34,999	2,037	24.6 (21.9-27.6)	26.4 (23.3-29.8)	3.5 (2.4-5.1)	26.8 (23.5-30.3)	18.7 (15.7-22.0)				
	\$35,000-\$49,999	1,226	28.5 (24.9-32.5)	25.2 (21.8-29.0)	2.0 (1.2-3.4)	27.2 (23.5-31.1)	17.1 (13.9-20.7)				
	\$50,000-\$74,999	1,161	27.1 (23.6-30.8)	29.1 (25.4-33.2)	1.8 (1.1-3.1)	24.5 (21.3-28.1)	17.5 (14.7-20.7)				
	\$75,000 and more	1,601	25.9 (23.0-29.0)	31.7 (28.7-34.8)	2.3 (1.4-3.5)	22.7 (20.1-25.5)	17.5 (15.2-20.1)				
Employment Status	Employed	4,166	26.3 (24.3-28.4)	30.5 (28.4-32.7)	1.9 (1.4-2.6)	24.3 (22.4-26.3)	17.0 (15.3-18.9)				
	Out of work	620	21.9 (17.5-27.0)	22.7 (18.0-28.2)	1.8 (0.8-3.9)*	35.2 (29.0-41.9)	18.5 (14.0-23.9)				
	Homemaker	791	27.2 (23.1-31.8)	26.2 (22.0-31.0)	4.8 (3.1-7.2)	25.6 (21.7-29.9)	16.2 (13.2-19.7)				
	Student	148	23.5 (15.7-33.6)	21.3 (14.1-30.7)	1.3 (0.3-5.4)*	27.0 (18.2-38.0)	27.0 (17.5-39.1)				
	Retired	2,982	28.5 (26.1-30.9)	30.9 (28.4-33.5)	3.9 (2.9-5.2)	22.3 (20.2-24.6)	14.4 (12.6-16.4)				
	Unable to work	1,243	19.5 (16.7-22.7)	27.1 (22.9-31.6)	3.1 (1.7-5.4)	23.7 (20.2-27.6)	26.6 (22.8-30.8)				
Veteran Status	Yes	1,305	26.0 (22.7-29.7)	28.8 (25.2-32.7)	3.6 (2.5-5.2)	21.8 (18.4-25.6)	19.8 (16.3-23.7)				
	No	8,658	25.5 (23.9-27.1)	29.0 (27.3-30.7)	2.3 (1.9-2.9)	25.6 (24.0-27.3)	17.6 (16.2-19.1)				
Diabetes	Yes	1,646	24.4 (21.6-27.4)	25.6 (22.6-28.8)	2.9 (2.0-4.1)	30.3 (26.9-34.0)	16.8 (14.6-19.4)				
	No	8,307	25.7 (24.2-27.4)	29.3 (27.6-31.0)	2.4 (2.0-3.0)	24.6 (23.1-26.2)	17.9 (16.5-19.5)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Mississippi, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	1,475	25.1	(22.3-28.1)	26.6	(23.5-30.0)	2.9	(1.8-4.4)	26.5	(23.2-30.1)	19.0	(16.3-21.9)
	No	8,339	25.7	(24.1-27.3)	29.2	(27.5-30.9)	2.4	(1.9-3.0)	25.1	(23.6-26.7)	17.6	(16.2-19.1)
Asthma	Yes	1,227	22.9	(19.4-26.7)	27.8	(23.7-32.3)	2.8	(1.7-4.5)	25.5	(21.6-29.8)	21.0	(17.6-25.0)
	No	8,712	25.9	(24.3-27.5)	29.2	(27.5-30.8)	2.5	(2.0-3.0)	25.2	(23.6-26.8)	17.3	(15.9-18.8)
Arthritis	Yes	4,294	23.4	(21.5-25.4)	27.6	(25.5-29.8)	3.3	(2.5-4.3)	26.3	(24.2-28.5)	19.4	(17.6-21.4)
	No	5,623	26.5	(24.7-28.5)	29.5	(27.5-31.5)	2.1	(1.6-2.7)	24.7	(22.8-26.6)	17.2	(15.6-19.0)
Frequent Mental Distress	Yes	1,256	18.5	(15.5-22.0)	23.4	(19.7-27.5)	2.3	(1.3-4.1)	26.4	(22.5-30.7)	29.4	(25.2-34.0)
	No	8,563	26.4	(24.8-28.0)	29.8	(28.2-31.5)	2.4	(2.0-3.0)	25.0	(23.5-26.6)	16.3	(15.0-17.7)
Serious Psychological Distress	Yes	662	16.6	(13.1-21.0)	24.2	(19.0-30.2)	2.8	(1.3-5.9)*	19.8	(15.3-25.4)	36.5	(30.0-43.6)
	No	9,065	26.3	(24.8-27.9)	29.2	(27.6-30.9)	2.4	(2.0-3.0)	25.3	(23.8-26.9)	16.7	(15.4-18.1)
Receiving Treatment for Mental Health Condition	Yes	1,470	19.4	(16.5-22.7)	20.7	(17.6-24.1)	2.4	(1.4-3.9)	22.3	(19.3-25.6)	35.3	(31.6-39.1)
	No	8,462	26.1	(24.5-27.8)	29.6	(27.9-31.4)	2.5	(2.0-3.0)	24.9	(23.3-26.6)	16.9	(15.5-18.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,499	70.5	(68.3-72.5)	24.0	(22.1-26.1)	1.4	(0.9-2.0)	2.7	(2.2-3.5)	1.4	(1.0-2.1)
Age (years)	18-24	130	57.2	(46.3-67.4)	39.1	(29.2-50.1)	0.2	(0.1-0.8)*	1.8	(0.6-5.3)*	1.7	(0.5-5.2)*
	25-34	405	65.2	(59.1-70.8)	30.3	(24.7-36.4)	0.8	(0.3-2.4)*	1.4	(0.7-3.1)*	2.3	(1.1-4.9)*
	35-54	1,492	75.2	(72.1-78.0)	19.0	(16.5-21.9)	1.7	(1.0-2.9)	2.9	(2.0-4.2)	1.2	(0.7-2.0)
	55+	2,445	72.1	(69.7-74.5)	21.0	(18.9-23.2)	1.7	(1.1-2.6)	3.6	(2.7-5.0)	1.5	(0.8-3.0)*
Sex	Men	1,700	63.5	(59.9-66.8)	30.7	(27.4-34.2)	1.3	(0.8-2.1)	2.9	(2.1-4.1)	1.6	(1.0-2.6)
	Women	2,799	77.0	(74.6-79.2)	17.8	(15.8-20.0)	1.4	(0.9-2.2)	2.6	(1.9-3.5)	1.3	(0.7-2.3)*
Race/ Ethnicity	Hispanic or Latino	85	66.2	(51.3-78.4)	25.1	(14.2-40.3)	3.7	(0.8-16.0)*	2.7	(0.9-8.2)*	2.3	(0.6-8.7)*
	Not Hispanic or Latino:	3,842	69.9	(67.7-72.1)	24.8	(22.7-27.0)	1.2	(0.9-1.7)	2.9	(2.4-3.6)	1.1	(0.8-1.6)
	White											
	Black or African American	368	62.3	(55.0-69.0)	19.6	(14.2-26.3)	1.6	(0.5-4.9)*	3.3	(1.5-7.2)*	13.3	(10.4-16.8)
	All Other Races	173	77.2	(68.1-84.2)	16.8	(10.9-25.0)	1.7	(0.7-3.7)*	3.2	(0.9-11.3)*	1.2	(0.3-5.0)*
Education	<High School	445	61.5	(53.2-69.1)	27.7	(21.0-35.7)	1.8	(0.8-4.1)*	6.2	(3.8-10.0)	2.7	(0.9-7.8)*
	High School Graduate (or GED)	1,541	66.4	(62.7-70.0)	26.7	(23.4-30.3)	1.6	(1.0-2.7)	3.1	(2.2-4.4)	2.2	(1.3-3.5)
	Some College	1,218	70.4	(66.4-74.2)	25.3	(21.7-29.4)	1.3	(0.7-2.3)*	2.2	(1.4-3.5)	0.8	(0.3-1.8)*
	College Graduate	1,288	76.4	(72.7-79.8)	20.1	(16.9-23.7)	1.1	(0.5-2.5)*	1.7	(1.0-3.0)	0.7	(0.3-1.8)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	807	69.5	(63.9-74.5)	24.1	(19.5-29.5)	1.1	(0.5-2.1)*	4.3	(2.6-6.8)	1.1	(0.6-2.0)*
	\$20,000-\$34,999	945	69.5	(64.8-73.8)	23.7	(19.7-28.2)	1.6	(0.8-3.0)*	3.5	(2.2-5.5)	1.7	(0.9-3.2)*
	\$35,000-\$49,999	687	71.7	(65.9-76.8)	23.5	(18.5-29.3)	1.3	(0.6-2.9)*	2.7	(1.5-4.7)	0.9	(0.3-2.7)*
	\$50,000-\$74,999	646	73.8	(68.6-78.4)	22.7	(18.2-27.9)	1.4	(0.6-3.1)*	1.1	(0.5-2.3)*	1.0	(0.3-3.1)*
	\$75,000 and more	841	73.7	(69.3-77.6)	22.7	(18.9-27.0)	0.7	(0.3-1.9)*	2.4	(1.4-4.3)	0.5	(0.1-1.6)*
Employment Status	Employed	2,210	69.9	(66.9-72.6)	25.4	(22.7-28.4)	1.2	(0.7-2.0)	2.4	(1.7-3.4)	1.2	(0.6-2.1)*
	Out of work	217	73.0	(64.0-80.4)	20.0	(13.6-28.6)	0.0	(0.0-0.0)	3.5	(1.7-6.9)*	3.5	(1.4-8.4)*
	Homemaker	317	74.6	(67.6-80.4)	19.0	(14.0-25.2)	1.8	(0.5-6.7)*	3.9	(2.0-7.6)*	0.7	(0.2-2.2)*
	Student	77	73.7	(60.1-83.9)	23.9	(14.3-37.2)	1.4	(0.2-7.5)*	1.0	(0.2-4.2)*	0.0	(0.0-0.0)
	Retired	1,291	67.9	(64.3-71.4)	25.1	(21.9-28.6)	2.7	(1.7-4.4)	2.9	(1.8-4.5)	1.4	(0.6-3.5)*
	Unable to work	375	73.2	(66.6-78.8)	19.9	(15.1-25.8)	0.3	(0.0-1.9)*	4.7	(2.4-9.1)*	1.9	(0.8-4.3)*
Veteran Status	Yes	636	64.8	(59.8-69.5)	28.9	(24.4-33.8)	2.3	(1.3-4.0)	3.1	(1.8-5.2)	1.0	(0.4-2.4)*
	No	3,858	71.3	(69.0-73.5)	23.3	(21.2-25.6)	1.2	(0.8-1.8)	2.6	(2.0-3.4)	1.5	(1.0-2.3)
Diabetes	Yes	540	76.6	(71.7-81.0)	17.9	(14.1-22.5)	1.0	(0.3-3.3)*	3.5	(2.0-6.1)	0.9	(0.4-1.8)*
	No	3,955	69.9	(67.7-72.1)	24.5	(22.5-26.7)	1.4	(0.9-2.0)	2.7	(2.1-3.4)	1.5	(1.0-2.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	585	70.0	(64.9-74.7)	22.7	(18.6-27.6)	2.8	(1.4-5.8)*	3.1	(1.7-5.7)*	1.3	(0.5-2.9)*
	No	3,878	70.5	(68.2-72.7)	24.1	(22.0-26.3)	1.2	(0.8-1.8)	2.7	(2.1-3.5)	1.5	(1.0-2.3)
Asthma	Yes	598	70.2	(63.6-76.1)	23.6	(18.1-30.1)	0.6	(0.3-1.4)*	4.1	(2.5-6.7)	1.5	(0.5-4.1)*
	No	3,887	70.6	(68.4-72.7)	24.0	(21.9-26.1)	1.5	(1.0-2.2)	2.5	(2.0-3.3)	1.4	(0.9-2.2)
Arthritis	Yes	1,841	70.9	(67.7-73.8)	23.1	(20.4-26.1)	1.3	(0.7-2.3)*	3.2	(2.2-4.8)	1.5	(0.8-3.1)*
	No	2,647	70.2	(67.5-72.8)	24.5	(22.0-27.1)	1.4	(0.9-2.1)	2.5	(1.9-3.4)	1.4	(0.9-2.2)
Frequent Mental Distress	Yes	566	72.2	(66.0-77.6)	21.8	(16.7-28.0)	0.5	(0.2-1.3)*	4.5	(2.8-7.0)	1.0	(0.5-2.0)*
	No	3,892	70.3	(68.0-72.4)	24.3	(22.2-26.5)	1.5	(1.0-2.2)	2.5	(1.9-3.2)	1.5	(1.0-2.3)
Serious Psychological Distress	Yes	204	58.7	(47.4-69.2)	34.8	(24.3-46.9)	0.4	(0.1-1.7)*	3.8	(1.9-7.1)*	2.4	(1.1-4.8)*
	No	4,251	71.2	(69.0-73.2)	23.5	(21.6-25.6)	1.3	(0.9-1.8)	2.7	(2.1-3.5)	1.3	(0.8-2.0)
Receiving Treatment for Mental Health Condition	Yes	658	83.2	(78.6-86.9)	13.9	(10.5-18.3)	0.1	(0.0-0.5)*	2.2	(1.1-4.0)*	0.6	(0.3-1.6)*
	No	3,837	68.5	(66.2-70.8)	25.5	(23.4-27.8)	1.5	(1.1-2.2)	2.9	(2.2-3.7)	1.6	(1.0-2.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,522	21.6	(19.8-23.5)	35.3	(33.1-37.5)	2.9	(2.1-4.0)	28.5	(26.5-30.7)	11.7	(10.4-13.2)
Age (years)	18-24	130	18.6	(11.5-28.5)	37.6	(27.5-49.0)	5.0	(1.3-17.4)*	30.4	(21.1-41.7)	8.4	(4.6-14.8)
	25-34	404	17.0	(12.6-22.6)	35.3	(29.7-41.4)	3.6	(1.7-7.1)*	33.0	(27.5-39.0)	11.1	(7.5-16.3)
	35-54	1,497	19.6	(17.0-22.6)	34.7	(31.4-38.1)	2.2	(1.3-3.8)	29.8	(26.8-33.1)	13.6	(11.4-16.2)
	55+	2,464	27.5	(25.2-29.9)	34.8	(32.2-37.4)	2.6	(1.9-3.6)	23.8	(21.6-26.1)	11.3	(9.7-13.2)
Sex	Men	1,707	24.3	(21.4-27.4)	37.6	(34.1-41.2)	3.4	(2.0-5.7)	24.9	(21.9-28.2)	9.8	(7.9-12.2)
	Women	2,815	19.1	(17.0-21.3)	33.1	(30.5-35.7)	2.4	(1.7-3.5)	32.0	(29.4-34.7)	13.5	(11.7-15.4)
Race/ Ethnicity	Hispanic or Latino	84	17.5	(8.1-34.0)*	30.5	(17.5-47.5)	33.4	(22.6-46.3)	14.3	(6.8-27.5)*	4.3	(1.8-10.0)*
	Not Hispanic or Latino:	3,865	20.6	(18.8-22.5)	35.8	(33.4-38.2)	2.8	(1.9-4.0)	29.5	(27.3-31.8)	11.4	(10.0-13.0)
	White											
	Black or African American	370	29.8	(23.7-36.8)	31.3	(24.4-39.1)	1.8	(0.7-4.6)*	24.8	(18.8-31.9)	12.3	(8.1-18.3)
	All Other Races	173	15.1	(9.5-23.3)	25.2	(16.6-36.4)	1.7	(0.6-4.6)*	38.6	(27.9-50.5)	19.4	(12.5-28.8)
Education	<High School	451	26.6	(20.9-33.1)	27.2	(20.7-34.7)	4.8	(1.3-16.0)*	27.1	(20.4-35.1)	14.4	(10.2-19.9)
	High School Graduate (or GED)	1,547	26.5	(23.4-30.0)	33.6	(30.0-37.4)	1.9	(1.3-2.9)	25.6	(22.3-29.2)	12.3	(10.0-15.1)
	Some College	1,234	18.8	(15.7-22.4)	36.4	(32.3-40.8)	2.3	(1.4-3.8)	29.8	(26.0-34.0)	12.6	(9.8-15.9)
	College Graduate	1,283	18.2	(15.0-21.8)	38.1	(34.2-42.2)	4.1	(2.4-6.9)	30.0	(26.5-33.7)	9.7	(7.7-12.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	811	23.3	(19.1-28.0)	30.1	(24.9-35.9)	4.2	(2.2-7.7)*	24.3	(19.3-30.0)	18.2	(14.4-22.8)
	\$20,000-\$34,999	960	19.3	(16.3-22.7)	35.4	(30.8-40.4)	1.8	(1.0-3.2)*	32.3	(27.6-37.4)	11.2	(8.6-14.4)
	\$35,000-\$49,999	688	22.5	(18.3-27.4)	34.7	(29.2-40.6)	2.8	(0.7-10.3)*	29.0	(23.9-34.6)	11.0	(7.4-15.9)
	\$50,000-\$74,999	648	26.3	(21.4-31.9)	34.7	(29.7-40.1)	2.7	(1.5-4.9)*	26.4	(22.1-31.2)	9.8	(7.1-13.5)
	\$75,000 and more	840	18.5	(14.7-23.1)	39.2	(34.7-43.9)	3.6	(1.9-6.4)*	29.6	(25.7-33.9)	9.1	(6.8-11.9)
Employment Status	Employed	2,214	21.7	(19.2-24.5)	35.1	(32.2-38.1)	3.0	(2.0-4.4)	29.2	(26.5-32.0)	11.1	(9.2-13.3)
	Out of work	220	17.4	(11.3-25.9)	38.5	(29.6-48.2)	2.4	(0.4-12.1)*	26.0	(18.5-35.2)	15.7	(10.1-23.5)
	Homemaker	317	18.5	(13.7-24.4)	43.1	(35.6-51.0)	2.0	(0.7-5.2)*	26.0	(19.4-33.9)	10.4	(7.0-15.2)
	Student	77	19.7	(9.6-36.4)*	27.6	(16.6-42.1)	7.9	(1.6-31.6)*	36.3	(23.0-52.0)	8.5	(3.7-18.5)*
	Retired	1,306	26.1	(23.0-29.5)	37.9	(34.0-41.9)	2.7	(1.8-4.0)	24.2	(21.1-27.6)	9.1	(7.1-11.6)
	Unable to work	376	17.5	(13.2-22.8)	25.5	(19.1-33.2)	1.3	(0.5-3.0)*	32.2	(25.2-40.1)	23.5	(17.6-30.7)
Veteran Status	Yes	637	22.5	(18.9-26.6)	34.4	(29.5-39.7)	1.5	(0.9-2.8)	31.4	(26.4-36.9)	10.1	(7.3-13.9)
	No	3,880	21.4	(19.4-23.5)	35.3	(33.0-37.8)	3.1	(2.2-4.5)	28.2	(26.0-30.5)	12.0	(10.5-13.6)
Diabetes	Yes	543	22.5	(18.4-27.2)	30.4	(25.2-36.1)	1.8	(0.8-4.0)*	28.3	(22.9-34.3)	17.0	(12.4-22.8)
	No	3,975	21.5	(19.6-23.5)	35.7	(33.4-38.0)	3.0	(2.1-4.2)	28.6	(26.5-30.8)	11.3	(9.9-12.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Missouri, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	593	23.9	(19.9-28.3)	33.4	(28.3-39.0)	2.3	(1.3-3.8)	24.9	(20.3-30.1)	15.5	(11.7-20.3)
	No	3,891	21.4	(19.5-23.5)	35.5	(33.1-37.8)	3.0	(2.1-4.3)	28.7	(26.6-31.0)	11.4	(9.9-13.0)
Asthma	Yes	599	20.1	(15.5-25.7)	28.1	(22.2-34.9)	3.8	(1.2-11.3)*	33.5	(27.4-40.2)	14.4	(10.7-19.0)
	No	3,909	21.6	(19.7-23.7)	36.6	(34.3-39.0)	2.7	(1.9-3.7)	27.8	(25.7-30.0)	11.3	(9.9-12.9)
Arthritis	Yes	1,842	21.0	(18.6-23.6)	33.6	(30.4-37.0)	2.7	(1.7-4.2)	28.1	(25.0-31.4)	14.6	(12.3-17.3)
	No	2,669	21.6	(19.3-24.2)	35.9	(33.2-38.8)	3.0	(1.9-4.6)	28.9	(26.4-31.6)	10.5	(8.8-12.4)
Frequent Mental Distress	Yes	568	15.3	(12.0-19.3)	26.3	(21.1-32.2)	1.6	(0.8-3.1)*	37.3	(31.2-43.8)	19.6	(14.4-26.0)
	No	3,912	22.4	(20.4-24.5)	36.5	(34.1-38.9)	3.1	(2.2-4.4)	27.5	(25.3-29.7)	10.6	(9.3-12.1)
Serious Psychological Distress	Yes	204	15.3	(10.0-22.6)	23.3	(15.1-34.2)	1.6	(0.5-5.0)*	32.8	(22.6-44.9)	27.0	(16.4-41.1)
	No	4,271	22.0	(20.1-24.0)	35.8	(33.5-38.1)	2.9	(2.1-4.2)	28.4	(26.3-30.6)	10.9	(9.6-12.3)
Receiving Treatment for Mental Health Condition	Yes	654	15.4	(11.9-19.6)	28.3	(23.0-34.2)	3.9	(1.2-12.0)*	33.3	(28.1-39.0)	19.2	(15.4-23.6)
	No	3,863	22.9	(20.9-25.0)	35.9	(33.5-38.3)	2.7	(2.0-3.7)	27.8	(25.6-30.1)	10.7	(9.3-12.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,233	64.3	(62.5-66.1)	26.4	(24.8-28.2)	3.4	(2.7-4.2)	3.8	(3.1-4.7)	2.0	(1.5-2.8)
Age (years)	18-24	196	54.1	(45.8-62.2)	33.3	(26.1-41.4)	4.5	(2.1-9.5)*	5.9	(3.3-10.3)	2.1	(0.7-6.1)*
	25-34	499	56.5	(51.0-61.9)	33.8	(28.8-39.1)	2.5	(1.3-4.8)*	5.0	(3.0-8.2)	2.2	(0.8-5.7)*
	35-54	1,923	66.9	(64.3-69.4)	24.6	(22.3-27.1)	3.5	(2.6-4.6)	2.9	(1.9-4.4)	2.1	(1.3-3.3)
	55+	2,590	69.2	(67.2-71.3)	21.9	(20.1-23.8)	3.6	(2.8-4.6)	3.4	(2.7-4.3)	1.9	(1.3-2.8)
Sex	Men	2,060	56.5	(53.6-59.4)	32.0	(29.2-34.8)	4.2	(3.2-5.5)	4.4	(3.3-5.9)	2.9	(1.9-4.4)
	Women	3,173	71.7	(69.5-73.7)	21.2	(19.3-23.2)	2.7	(1.9-3.7)	3.2	(2.4-4.4)	1.2	(0.8-1.9)
Race/ Ethnicity	Hispanic or Latino	79	63.5	(45.6-78.3)	29.7	(17.0-46.5)	3.1	(0.8-10.9)*	2.9	(0.8-9.5)*	0.9	(0.1-6.0)*
	Not Hispanic or Latino:	4,674	65.4	(63.5-67.2)	26.5	(24.8-28.3)	3.2	(2.6-3.9)	3.4	(2.7-4.3)	1.5	(1.0-2.2)
	White											
	Black or African American	12	--	--	--	--	--	--	--	--	--	--
	All Other Races	426	57.8	(50.8-64.6)	25.9	(20.3-32.4)	6.4	(3.0-13.1)*	5.8	(3.5-9.3)	4.1	(2.4-7.0)
Education	<High School	343	53.1	(45.2-60.8)	28.1	(21.6-35.8)	5.8	(2.9-11.2)*	6.8	(4.1-10.9)	6.2	(3.2-11.7)*
	High School Graduate (or GED)	1,609	60.3	(56.9-63.7)	29.6	(26.6-32.9)	3.3	(2.3-4.6)	4.5	(3.1-6.5)	2.3	(1.4-3.5)
	Some College	1,529	62.4	(58.9-65.7)	28.1	(25.0-31.4)	4.4	(3.1-6.2)	3.8	(2.5-5.6)	1.3	(0.7-2.5)*
	College Graduate	1,744	71.8	(68.8-74.6)	21.9	(19.3-24.7)	2.3	(1.6-3.4)	2.4	(1.6-3.7)	1.6	(0.7-3.4)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	887	58.2	(53.6-62.7)	28.7	(24.4-33.3)	4.1	(2.6-6.4)	5.5	(3.8-8.0)	3.5	(1.7-7.1)*
	\$20,000-\$34,999	1,176	64.4	(60.5-68.1)	28.0	(24.5-31.7)	2.7	(1.9-3.8)	3.1	(2.1-4.6)	1.9	(1.1-3.3)
	\$35,000-\$49,999	888	61.5	(57.1-65.7)	26.5	(22.7-30.6)	4.1	(2.5-6.6)	5.7	(3.5-9.3)	2.3	(0.9-5.4)*
	\$50,000-\$74,999	851	69.2	(64.9-73.1)	25.2	(21.6-29.2)	1.7	(0.8-3.6)*	2.7	(1.4-5.2)*	1.2	(0.4-3.4)*
	\$75,000 and more	887	69.9	(65.5-74.0)	23.4	(19.6-27.6)	2.7	(1.7-4.3)	3.1	(1.6-6.0)*	0.9	(0.4-2.3)*
Employment Status	Employed	3,002	65.3	(62.9-67.6)	25.8	(23.7-28.0)	3.6	(2.7-4.7)	3.7	(2.8-5.0)	1.7	(1.0-2.7)
	Out of work	169	67.1	(57.1-75.7)	22.0	(14.9-31.3)	2.9	(0.4-16.7)*	4.1	(1.9-8.5)*	4.0	(1.1-13.8)*
	Homemaker	400	62.0	(55.3-68.2)	28.7	(22.9-35.3)	1.4	(0.6-3.4)*	6.2	(3.2-11.8)*	1.7	(0.6-4.9)*
	Student	80	67.1	(53.0-78.6)	27.9	(17.5-41.2)	3.7	(0.8-15.0)*	1.4	(0.2-8.3)*	0.0	(0.0-0.0)
	Retired	1,285	60.5	(57.3-63.6)	28.6	(25.7-31.6)	4.3	(3.1-5.8)	3.8	(2.7-5.3)	2.9	(1.7-4.8)
	Unable to work	279	63.2	(56.3-69.7)	27.1	(20.9-34.3)	2.1	(0.9-5.2)*	4.1	(2.3-7.1)	3.4	(1.5-7.5)*
Veteran Status	Yes	806	59.6	(55.5-63.7)	28.8	(25.0-33.0)	4.1	(2.7-6.2)	4.7	(3.0-7.3)	2.7	(1.5-4.7)
	No	4,425	65.1	(63.1-67.1)	26.0	(24.2-27.9)	3.3	(2.5-4.1)	3.7	(2.9-4.7)	1.9	(1.3-2.8)
Diabetes	Yes	467	61.3	(55.7-66.5)	30.2	(25.2-35.8)	4.2	(2.4-7.1)	2.7	(1.5-4.6)	1.6	(0.6-4.6)*
	No	4,764	64.5	(62.6-66.4)	26.1	(24.4-27.9)	3.4	(2.7-4.2)	3.9	(3.1-4.9)	2.1	(1.5-2.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	505	60.9 (55.8-65.8)	27.4 (22.9-32.4)	4.3 (2.7-6.9)	3.6 (2.1-6.0)	3.7 (2.0-6.9)*				
	No	4,690	64.7 (62.8-66.6)	26.4 (24.6-28.2)	3.2 (2.5-4.0)	3.9 (3.1-4.8)	1.9 (1.3-2.7)				
Asthma	Yes	680	62.1 (57.1-66.9)	24.3 (20.0-29.2)	3.5 (2.0-6.1)	5.8 (3.7-9.1)	4.3 (2.0-8.9)*				
	No	4,548	64.6 (62.6-66.5)	26.7 (25.0-28.6)	3.4 (2.7-4.2)	3.5 (2.8-4.5)	1.8 (1.3-2.4)				
Arthritis	Yes	1,895	68.8 (65.9-71.5)	24.1 (21.6-26.8)	2.7 (2.0-3.7)	2.7 (2.0-3.7)	1.7 (1.0-3.0)				
	No	3,324	62.6 (60.3-64.8)	27.4 (25.3-29.5)	3.7 (2.8-4.8)	4.2 (3.2-5.4)	2.2 (1.5-3.2)				
Frequent Mental Distress	Yes	503	64.3 (58.0-70.2)	25.6 (20.6-31.3)	3.3 (1.3-8.0)*	3.9 (1.9-7.6)*	2.9 (1.6-5.4)*				
	No	4,666	64.3 (62.4-66.2)	26.5 (24.8-28.4)	3.4 (2.7-4.2)	3.8 (3.0-4.8)	1.9 (1.3-2.8)				
Serious Psychological Distress	Yes	184	47.7 (37.2-58.3)	32.7 (24.0-42.8)	4.0 (0.7-18.9)*	3.5 (1.6-7.8)*	12.1 (7.8-18.3)				
	No	4,970	64.8 (62.9-66.6)	26.3 (24.6-28.1)	3.3 (2.6-4.0)	3.9 (3.1-4.9)	1.8 (1.2-2.6)				
Receiving Treatment for Mental Health Condition	Yes	718	69.2 (64.3-73.8)	20.9 (17.0-25.4)	2.7 (1.2-6.1)*	4.1 (2.3-7.3)	3.0 (1.2-7.2)*				
	No	4,501	63.8 (61.8-65.7)	27.1 (25.3-29.0)	3.5 (2.8-4.3)	3.8 (3.0-4.7)	1.9 (1.3-2.6)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,247	19.8	(18.3-21.4)	39.8	(37.9-41.7)	4.8	(4.1-5.6)	26.2	(24.6-27.8)	9.4	(8.5-10.4)
Age (years)	18-24	198	21.7	(15.6-29.3)	45.1	(36.6-53.9)	2.8	(1.2-6.2)*	27.4	(20.5-35.6)	3.0	(1.2-7.2)*
	25-34	503	14.6	(10.8-19.4)	43.1	(37.6-48.7)	4.2	(2.6-6.8)	28.5	(23.9-33.5)	9.7	(6.9-13.4)
	35-54	1,926	16.8	(14.8-19.0)	39.7	(37.1-42.4)	5.7	(4.5-7.0)	26.9	(24.6-29.3)	10.9	(9.4-12.6)
	55+	2,596	25.2	(23.2-27.2)	36.6	(34.5-38.8)	5.0	(4.2-6.1)	23.4	(21.6-25.3)	9.8	(8.6-11.2)
Sex	Men	2,076	23.0	(20.5-25.6)	42.5	(39.6-45.4)	4.9	(3.9-6.1)	22.8	(20.4-25.3)	6.9	(5.6-8.5)
	Women	3,171	16.9	(15.2-18.7)	37.2	(34.9-39.5)	4.7	(3.9-5.7)	29.5	(27.4-31.6)	11.7	(10.5-13.1)
Race/ Ethnicity	Hispanic or Latino	79	28.2	(14.7-47.3)*	35.9	(22.3-52.2)	2.9	(0.7-11.3)*	23.9	(13.7-38.3)	9.0	(4.7-16.5)*
	Not Hispanic or Latino:	4,676	17.7	(16.2-19.2)	41.0	(39.0-42.9)	4.9	(4.2-5.7)	27.2	(25.5-29.0)	9.2	(8.3-10.3)
	White											
	Black or African American	13	--	--	--	--	--	--	--	--	--	--
	All Other Races	438	19.7	(14.9-25.7)	35.5	(28.5-43.1)	5.3	(3.4-8.1)	28.7	(23.1-35.1)	10.8	(7.5-15.5)
Education	<High School	340	24.3	(17.9-32.0)	33.3	(26.3-41.1)	3.6	(1.5-8.5)*	27.5	(20.8-35.3)	11.4	(7.1-17.8)
	High School Graduate (or GED)	1,621	24.9	(21.9-28.2)	37.3	(33.9-40.9)	5.0	(3.8-6.6)	23.4	(20.7-26.3)	9.4	(7.7-11.4)
	Some College	1,531	18.0	(15.5-20.7)	40.4	(37.0-43.9)	5.3	(4.1-6.7)	27.3	(24.2-30.6)	9.0	(7.5-10.8)
	College Graduate	1,748	14.2	(12.1-16.6)	43.1	(40.1-46.2)	4.5	(3.5-5.8)	28.4	(25.8-31.1)	9.8	(8.3-11.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	880	19.8	(16.3-23.7)	36.9	(32.3-41.8)	4.9	(3.4-7.1)	26.0	(22.1-30.2)	12.4	(9.9-15.4)
	\$20,000-\$34,999	1,187	20.0	(17.1-23.3)	38.4	(34.6-42.4)	4.3	(3.1-6.0)	26.4	(23.1-30.1)	10.8	(8.8-13.2)
	\$35,000-\$49,999	884	17.9	(14.7-21.6)	40.8	(36.5-45.2)	5.3	(3.9-7.3)	26.9	(23.2-31.0)	9.1	(7.2-11.4)
	\$50,000-\$74,999	846	16.6	(13.7-20.0)	43.5	(39.2-47.9)	4.6	(3.0-7.0)	26.8	(23.4-30.6)	8.3	(6.6-10.6)
	\$75,000 and more	891	22.1	(18.1-26.7)	39.5	(35.2-43.9)	4.6	(3.3-6.4)	25.0	(21.5-28.9)	8.8	(6.7-11.5)
Employment Status	Employed	3,019	19.1	(17.1-21.2)	40.4	(38.0-42.9)	4.9	(4.0-5.9)	26.4	(24.4-28.6)	9.2	(8.0-10.5)
	Out of work	169	13.2	(7.8-21.4)	43.9	(33.0-55.4)	7.4	(4.0-13.5)*	23.1	(15.9-32.4)	12.4	(6.4-22.8)*
	Homemaker	393	19.3	(14.8-24.7)	42.4	(36.2-48.9)	4.9	(2.7-8.6)	29.0	(23.5-35.2)	4.4	(2.9-6.6)
	Student	80	32.4	(20.0-47.9)	42.5	(28.7-57.6)	1.0	(0.3-3.6)*	18.7	(9.8-32.6)*	5.5	(2.0-14.3)*
	Retired	1,294	23.3	(20.8-26.0)	38.1	(35.0-41.3)	5.1	(3.9-6.6)	25.4	(22.7-28.3)	8.1	(6.5-10.1)
	Unable to work	274	15.0	(11.0-20.3)	24.9	(19.0-32.0)	4.5	(2.2-9.0)*	29.1	(22.9-36.2)	26.4	(20.9-32.8)
Veteran Status	Yes	810	21.5	(18.0-25.4)	38.4	(34.2-42.8)	6.9	(5.2-9.2)	25.7	(21.8-30.0)	7.5	(5.6-10.0)
	No	4,435	19.5	(17.8-21.3)	40.1	(38.0-42.2)	4.5	(3.8-5.3)	26.3	(24.6-28.1)	9.7	(8.7-10.9)
Diabetes	Yes	463	20.3	(16.2-25.1)	37.6	(32.3-43.3)	3.7	(2.3-5.9)	25.2	(20.4-30.7)	13.2	(10.0-17.2)
	No	4,781	19.8	(18.2-21.5)	39.9	(38.0-41.9)	4.9	(4.2-5.7)	26.2	(24.6-28.0)	9.1	(8.2-10.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Montana, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	505	22.9	(19.0-27.2)	33.7	(28.8-38.9)	4.7	(2.9-7.5)	22.6	(18.5-27.2)	16.2	(12.6-20.6)
	No	4,704	19.7	(18.0-21.4)	40.3	(38.3-42.3)	4.8	(4.1-5.6)	26.5	(24.8-28.2)	8.8	(7.9-9.8)
Asthma	Yes	684	20.2	(15.9-25.3)	35.6	(30.8-40.8)	4.8	(3.2-7.1)	26.1	(22.0-30.7)	13.2	(10.3-16.8)
	No	4,557	19.7	(18.2-21.4)	40.4	(38.4-42.5)	4.8	(4.1-5.6)	26.2	(24.5-28.0)	8.9	(7.9-9.9)
Arthritis	Yes	1,903	19.2	(17.2-21.4)	38.1	(35.2-41.0)	4.0	(3.1-5.1)	26.6	(24.1-29.2)	12.1	(10.4-14.0)
	No	3,329	19.9	(18.0-22.1)	40.4	(38.0-42.7)	5.1	(4.3-6.2)	26.2	(24.2-28.2)	8.4	(7.3-9.7)
Frequent Mental Distress	Yes	507	14.4	(10.7-19.1)	33.7	(27.7-40.2)	4.2	(2.1-8.3)*	25.8	(21.2-31.0)	21.9	(17.4-27.2)
	No	4,675	20.3	(18.7-22.1)	40.5	(38.5-42.5)	4.8	(4.1-5.6)	26.2	(24.5-28.0)	8.2	(7.3-9.2)
Serious Psychological Distress	Yes	186	16.4	(10.5-24.8)	22.1	(15.1-31.1)	3.2	(1.2-8.2)*	29.2	(21.2-38.7)	29.1	(20.7-39.1)
	No	4,984	19.8	(18.2-21.4)	40.6	(38.7-42.6)	4.8	(4.1-5.6)	26.1	(24.4-27.7)	8.7	(7.8-9.7)
Receiving Treatment for Mental Health Condition	Yes	716	15.7	(12.3-20.0)	35.3	(30.4-40.6)	5.1	(3.2-7.9)	27.8	(23.8-32.1)	16.1	(13.0-19.8)
	No	4,517	20.4	(18.7-22.1)	40.3	(38.3-42.4)	4.8	(4.1-5.6)	26.1	(24.3-27.8)	8.5	(7.5-9.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,730	61.4	(58.7-64.0)	29.5	(26.9-32.3)	1.8	(1.1-2.8)	5.7	(4.3-7.6)	1.6	(1.1-2.5)
Age (years)	18-24	109	45.1	(30.8-60.3)	34.7	(22.1-49.9)	4.6	(0.9-20.0)*	15.3	(5.9-34.4)*	0.2	(0.0-1.5)*
	25-34	396	51.6	(44.2-58.8)	37.3	(30.0-45.3)	1.0	(0.3-3.1)*	8.8	(4.8-15.7)*	1.3	(0.4-3.9)*
	35-54	1,531	65.5	(61.1-69.6)	27.1	(23.0-31.7)	1.5	(0.7-3.0)*	3.6	(2.5-5.2)	2.3	(1.1-4.7)*
	55+	2,684	66.2	(63.4-68.9)	25.2	(22.8-27.8)	1.8	(1.2-2.8)	4.8	(3.7-6.2)	2.0	(1.3-3.0)
Sex	Men	1,831	54.7	(50.4-58.9)	32.7	(28.5-37.1)	2.1	(1.2-3.5)	8.2	(5.5-12.0)	2.4	(1.4-4.1)
	Women	2,899	67.6	(64.4-70.7)	26.5	(23.4-29.8)	1.5	(0.7-3.0)*	3.4	(2.5-4.7)	1.0	(0.5-1.9)*
Race/ Ethnicity	Hispanic or Latino	149	53.7	(38.6-68.1)	43.0	(28.7-58.7)	0.7	(0.2-1.7)*	2.2	(0.7-7.1)*	0.4	(0.2-0.8)*
	Not Hispanic or Latino:	4,419	62.7	(60.2-65.3)	28.2	(25.8-30.7)	1.7	(1.1-2.6)	5.6	(4.1-7.5)	1.8	(1.2-2.6)
	White											
	Black or African American	36	--	--	--	--	--	--	--	--	--	--
	All Other Races	113	48.3	(27.1-70.1)	33.1	(12.9-62.2)*	0.7	(0.2-2.2)*	13.7	(7.7-23.2)	4.2	(2.0-8.7)*
Education	<High School	331	58.2	(46.1-69.5)	33.4	(22.1-47.1)	0.8	(0.2-2.5)*	5.7	(2.9-10.9)*	1.9	(0.7-4.6)*
	High School Graduate (or GED)	1,701	57.3	(52.8-61.7)	33.3	(28.9-38.0)	1.1	(0.7-1.8)	6.6	(4.6-9.3)	1.7	(1.0-2.8)
	Some College	1,404	58.4	(53.1-63.5)	30.7	(25.4-36.6)	3.5	(1.7-7.1)*	6.8	(3.5-12.6)*	0.6	(0.3-1.1)*
	College Graduate	1,290	68.0	(63.3-72.3)	24.4	(20.6-28.6)	1.2	(0.5-3.0)*	3.8	(2.1-6.9)*	2.6	(1.1-5.8)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	778	58.4	(51.5-65.0)	28.5	(22.1-36.0)	2.7	(1.2-5.7)*	8.5	(5.4-13.2)	1.9	(1.0-3.6)*
	\$20,000-\$34,999	1,072	57.1	(51.2-62.7)	31.7	(26.1-37.9)	0.8	(0.3-2.2)*	8.6	(4.3-16.6)*	1.8	(0.9-3.8)*
	\$35,000-\$49,999	772	62.4	(56.3-68.2)	32.6	(26.9-38.9)	1.4	(0.5-3.9)*	3.1	(1.8-5.5)	0.5	(0.1-2.1)*
	\$50,000-\$74,999	723	63.1	(55.8-69.8)	31.1	(24.2-39.0)	2.2	(0.6-7.6)*	2.6	(1.3-4.9)*	1.0	(0.3-3.1)*
	\$75,000 and more	911	69.6	(64.3-74.4)	22.4	(18.3-27.2)	1.5	(0.7-3.3)*	4.3	(2.4-7.8)*	2.1	(0.7-6.2)*
Employment Status	Employed	2,688	62.6	(59.2-65.9)	28.6	(25.4-32.1)	1.4	(0.9-2.4)	5.5	(3.6-8.4)	1.8	(1.0-3.3)*
	Out of work	130	61.9	(45.9-75.7)	26.7	(14.0-45.0)*	1.3	(0.2-8.8)*	8.7	(3.8-18.9)*	1.3	(0.3-5.0)*
	Homemaker	367	52.6	(42.4-62.6)	41.5	(30.9-52.9)	2.0	(0.5-7.4)*	3.1	(1.3-7.5)*	0.8	(0.3-2.2)*
	Student	55	57.6	(38.3-74.9)	29.6	(15.3-49.3)*	7.8	(1.3-34.4)*	5.0	(1.5-15.5)*	0.0	(0.0-0.0)
	Retired	1,235	61.4	(57.2-65.5)	28.7	(25.0-32.8)	1.6	(1.0-2.7)	6.6	(4.5-9.5)	1.7	(1.0-2.9)
	Unable to work	251	58.9	(49.8-67.4)	31.8	(23.9-41.0)	1.3	(0.4-4.9)*	5.8	(2.8-11.5)*	2.2	(0.7-6.4)*
Veteran Status	Yes	639	60.4	(53.2-67.2)	30.6	(24.0-38.1)	1.4	(0.7-2.8)*	4.6	(2.3-9.0)*	2.9	(1.0-8.5)*
	No	4,086	61.5	(58.6-64.3)	29.4	(26.6-32.4)	1.8	(1.1-3.0)	5.9	(4.3-8.1)	1.4	(0.9-2.1)
Diabetes	Yes	521	63.1	(56.4-69.3)	29.5	(23.5-36.3)	1.3	(0.6-2.9)*	4.9	(2.7-8.7)	1.2	(0.6-2.4)*
	No	4,205	61.3	(58.4-64.0)	29.5	(26.7-32.5)	1.8	(1.1-2.9)	5.7	(4.2-7.8)	1.7	(1.1-2.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	564	62.4	(57.0-67.5)	26.0	(21.4-31.2)	1.4	(0.7-2.7)*	8.5	(5.4-13.1)	1.7	(0.9-3.3)*
	No	4,112	61.4	(58.5-64.2)	29.7	(26.9-32.7)	1.8	(1.1-2.9)	5.4	(3.9-7.5)	1.6	(1.0-2.6)
Asthma	Yes	455	71.5	(64.5-77.6)	20.3	(15.3-26.4)	1.6	(0.5-4.9)*	5.6	(3.1-9.9)	1.1	(0.4-2.7)*
	No	4,259	60.4	(57.5-63.2)	30.5	(27.6-33.4)	1.8	(1.1-2.8)	5.7	(4.1-7.8)	1.7	(1.1-2.6)
Arthritis	Yes	1,773	63.7	(59.4-67.7)	28.4	(24.3-32.9)	2.0	(1.0-4.0)*	4.6	(3.3-6.4)	1.4	(0.8-2.5)
	No	2,938	60.5	(57.2-63.7)	30.0	(26.7-33.4)	1.7	(1.0-2.9)	6.1	(4.2-8.7)	1.8	(1.0-3.0)
Frequent Mental Distress	Yes	356	53.5	(44.8-62.0)	36.0	(27.9-44.9)	0.8	(0.2-3.1)*	5.3	(3.0-8.9)	4.5	(1.1-16.6)*
	No	4,321	62.1	(59.2-64.8)	29.1	(26.3-32.0)	1.8	(1.2-2.9)	5.7	(4.1-7.7)	1.3	(0.9-1.9)
Serious Psychological Distress	Yes	128	54.3	(37.0-70.7)	35.8	(19.2-56.7)	0.2	(0.0-1.2)*	8.6	(3.9-17.9)*	1.1	(0.3-4.4)*
	No	4,560	62.0	(59.3-64.7)	29.2	(26.6-32.0)	1.7	(1.1-2.6)	5.4	(4.0-7.3)	1.7	(1.1-2.6)
Receiving Treatment for Mental Health Condition	Yes	519	74.0	(67.5-79.6)	21.0	(15.7-27.4)	0.5	(0.1-2.9)*	4.2	(2.2-8.1)*	0.3	(0.1-0.9)*
	No	4,203	60.3	(57.4-63.1)	30.2	(27.4-33.2)	1.9	(1.2-2.9)	5.9	(4.3-7.9)	1.8	(1.1-2.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,745	25.1	(22.7-27.6)	41.4	(38.7-44.2)	2.2	(1.5-3.2)	23.7	(21.6-26.0)	7.6	(6.3-9.0)
Age (years)	18-24	109	25.2	(13.0-43.2)*	48.7	(33.9-63.8)	1.6	(0.5-5.0)*	16.3	(8.8-28.2)	8.2	(2.7-22.1)*
	25-34	400	27.0	(20.4-34.9)	37.9	(30.8-45.5)	2.5	(0.9-6.6)*	26.1	(19.8-33.6)	6.5	(3.3-12.2)*
	35-54	1,529	19.8	(16.6-23.5)	44.6	(40.2-49.1)	2.0	(1.0-3.9)*	25.1	(21.8-28.7)	8.4	(6.7-10.6)
	55+	2,697	31.5	(28.8-34.3)	36.9	(34.0-39.8)	2.5	(1.7-3.6)	21.6	(19.4-24.0)	7.5	(6.1-9.3)
Sex	Men	1,845	26.7	(22.9-30.9)	44.5	(40.2-49.0)	2.9	(1.7-4.9)	19.6	(16.6-23.0)	6.2	(4.5-8.6)
	Women	2,900	23.5	(20.8-26.5)	38.4	(35.2-41.8)	1.6	(1.0-2.5)	27.7	(24.8-30.8)	8.8	(7.2-10.7)
Race/ Ethnicity	Hispanic or Latino	151	38.9	(25.5-54.2)	31.6	(17.8-49.6)	2.2	(0.5-9.9)*	22.2	(10.5-40.8)*	5.1	(2.7-9.4)*
	Not Hispanic or Latino:	4,429	23.7	(21.4-26.2)	41.5	(38.9-44.3)	2.0	(1.4-2.9)	24.8	(22.5-27.1)	8.0	(6.7-9.5)
	White											
	Black or African American	36	--	--	--	--	--	--	--	--	--	--
	All Other Races	112	22.0	(11.3-38.4)*	43.0	(22.3-66.5)	2.0	(0.5-7.6)*	22.3	(10.6-41.0)*	10.8	(4.6-23.3)*
Education	<High School	344	39.7	(29.6-50.8)	39.9	(28.7-52.2)	0.9	(0.3-2.1)*	12.4	(8.2-18.4)	7.1	(4.0-12.4)
	High School Graduate (or GED)	1,689	28.0	(24.0-32.4)	40.4	(36.0-45.0)	1.9	(1.1-3.0)	21.6	(17.9-25.7)	8.1	(6.1-10.7)
	Some College	1,413	27.0	(22.3-32.2)	40.3	(35.0-46.0)	2.2	(1.2-4.0)	23.0	(19.2-27.2)	7.5	(5.4-10.3)
	College Graduate	1,295	17.2	(13.7-21.4)	43.6	(39.0-48.3)	3.0	(1.5-6.1)*	28.5	(24.6-32.9)	7.6	(5.4-10.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	779	30.6	(23.5-38.6)	34.8	(28.3-42.0)	1.9	(0.9-3.9)*	21.1	(15.9-27.3)	11.7	(8.7-15.6)
	\$20,000-\$34,999	1,075	26.1	(20.5-32.6)	41.0	(34.9-47.4)	2.3	(0.8-6.4)*	22.8	(18.8-27.4)	7.9	(5.3-11.5)
	\$35,000-\$49,999	778	24.6	(19.9-29.9)	42.0	(36.1-48.0)	3.8	(2.0-6.8)*	19.9	(15.6-25.0)	9.8	(6.2-15.2)
	\$50,000-\$74,999	732	19.7	(14.7-25.9)	47.3	(40.2-54.5)	0.8	(0.4-1.7)*	25.6	(20.8-30.9)	6.7	(4.6-9.6)
	\$75,000 and more	909	22.8	(18.5-27.9)	42.5	(37.3-47.8)	2.5	(1.2-5.3)*	26.4	(22.0-31.3)	5.7	(3.6-9.0)
Employment Status	Employed	2,680	25.3	(22.2-28.8)	41.2	(37.8-44.7)	2.6	(1.6-4.1)	24.6	(21.9-27.5)	6.3	(5.0-8.0)
	Out of work	129	15.2	(9.0-24.5)	48.1	(31.8-64.8)	1.0	(0.1-6.7)*	18.2	(10.3-30.2)	17.5	(7.5-36.0)*
	Homemaker	369	30.9	(21.5-42.3)	38.5	(28.6-49.5)	1.0	(0.2-5.8)*	24.4	(17.2-33.3)	5.2	(3.3-8.0)
	Student	55	14.5	(6.2-30.2)*	50.6	(30.7-70.4)	0.0	(0.0-0.0)	23.4	(11.4-42.1)*	11.5	(2.5-39.8)*
	Retired	1,258	29.2	(25.4-33.3)	43.6	(39.1-48.2)	1.7	(1.0-2.8)	18.8	(15.8-22.3)	6.7	(4.9-9.0)
	Unable to work	250	18.7	(13.1-26.0)	26.3	(18.7-35.5)	3.6	(1.3-9.6)*	27.4	(20.1-36.2)	24.0	(16.7-33.3)
Veteran Status	Yes	644	25.5	(20.4-31.4)	47.1	(39.8-54.4)	1.3	(0.6-2.8)*	17.8	(13.5-23.1)	8.3	(5.0-13.5)
	No	4,098	25.0	(22.4-27.8)	40.6	(37.7-43.6)	2.4	(1.6-3.5)	24.5	(22.2-27.1)	7.5	(6.2-9.1)
Diabetes	Yes	520	24.6	(19.3-30.8)	40.5	(33.8-47.5)	1.5	(0.6-3.8)*	23.9	(18.7-29.9)	9.5	(6.4-14.0)
	No	4,221	25.1	(22.6-27.8)	41.5	(38.6-44.4)	2.3	(1.5-3.3)	23.8	(21.5-26.2)	7.4	(6.1-8.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nebraska, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	569	23.9	(19.7-28.7)	33.9	(28.4-39.8)	1.6	(0.8-3.1)*	29.8	(24.2-36.0)	10.9	(7.7-15.2)
	No	4,119	25.2	(22.6-27.9)	41.9	(39.0-44.8)	2.3	(1.5-3.4)	23.4	(21.1-25.8)	7.3	(6.0-8.9)
Asthma	Yes	458	21.9	(15.9-29.4)	39.4	(31.7-47.7)	1.3	(0.5-3.2)*	23.7	(17.6-31.0)	13.7	(8.4-21.5)
	No	4,271	25.4	(22.9-28.1)	41.7	(38.7-44.6)	2.3	(1.6-3.4)	23.7	(21.4-26.1)	6.9	(5.7-8.3)
Arthritis	Yes	1,783	21.8	(18.7-25.4)	40.5	(36.0-45.2)	3.0	(1.7-5.5)*	24.4	(21.3-27.9)	10.2	(8.1-12.7)
	No	2,941	26.4	(23.4-29.7)	41.7	(38.3-45.1)	1.8	(1.2-2.9)	23.4	(20.8-26.3)	6.6	(5.2-8.5)
Frequent Mental Distress	Yes	353	20.8	(14.6-28.6)	32.3	(24.1-41.8)	0.6	(0.3-1.5)*	31.5	(23.5-40.9)	14.8	(10.4-20.6)
	No	4,339	25.4	(22.9-28.1)	42.2	(39.3-45.1)	2.3	(1.6-3.4)	23.0	(20.8-25.4)	7.0	(5.7-8.5)
Serious Psychological Distress	Yes	129	20.9	(10.4-37.5)*	31.2	(14.1-55.6)*	1.5	(0.5-4.0)*	28.7	(16.9-44.4)	17.8	(10.2-29.2)
	No	4,572	25.4	(23.0-28.0)	41.8	(39.0-44.6)	2.3	(1.5-3.3)	23.2	(21.1-25.5)	7.3	(6.1-8.8)
Receiving Treatment for Mental Health Condition	Yes	518	19.4	(13.2-27.6)	33.2	(26.4-40.9)	2.0	(0.7-5.2)*	26.2	(20.8-32.5)	19.2	(14.6-24.7)
	No	4,218	25.8	(23.3-28.5)	41.6	(38.7-44.6)	2.2	(1.5-3.3)	23.4	(21.1-25.9)	6.9	(5.7-8.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,421	64.8	(61.8-67.6)	27.2	(24.6-30.0)	1.8	(1.2-2.5)	4.3	(3.1-6.0)	1.9	(1.1-3.4)
Age (years)												
	18-24	86	43.4	(30.5-57.3)	38.1	(25.7-52.3)	2.1	(0.7-6.6)*	9.6	(3.1-25.7)*	6.8	(2.2-19.1)*
	25-34	292	63.9	(54.6-72.2)	31.3	(23.8-40.0)	0.3	(0.1-0.7)*	3.5	(1.5-8.0)*	1.1	(0.4-3.0)*
	35-54	1,141	67.7	(63.6-71.5)	25.4	(21.7-29.4)	2.2	(1.2-3.9)	3.7	(2.3-6.0)	1.1	(0.5-2.4)*
	55+	1,860	66.8	(63.5-69.9)	24.5	(21.7-27.5)	2.6	(1.6-4.0)	3.8	(2.6-5.5)	2.3	(1.2-4.6)*
Sex												
	Men	1,346	60.0	(55.2-64.7)	28.9	(24.9-33.3)	2.6	(1.6-4.1)	6.1	(3.9-9.2)	2.4	(1.1-5.5)*
	Women	2,075	69.0	(65.6-72.3)	25.7	(22.6-29.1)	1.0	(0.6-1.7)	2.7	(1.8-4.1)	1.5	(0.7-3.1)*
Race/Ethnicity												
	Hispanic or Latino	245	53.5	(42.2-64.4)	26.3	(18.2-36.5)	1.3	(0.4-4.5)*	8.0	(4.5-13.7)	10.9	(6.2-18.5)
	Not Hispanic or Latino:	2,809	64.0	(60.9-67.1)	28.1	(25.2-31.2)	2.0	(1.4-2.9)	4.3	(3.0-6.3)	1.5	(0.9-2.6)
	White											
	Black or African American	98	70.5	(57.3-81.0)	24.4	(14.8-37.4)	0.5	(0.4-0.8)	1.7	(0.8-3.6)*	2.8	(0.8-10.0)*
	All Other Races	234	67.3	(55.9-77.1)	21.8	(14.4-31.5)	2.1	(0.8-5.7)*	4.1	(1.5-10.3)*	4.6	(1.9-11.1)*
Education												
	<High School	188	61.6	(48.2-73.4)	26.2	(17.0-38.3)	1.4	(0.4-5.4)*	9.5	(4.7-18.2)*	1.3	(0.3-4.7)*
	High School Graduate (or GED)	947	61.0	(55.1-66.5)	30.2	(25.1-36.0)	2.2	(1.2-4.1)*	4.0	(2.0-8.1)*	2.6	(0.9-7.2)*
	Some College	1,163	66.4	(61.2-71.2)	27.6	(23.2-32.5)	1.9	(1.0-3.4)*	2.7	(1.6-4.6)	1.4	(0.7-3.2)*
	College Graduate	1,118	67.5	(62.4-72.2)	24.9	(20.7-29.8)	1.0	(0.5-2.2)*	4.6	(2.5-8.1)	1.9	(0.6-5.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	438	67.8	(58.4-75.9)	21.2	(14.9-29.4)	1.0	(0.4-2.6)*	5.2	(2.4-10.8)*	4.8	(1.6-13.7)*
	\$20,000-\$34,999	625	69.8	(62.5-76.2)	22.3	(17.1-28.5)	1.2	(0.6-2.3)*	4.4	(2.5-7.7)	2.3	(0.8-6.8)*
	\$35,000-\$49,999	486	63.1	(54.6-71.0)	30.3	(23.2-38.6)	1.3	(0.4-3.9)*	3.1	(1.4-6.7)*	2.1	(0.8-5.1)*
	\$50,000-\$74,999	570	69.8	(63.5-75.4)	24.7	(19.5-30.7)	2.5	(1.0-5.8)*	2.7	(1.2-6.1)*	0.3	(0.1-0.9)*
	\$75,000 and more	915	63.2	(57.8-68.3)	29.5	(24.4-35.1)	1.1	(0.5-2.3)*	5.5	(2.8-10.4)*	0.7	(0.2-2.8)*
Employment Status	Employed	1,643	63.2	(59.0-67.1)	27.9	(24.3-31.9)	2.2	(1.4-3.6)	4.5	(2.9-6.8)	2.2	(0.9-5.4)*
	Out of work	313	72.3	(63.0-80.0)	22.6	(16.0-30.9)	0.5	(0.1-3.5)*	3.7	(1.2-10.7)*	1.0	(0.4-2.8)*
	Homemaker	210	62.1	(50.5-72.5)	33.7	(23.9-45.1)	0.5	(0.1-1.6)*	0.2	(0.0-1.5)*	3.5	(0.5-21.5)*
	Student	43	--	--	--	--	--	--	--	--	--	--
	Retired	1,025	63.2	(58.7-67.5)	26.8	(22.9-31.1)	2.5	(1.3-4.6)*	5.4	(3.5-8.2)	2.1	(0.8-5.1)*
	Unable to work	178	75.1	(64.2-83.6)	13.7	(7.4-23.9)*	2.1	(0.7-5.7)*	8.0	(3.2-18.3)*	1.2	(0.3-4.6)*
Veteran Status	Yes	567	61.8	(55.2-68.0)	30.7	(24.8-37.2)	1.6	(0.8-3.6)*	4.6	(2.2-9.2)*	1.3	(0.5-3.5)*
	No	2,853	65.3	(62.0-68.5)	26.6	(23.7-29.7)	1.8	(1.2-2.7)	4.3	(3.0-6.1)	2.0	(1.1-3.8)*
Diabetes	Yes	358	64.3	(56.7-71.2)	24.9	(18.8-32.2)	4.8	(2.3-9.7)*	4.1	(2.2-7.6)*	1.9	(0.8-4.7)*
	No	3,059	64.8	(61.6-67.9)	27.5	(24.7-30.4)	1.5	(1.0-2.2)	4.3	(3.1-6.1)	1.9	(1.0-3.6)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	416	60.6 (53.6-67.2)	29.1 (22.9-36.1)	2.3 (1.0-5.1)*	6.2 (3.4-11.2)*	1.8 (0.7-4.6)*				
	No	2,978	65.2 (62.0-68.2)	27.1 (24.3-30.1)	1.7 (1.2-2.6)	4.0 (2.8-5.7)	2.0 (1.0-3.7)*				
Asthma	Yes	462	66.5 (58.2-73.9)	26.0 (18.9-34.6)	0.6 (0.3-1.3)*	2.0 (0.9-4.3)*	4.9 (1.3-16.6)*				
	No	2,958	64.4 (61.2-67.5)	27.4 (24.6-30.4)	2.0 (1.3-2.9)	4.7 (3.4-6.6)	1.4 (0.9-2.3)				
Arthritis	Yes	1,182	69.2 (64.6-73.3)	24.3 (20.5-28.7)	1.6 (0.8-2.9)*	3.8 (2.2-6.5)	1.2 (0.6-2.2)*				
	No	2,232	63.3 (59.6-66.8)	28.2 (25.0-31.6)	1.8 (1.2-2.8)	4.5 (3.1-6.6)	2.2 (1.1-4.3)*				
Frequent Mental Distress	Yes	403	66.3 (58.1-73.6)	27.2 (20.5-35.1)	1.2 (0.5-2.9)*	4.2 (2.3-7.5)*	1.1 (0.4-3.3)*				
	No	2,979	64.7 (61.5-67.8)	27.4 (24.5-30.4)	1.7 (1.1-2.5)	4.2 (2.9-6.0)	2.1 (1.1-3.8)*				
Serious Psychological Distress	Yes	131	67.4 (53.7-78.6)	24.2 (13.9-38.9)	2.6 (0.9-7.7)*	2.6 (1.1-6.0)*	3.1 (1.1-8.8)*				
	No	3,237	64.5 (61.4-67.5)	27.6 (24.9-30.5)	1.6 (1.1-2.4)	4.3 (3.1-6.0)	1.9 (1.0-3.5)*				
Receiving Treatment for Mental Health Condition	Yes	369	74.8 (66.7-81.5)	21.1 (14.7-29.2)	1.1 (0.3-3.5)*	1.5 (0.6-4.1)*	1.5 (0.6-4.1)*				
	No	3,049	63.8 (60.6-66.8)	27.8 (25.0-30.8)	1.8 (1.3-2.6)	4.6 (3.3-6.4)	2.0 (1.1-3.6)*				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,420	22.0	(19.5-24.6)	34.6	(31.8-37.6)	2.3	(1.8-3.1)	28.9	(26.4-31.5)	12.2	(10.6-14.0)
Age (years)	18-24	86	20.5	(11.5-33.7)	45.1	(31.9-59.1)	0.7	(0.1-4.8)*	27.1	(17.0-40.4)	6.6	(3.1-13.6)*
	25-34	290	19.5	(13.3-27.8)	42.1	(33.7-51.1)	1.5	(0.7-3.4)*	24.6	(18.4-32.1)	12.2	(7.8-18.5)
	35-54	1,142	20.9	(17.6-24.6)	32.2	(28.3-36.2)	2.5	(1.5-4.2)	32.6	(28.7-36.6)	11.9	(9.4-14.9)
	55+	1,861	25.6	(22.8-28.6)	28.8	(25.9-31.8)	3.3	(2.3-4.7)	28.1	(25.2-31.2)	14.2	(12.1-16.7)
Sex	Men	1,347	25.1	(21.1-29.5)	38.3	(33.8-42.9)	2.4	(1.6-3.7)	23.9	(20.5-27.6)	10.3	(8.0-13.3)
	Women	2,073	19.0	(16.4-21.9)	31.0	(27.5-34.7)	2.3	(1.5-3.4)	33.9	(30.5-37.5)	13.9	(11.8-16.2)
Race/ Ethnicity	Hispanic or Latino	242	29.2	(21.0-39.1)	34.4	(24.9-45.4)	3.9	(2.0-7.6)*	23.0	(15.8-32.3)	9.4	(5.9-14.8)
	Not Hispanic or Latino:	2,811	18.8	(16.5-21.2)	34.9	(31.8-38.2)	2.4	(1.7-3.3)	31.9	(29.0-34.8)	12.1	(10.3-14.1)
	White											
	Black or African American	98	27.1	(17.7-39.3)	25.8	(15.6-39.5)	0.7	(0.5-1.0)	18.9	(9.7-33.5)*	27.5	(15.7-43.4)
	All Other Races	234	42.8	(30.5-56.0)	25.9	(17.2-37.0)	1.4	(0.5-3.9)*	20.4	(13.0-30.5)	9.6	(5.8-15.4)
Education	<High School	188	30.3	(19.9-43.1)	25.3	(16.1-37.4)	2.8	(1.0-7.3)*	34.4	(23.8-46.7)	7.3	(4.1-12.7)
	High School Graduate (or GED)	947	21.1	(17.0-26.0)	40.7	(34.8-46.9)	1.8	(1.0-3.2)	26.3	(21.8-31.4)	10.0	(7.5-13.1)
	Some College	1,172	23.2	(19.1-27.9)	32.4	(27.8-37.4)	3.0	(1.9-4.6)	26.8	(22.8-31.2)	14.7	(11.4-18.7)
	College Graduate	1,107	18.8	(14.9-23.4)	33.8	(29.2-38.7)	1.9	(1.0-3.4)*	32.2	(27.9-36.8)	13.3	(10.5-16.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	440	28.0	(20.0-37.5)	25.0	(18.5-33.0)	3.2	(1.4-7.1)*	26.4	(19.3-34.9)	17.5	(12.3-24.3)
	\$20,000-\$34,999	633	18.2	(13.5-24.0)	38.1	(30.2-46.6)	2.2	(1.1-4.2)*	28.7	(22.5-35.9)	12.8	(9.1-17.9)
	\$35,000-\$49,999	480	24.6	(18.0-32.7)	34.7	(27.5-42.8)	2.6	(1.4-4.7)*	28.5	(22.3-35.5)	9.6	(6.8-13.5)
	\$50,000-\$74,999	569	20.3	(15.1-26.6)	34.7	(28.3-41.6)	2.4	(1.2-4.7)*	27.2	(21.9-33.2)	15.5	(11.0-21.4)
	\$75,000 and more	916	20.7	(16.7-25.5)	37.7	(32.7-43.0)	1.6	(0.8-3.2)*	30.3	(25.9-35.1)	9.7	(7.1-13.0)
Employment Status	Employed	1,633	22.6	(19.2-26.5)	34.4	(30.5-38.5)	2.5	(1.7-3.8)	29.1	(25.7-32.7)	11.4	(9.2-13.9)
	Out of work	311	18.6	(12.1-27.4)	36.2	(27.2-46.3)	0.5	(0.1-3.0)*	29.2	(21.4-38.4)	15.5	(10.5-22.3)
	Homemaker	215	18.7	(12.4-27.4)	41.9	(31.3-53.3)	2.5	(0.9-6.5)*	28.1	(19.4-38.8)	8.8	(5.2-14.6)
	Student	44	--	--	--	--	--	--	--	--	--	--
	Retired	1,032	25.9	(22.0-30.3)	32.6	(28.3-37.1)	3.2	(2.0-5.3)	28.1	(24.3-32.3)	10.1	(7.8-13.2)
	Unable to work	177	16.4	(10.2-25.4)	26.8	(17.4-39.0)	2.4	(0.9-6.4)*	30.3	(20.7-41.9)	24.1	(13.8-38.6)
Veteran Status	Yes	567	22.2	(17.5-27.8)	29.6	(24.0-36.0)	1.2	(0.6-2.2)*	29.4	(23.6-35.9)	17.6	(11.8-25.4)
	No	2,852	21.9	(19.2-24.8)	35.5	(32.3-38.8)	2.6	(1.9-3.5)	28.7	(26.0-31.6)	11.4	(9.8-13.2)
Diabetes	Yes	355	21.5	(15.9-28.5)	27.8	(20.9-35.9)	3.0	(1.3-6.9)*	29.7	(22.3-38.4)	17.9	(12.4-25.3)
	No	3,061	22.1	(19.5-24.9)	35.3	(32.2-38.4)	2.2	(1.6-2.9)	28.8	(26.2-31.6)	11.7	(10.0-13.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Nevada, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	417	19.5	(14.5-25.6)	29.7	(23.9-36.2)	2.1	(0.8-5.1)*	35.5	(28.8-42.8)	13.3	(9.5-18.2)
	No	2,977	22.2	(19.6-25.1)	34.9	(31.9-38.1)	2.4	(1.7-3.2)	28.3	(25.7-31.1)	12.1	(10.4-14.1)
Asthma	Yes	468	18.2	(12.8-25.2)	35.9	(28.4-44.2)	1.5	(0.6-3.4)*	29.7	(23.3-37.0)	14.7	(10.8-19.7)
	No	2,951	22.6	(20.0-25.5)	34.5	(31.4-37.7)	2.5	(1.8-3.4)	28.7	(26.0-31.5)	11.8	(10.0-13.8)
Arthritis	Yes	1,182	18.2	(14.6-22.3)	28.5	(24.4-33.1)	2.8	(1.6-4.8)	33.1	(28.9-37.6)	17.4	(14.2-21.3)
	No	2,233	23.5	(20.5-26.7)	36.6	(33.1-40.2)	2.2	(1.6-3.1)	27.4	(24.5-30.5)	10.4	(8.5-12.5)
Frequent Mental Distress	Yes	402	15.4	(9.9-23.2)	33.3	(25.2-42.5)	4.1	(2.0-8.0)*	25.9	(19.6-33.4)	21.3	(15.6-28.4)
	No	2,980	22.9	(20.3-25.7)	34.8	(31.8-38.0)	2.0	(1.5-2.8)	29.4	(26.8-32.3)	10.8	(9.2-12.8)
Serious Psychological Distress	Yes	130	13.1	(6.5-24.5)*	22.2	(14.0-33.5)	0.7	(0.1-3.5)*	36.0	(23.3-51.0)	28.0	(17.4-41.8)
	No	3,242	21.9	(19.5-24.6)	35.6	(32.7-38.7)	2.5	(1.8-3.3)	28.5	(26.0-31.1)	11.5	(9.9-13.4)
Receiving Treatment for Mental Health Condition	Yes	368	13.6	(9.6-18.8)	20.0	(14.2-27.4)	1.2	(0.5-2.8)*	35.5	(27.6-44.4)	29.7	(22.3-38.4)
	No	3,049	23.1	(20.5-25.9)	35.5	(32.5-38.6)	2.5	(1.8-3.3)	28.1	(25.5-30.8)	10.9	(9.3-12.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,310	67.0	(65.3-68.7)	25.9	(24.3-27.5)	2.1	(1.7-2.7)	3.7	(2.9-4.6)	1.3	(0.9-1.8)
Age (years)	18-24	164	55.3	(46.6-63.7)	40.0	(31.7-48.9)	2.5	(0.9-6.8)*	2.2	(0.8-6.2)*	0.0	(0.0-0.0)
	25-34	472	63.8	(58.5-68.7)	27.5	(23.0-32.4)	1.6	(0.7-3.6)*	6.5	(3.9-10.7)	0.6	(0.1-3.1)*
	35-54	2,094	68.4	(66.0-70.8)	24.3	(22.1-26.6)	2.3	(1.6-3.2)	3.6	(2.7-4.9)	1.4	(0.8-2.2)
	55+	2,530	71.6	(69.5-73.6)	21.7	(19.9-23.7)	2.2	(1.6-3.0)	2.6	(2.0-3.4)	1.9	(1.2-2.9)
Sex	Men	2,048	60.9	(58.2-63.6)	30.4	(27.8-33.0)	2.7	(1.9-3.7)	4.8	(3.6-6.6)	1.2	(0.7-2.0)
	Women	3,262	72.7	(70.7-74.7)	21.7	(19.9-23.7)	1.6	(1.2-2.2)	2.6	(2.0-3.4)	1.4	(0.9-2.1)
Race/ Ethnicity	Hispanic or Latino	78	56.5	(43.6-68.6)	35.4	(23.6-49.3)	2.5	(0.4-15.9)*	4.3	(0.6-23.5)*	1.3	(0.2-6.3)*
	Not Hispanic or Latino:	5,014	68.1	(66.4-69.8)	25.6	(24.0-27.2)	2.1	(1.6-2.6)	3.3	(2.6-4.1)	0.9	(0.7-1.3)
	White											
	Black or African American	21	--	--	--	--	--	--	--	--	--	--
	All Other Races	155	63.1	(53.2-72.0)	26.9	(19.2-36.3)	2.9	(1.2-7.2)*	4.4	(1.3-13.6)*	2.7	(0.6-11.3)*
Education	<High School	304	61.0	(53.5-68.0)	27.6	(21.2-35.1)	1.2	(0.5-3.0)*	9.3	(5.7-14.8)	0.9	(0.4-2.4)*
	High School Graduate (or GED)	1,451	61.9	(58.6-65.1)	29.4	(26.4-32.7)	2.8	(1.8-4.3)	4.1	(3.0-5.7)	1.7	(1.0-2.7)
	Some College	1,333	66.0	(62.5-69.4)	26.3	(23.2-29.6)	2.7	(1.8-3.9)	3.5	(2.1-5.9)	1.5	(0.8-2.8)*
	College Graduate	2,214	73.4	(70.9-75.7)	22.3	(20.1-24.7)	1.3	(0.9-2.0)	2.2	(1.4-3.3)	0.8	(0.4-1.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	632	63.2	(57.8-68.4)	26.3	(21.5-31.7)	2.5	(1.4-4.5)*	4.5	(2.9-6.8)	3.5	(2.2-5.6)
	\$20,000-\$34,999	838	62.9	(58.4-67.1)	29.7	(25.6-34.1)	2.0	(1.0-3.7)*	4.6	(2.9-7.1)	0.9	(0.3-2.4)*
	\$35,000-\$49,999	784	65.1	(60.7-69.2)	26.7	(22.9-30.9)	2.7	(1.6-4.4)	3.9	(2.3-6.4)	1.7	(0.8-3.6)*
	\$50,000-\$74,999	843	71.2	(67.2-74.9)	25.0	(21.5-28.8)	1.4	(0.7-2.7)*	2.0	(0.9-4.2)*	0.4	(0.1-1.4)*
	\$75,000 and more	1,548	70.9	(67.9-73.8)	22.5	(19.9-25.3)	2.1	(1.4-3.1)	3.9	(2.5-6.1)	0.6	(0.2-1.8)*
Employment Status	Employed	3,098	67.0	(64.9-69.1)	25.6	(23.6-27.7)	2.4	(1.8-3.2)	4.1	(3.1-5.4)	0.9	(0.5-1.5)
	Out of work	180	59.7	(50.1-68.7)	32.8	(24.4-42.5)	1.0	(0.4-2.9)*	3.8	(1.6-9.0)*	2.6	(0.9-7.6)*
	Homemaker	325	65.5	(58.7-71.8)	27.4	(21.4-34.2)	2.8	(1.3-6.0)*	1.7	(0.7-4.2)*	2.6	(0.8-8.7)*
	Student	83	72.4	(60.6-81.7)	23.8	(15.2-35.3)	1.8	(0.4-8.1)*	2.0	(0.5-7.4)*	0.0	(0.0-0.0)
	Retired	1,322	68.3	(65.3-71.2)	25.4	(22.7-28.3)	1.7	(1.1-2.6)	2.7	(1.9-3.9)	2.0	(1.1-3.4)
	Unable to work	294	66.2	(59.2-72.6)	24.4	(18.9-30.9)	1.6	(0.6-4.7)*	5.9	(3.3-10.6)*	1.8	(0.8-4.0)*
Veteran Status	Yes	748	63.2	(59.1-67.2)	27.8	(24.1-31.8)	2.6	(1.6-4.2)	3.7	(2.2-6.1)	2.7	(1.4-5.1)*
	No	4,556	67.6	(65.8-69.4)	25.6	(23.8-27.4)	2.1	(1.6-2.7)	3.7	(2.9-4.7)	1.0	(0.7-1.5)
Diabetes	Yes	496	64.6	(59.5-69.4)	28.5	(23.8-33.7)	2.2	(1.2-3.8)	3.0	(1.7-5.2)	1.7	(0.8-3.7)*
	No	4,810	67.2	(65.4-68.9)	25.7	(24.1-27.4)	2.1	(1.7-2.7)	3.7	(3.0-4.7)	1.3	(0.9-1.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	554	68.4	(63.4-73.0)	24.3	(20.0-29.1)	1.5	(0.8-3.0)*	3.2	(1.9-5.4)	2.6	(1.1-5.9)*
	No	4,722	66.8	(65.0-68.6)	26.1	(24.4-27.9)	2.2	(1.7-2.8)	3.7	(2.9-4.7)	1.2	(0.8-1.7)
Asthma	Yes	770	67.9	(63.3-72.2)	25.1	(21.1-29.6)	1.8	(1.0-3.1)	4.6	(2.5-8.2)*	0.7	(0.3-1.3)*
	No	4,526	66.8	(64.9-68.5)	26.0	(24.3-27.8)	2.2	(1.7-2.8)	3.6	(2.8-4.5)	1.5	(1.0-2.1)
Arthritis	Yes	1,955	69.7	(67.0-72.3)	23.4	(21.0-25.9)	1.9	(1.3-2.7)	3.0	(2.1-4.3)	2.1	(1.2-3.4)
	No	3,334	65.9	(63.8-68.0)	26.9	(24.9-28.9)	2.3	(1.7-3.0)	3.9	(3.0-5.1)	1.0	(0.6-1.5)
Frequent Mental Distress	Yes	517	59.0	(53.3-64.4)	29.0	(24.0-34.6)	2.5	(1.2-5.0)*	6.6	(3.7-11.6)	2.9	(1.6-5.5)*
	No	4,736	67.9	(66.1-69.6)	25.7	(24.1-27.5)	2.1	(1.7-2.7)	3.3	(2.6-4.1)	1.0	(0.7-1.5)
Serious Psychological Distress	Yes	179	55.5	(46.3-64.3)	26.7	(18.8-36.3)	2.9	(1.1-7.6)*	11.7	(7.0-18.9)	3.3	(1.3-8.1)*
	No	5,077	67.4	(65.7-69.1)	25.9	(24.3-27.6)	2.1	(1.7-2.7)	3.4	(2.7-4.3)	1.2	(0.8-1.7)
Receiving Treatment for Mental Health Condition	Yes	792	76.9	(72.7-80.6)	16.9	(13.6-20.9)	2.0	(1.0-3.9)*	3.1	(1.8-5.4)	1.0	(0.5-2.2)*
	No	4,508	65.6	(63.7-67.4)	27.2	(25.5-29.0)	2.1	(1.7-2.7)	3.8	(3.0-4.8)	1.3	(0.9-1.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,296	21.5	(20.1-23.0)	38.2	(36.5-39.9)	3.6	(3.0-4.3)	26.1	(24.7-27.6)	10.6	(9.6-11.7)
Age (years)	18-24	166	14.5	(9.6-21.4)	45.2	(36.8-54.0)	3.8	(1.7-8.3)*	27.1	(20.1-35.4)	9.4	(5.2-16.3)
	25-34	473	16.6	(13.0-20.8)	43.5	(38.6-48.5)	4.2	(2.7-6.6)	26.9	(22.6-31.6)	8.9	(6.3-12.2)
	35-54	2,093	19.4	(17.4-21.7)	38.2	(35.8-40.6)	3.2	(2.4-4.1)	27.5	(25.4-29.7)	11.7	(10.3-13.4)
	55+	2,515	29.6	(27.5-31.7)	32.5	(30.4-34.7)	3.7	(3.0-4.7)	23.6	(21.8-25.4)	10.6	(9.3-12.0)
Sex	Men	2,047	22.7	(20.5-24.9)	43.3	(40.6-46.0)	4.0	(3.2-5.2)	22.4	(20.3-24.7)	7.6	(6.2-9.1)
	Women	3,249	20.5	(18.8-22.3)	33.3	(31.3-35.3)	3.2	(2.5-4.0)	29.7	(27.7-31.6)	13.4	(12.0-14.9)
Race/ Ethnicity	Hispanic or Latino	74	19.1	(9.9-33.8)*	31.8	(20.6-45.6)	6.7	(2.7-16.0)*	17.7	(10.2-28.9)	24.6	(14.7-38.2)
	Not Hispanic or Latino:	4,999	18.9	(17.7-20.3)	39.3	(37.6-41.0)	3.6	(3.0-4.3)	27.8	(26.3-29.4)	10.3	(9.3-11.4)
	White											
	Black or African American	21	--	--	--	--	--	--	--	--	--	--
	All Other Races	158	26.5	(18.7-36.2)	32.8	(24.5-42.2)	4.6	(2.0-10.4)*	24.3	(17.0-33.5)	11.8	(7.5-18.2)
Education	<High School	313	33.7	(27.0-41.1)	29.1	(22.5-36.7)	2.3	(1.2-4.7)*	20.0	(14.5-26.9)	14.9	(10.0-21.6)
	High School Graduate (or GED)	1,450	26.7	(23.9-29.8)	35.8	(32.6-39.1)	3.7	(2.6-5.2)	23.3	(20.5-26.4)	10.5	(8.8-12.5)
	Some College	1,324	18.4	(16.0-21.1)	39.9	(36.6-43.3)	3.8	(2.7-5.3)	27.1	(24.2-30.3)	10.7	(8.9-12.9)
	College Graduate	2,200	15.5	(13.7-17.4)	40.7	(38.1-43.2)	3.7	(2.9-4.7)	29.8	(27.6-32.2)	10.4	(8.8-12.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	630	25.5 (21.3-30.1)	28.5 (23.8-33.8)	4.1 (2.1-7.8)*	25.3 (20.9-30.3)	16.6 (13.3-20.6)				
	\$20,000-\$34,999	835	23.7 (20.3-27.5)	35.3 (31.2-39.6)	3.1 (2.0-4.7)	25.4 (21.7-29.4)	12.6 (9.9-15.9)				
	\$35,000-\$49,999	783	21.1 (17.8-24.8)	37.6 (33.4-42.0)	3.1 (2.0-4.9)	26.9 (23.2-31.0)	11.3 (8.7-14.5)				
	\$50,000-\$74,999	847	21.5 (18.2-25.2)	40.0 (36.1-44.1)	5.0 (3.6-6.9)	24.2 (21.0-27.7)	9.3 (7.3-11.8)				
	\$75,000 and more	1,538	18.3 (16.0-20.9)	40.8 (37.9-43.8)	3.4 (2.5-4.6)	29.1 (26.5-31.8)	8.4 (6.9-10.2)				
Employment Status	Employed	3,094	19.8 (18.1-21.7)	41.3 (39.1-43.4)	3.3 (2.6-4.1)	25.9 (24.0-27.7)	9.8 (8.6-11.1)				
	Out of work	180	15.5 (10.3-22.7)	30.3 (21.7-40.5)	2.8 (0.9-8.1)*	30.6 (22.2-40.5)	20.8 (13.4-30.9)				
	Homemaker	323	27.7 (21.9-34.4)	34.9 (29.1-41.2)	3.5 (1.8-6.5)*	24.5 (19.4-30.4)	9.4 (6.5-13.4)				
	Student	83	14.7 (7.3-27.4)*	41.3 (29.3-54.6)	6.5 (2.2-17.5)*	27.0 (17.4-39.2)	10.5 (4.0-25.2)*				
	Retired	1,311	28.7 (26.0-31.6)	32.6 (29.7-35.7)	4.4 (3.3-5.9)	25.8 (23.2-28.6)	8.4 (6.9-10.3)				
	Unable to work	298	22.0 (16.9-28.3)	22.1 (16.9-28.5)	4.8 (2.6-8.7)*	30.6 (24.2-37.8)	20.5 (15.7-26.2)				
Veteran Status	Yes	749	26.5 (23.2-30.1)	37.0 (33.0-41.1)	3.5 (2.4-5.2)	23.0 (19.5-26.8)	10.0 (7.7-12.8)				
	No	4,541	20.6 (19.1-22.2)	38.5 (36.7-40.3)	3.6 (3.0-4.4)	26.6 (25.0-28.2)	10.6 (9.6-11.8)				
Diabetes	Yes	502	23.1 (19.1-27.6)	36.3 (31.3-41.7)	4.4 (2.8-6.9)	22.6 (18.8-27.0)	13.5 (10.4-17.4)				
	No	4,791	21.4 (19.9-22.9)	38.3 (36.6-40.1)	3.6 (3.0-4.3)	26.4 (24.8-27.9)	10.4 (9.3-11.5)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Hampshire, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	543	24.6 (20.9-28.8)	30.7 (26.0-35.8)	4.1 (2.7-6.2)	28.1 (23.6-33.1)	12.5 (9.6-16.1)				
	No	4,720	21.1 (19.6-22.6)	39.0 (37.2-40.8)	3.5 (3.0-4.3)	26.0 (24.5-27.6)	10.3 (9.3-11.5)				
Asthma	Yes	764	16.0 (13.1-19.4)	36.2 (31.9-40.8)	3.7 (2.4-5.7)	29.0 (25.0-33.4)	15.0 (12.2-18.3)				
	No	4,518	22.5 (21.0-24.1)	38.6 (36.8-40.4)	3.6 (3.0-4.3)	25.6 (24.0-27.2)	9.7 (8.7-10.9)				
Arthritis	Yes	1,943	22.4 (20.3-24.6)	33.6 (31.1-36.2)	3.0 (2.3-4.0)	27.6 (25.2-30.0)	13.4 (11.6-15.4)				
	No	3,332	21.1 (19.3-22.9)	39.9 (37.8-42.0)	3.8 (3.1-4.7)	25.7 (23.9-27.6)	9.5 (8.4-10.8)				
Frequent Mental Distress	Yes	514	18.1 (14.3-22.8)	24.7 (19.9-30.2)	3.4 (1.9-6.1)	29.6 (24.7-35.1)	24.1 (19.3-29.6)				
	No	4,727	22.0 (20.5-23.6)	39.6 (37.8-41.4)	3.6 (3.0-4.3)	25.7 (24.2-27.3)	9.1 (8.1-10.1)				
Serious Psychological Distress	Yes	180	13.0 (8.6-19.2)	20.1 (13.1-29.6)	3.1 (1.0-8.8)*	30.7 (22.6-40.2)	33.2 (24.6-43.0)				
	No	5,059	21.9 (20.5-23.4)	38.9 (37.2-40.7)	3.6 (3.0-4.2)	25.8 (24.4-27.4)	9.7 (8.8-10.8)				
Receiving Treatment for Mental Health Condition	Yes	791	16.9 (13.9-20.5)	33.0 (28.8-37.4)	4.2 (2.8-6.3)	30.1 (26.3-34.2)	15.8 (13.0-19.2)				
	No	4,494	22.2 (20.7-23.8)	38.8 (37.0-40.7)	3.5 (2.9-4.2)	25.6 (24.0-27.2)	9.8 (8.8-11.0)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,672	67.0	(65.1-68.7)	25.1	(23.4-26.9)	2.5	(2.0-3.1)	3.9	(3.2-4.7)	1.6	(1.1-2.3)
Age (years)	18-24	253	53.0	(45.3-60.5)	39.2	(32.0-46.9)	0.5	(0.2-1.8)*	4.4	(2.4-7.8)	2.9	(1.1-7.7)*
	25-34	638	62.0	(57.1-66.7)	29.2	(24.9-33.8)	1.4	(0.5-3.7)*	6.2	(4.0-9.3)	1.2	(0.6-2.4)*
	35-54	2,125	69.5	(66.8-72.0)	23.4	(21.1-25.9)	2.7	(1.9-3.9)	3.1	(2.3-4.1)	1.3	(0.9-2.0)
	55+	2,642	72.3	(70.2-74.4)	19.3	(17.5-21.2)	3.6	(2.7-4.6)	3.4	(2.6-4.4)	1.4	(1.0-2.0)
Sex	Men	2,081	60.2	(57.2-63.2)	30.4	(27.6-33.3)	2.8	(2.1-3.8)	4.4	(3.3-5.9)	2.2	(1.3-3.6)
	Women	3,591	73.2	(71.1-75.1)	20.3	(18.5-22.1)	2.1	(1.6-2.9)	3.3	(2.6-4.3)	1.1	(0.7-1.7)
Race/ Ethnicity	Hispanic or Latino	1,789	68.3	(65.2-71.2)	24.0	(21.3-27.0)	2.2	(1.6-3.2)	3.9	(2.8-5.3)	1.6	(1.0-2.4)
	Not Hispanic or Latino:	3,296	67.0	(64.6-69.2)	25.2	(23.0-27.4)	2.6	(2.0-3.4)	3.2	(2.4-4.3)	2.1	(1.2-3.7)
	White											
	Black or African American	50	85.2	(72.7-92.6)	13.4	(6.4-25.9)*	0.1	(0.1-0.2)	0.2	(0.1-0.3)	1.1	(0.1-7.5)*
	All Other Races	507	58.3	(51.9-64.4)	31.2	(25.8-37.3)	2.9	(1.0-7.6)*	5.9	(4.0-8.7)	1.7	(0.9-3.3)*
Education	<High School	678	67.0	(61.7-72.0)	23.4	(19.1-28.4)	4.1	(2.3-7.0)	3.3	(2.0-5.3)	2.2	(1.1-4.1)*
	High School Graduate (or GED)	1,506	65.6	(61.9-69.1)	25.5	(22.3-29.1)	2.8	(1.9-4.0)	4.7	(3.5-6.5)	1.4	(0.9-2.1)
	Some College	1,520	63.4	(59.9-66.8)	28.0	(24.8-31.5)	2.4	(1.7-3.6)	4.2	(2.9-6.1)	1.9	(0.8-4.5)*
	College Graduate	1,962	71.2	(68.4-73.8)	22.9	(20.5-25.5)	1.7	(1.1-2.7)	2.8	(1.9-4.3)	1.3	(0.8-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,149	67.0 (62.8-70.9)	23.4 (20.0-27.3)	3.5 (2.2-5.5)	4.2 (2.9-6.0)	1.9 (1.2-2.9)				
	\$20,000-\$34,999	1,176	65.1 (61.3-68.8)	26.1 (22.7-29.8)	2.1 (1.3-3.3)	5.2 (3.5-7.5)	1.5 (0.5-4.2)*				
	\$35,000-\$49,999	867	67.9 (63.5-72.1)	26.7 (22.7-31.1)	1.4 (0.8-2.6)	3.2 (1.9-5.4)	0.7 (0.3-1.6)*				
	\$50,000-\$74,999	861	66.4 (61.9-70.6)	26.7 (22.6-31.3)	1.6 (0.9-2.9)*	3.0 (1.7-5.2)	2.3 (0.9-6.1)*				
	\$75,000 and more	1,113	69.2 (65.1-72.9)	24.1 (20.7-28.0)	2.8 (1.7-4.7)	2.6 (1.5-4.4)	1.3 (0.7-2.3)*				
Employment Status	Employed	3,115	66.8 (64.5-69.1)	25.0 (22.9-27.2)	2.4 (1.7-3.3)	4.2 (3.2-5.5)	1.6 (0.9-2.7)				
	Out of work	212	60.3 (50.3-69.4)	32.3 (23.7-42.3)	1.9 (0.6-5.7)*	2.9 (1.3-6.2)*	2.7 (0.6-10.4)*				
	Homemaker	544	68.6 (63.2-73.5)	22.1 (17.8-27.0)	1.7 (1.0-3.1)	6.1 (3.5-10.4)	1.5 (0.7-3.1)*				
	Student	129	64.6 (53.6-74.2)	32.8 (23.4-43.8)	0.3 (0.1-1.8)*	1.1 (0.3-3.9)*	1.2 (0.2-7.1)*				
	Retired	1,313	68.0 (64.5-71.2)	23.8 (20.7-27.2)	3.5 (2.5-5.0)	3.6 (2.5-5.3)	1.1 (0.6-2.0)*				
	Unable to work	347	73.1 (66.2-79.0)	17.7 (12.8-23.9)	3.8 (1.8-7.8)*	3.3 (2.0-5.5)	2.2 (1.1-4.1)*				
Veteran Status	Yes	780	65.2 (60.8-69.3)	26.2 (22.3-30.4)	3.2 (2.0-5.1)	4.5 (3.1-6.7)	0.9 (0.4-2.1)*				
	No	4,891	67.3 (65.3-69.3)	24.9 (23.1-26.8)	2.3 (1.8-3.0)	3.7 (3.0-4.6)	1.7 (1.1-2.5)				
Diabetes	Yes	557	69.7 (63.9-75.0)	22.5 (17.5-28.3)	3.6 (1.9-6.5)*	2.8 (1.7-4.5)	1.4 (0.7-3.0)*				
	No	5,113	66.7 (64.8-68.5)	25.3 (23.6-27.1)	2.4 (1.9-3.0)	4.0 (3.2-4.9)	1.7 (1.1-2.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	535	65.4 (59.8-70.6)	26.0 (21.3-31.4)	2.4 (1.4-4.1)	4.3 (2.7-6.8)	1.9 (1.0-3.6)*				
	No	5,075	66.9 (65.0-68.8)	25.1 (23.4-27.0)	2.5 (2.0-3.2)	3.8 (3.1-4.7)	1.6 (1.1-2.4)				
Asthma	Yes	794	69.6 (64.8-73.9)	22.0 (18.0-26.5)	2.0 (1.1-3.6)	4.6 (2.7-7.8)	1.8 (0.8-4.2)*				
	No	4,867	66.5 (64.5-68.4)	25.6 (23.8-27.5)	2.6 (2.0-3.2)	3.8 (3.1-4.6)	1.6 (1.1-2.4)				
Arthritis	Yes	1,998	69.2 (66.2-72.0)	22.5 (20.0-25.2)	2.3 (1.6-3.4)	4.5 (3.2-6.2)	1.5 (1.0-2.3)				
	No	3,661	66.1 (63.9-68.3)	26.0 (24.0-28.2)	2.6 (2.0-3.3)	3.6 (2.8-4.6)	1.6 (1.0-2.6)				
Frequent Mental Distress	Yes	598	64.5 (58.5-70.0)	22.6 (17.8-28.3)	3.0 (1.5-5.6)*	5.2 (3.4-7.9)	4.7 (2.2-9.9)*				
	No	5,016	67.4 (65.4-69.2)	25.4 (23.7-27.3)	2.4 (1.9-3.1)	3.6 (2.9-4.4)	1.2 (0.8-1.8)				
Serious Psychological Distress	Yes	231	56.2 (46.8-65.2)	28.0 (20.0-37.7)	2.3 (0.7-7.8)*	4.3 (2.3-7.8)*	9.2 (5.4-15.4)				
	No	5,367	67.3 (65.4-69.1)	25.0 (23.3-26.8)	2.4 (1.9-3.0)	3.9 (3.1-4.7)	1.5 (1.0-2.1)				
Receiving Treatment for Mental Health Condition	Yes	667	77.7 (72.6-82.1)	16.4 (12.5-21.4)	1.6 (0.7-3.4)*	2.8 (1.5-5.0)*	1.5 (0.8-2.7)				
	No	5,000	65.6 (63.7-67.6)	26.2 (24.4-28.0)	2.6 (2.1-3.3)	4.0 (3.2-4.9)	1.6 (1.1-2.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,723	23.7	(22.1-25.2)	33.4	(31.6-35.3)	2.6	(2.1-3.3)	28.8	(27.1-30.6)	11.4	(10.3-12.7)
Age (years)	18-24	256	17.9	(13.1-24.1)	43.6	(36.0-51.4)	3.3	(1.3-8.1)*	28.0	(21.1-36.1)	7.2	(3.8-13.1)*
	25-34	655	23.1	(19.5-27.1)	36.4	(31.8-41.3)	1.5	(0.7-3.2)*	28.6	(24.5-33.1)	10.4	(7.5-14.2)
	35-54	2,150	23.8	(21.7-26.1)	32.3	(29.8-34.8)	1.8	(1.2-2.6)	29.5	(27.0-32.1)	12.6	(10.9-14.5)
	55+	2,649	27.0	(25.0-29.2)	29.3	(27.2-31.5)	3.8	(3.0-4.8)	27.9	(25.8-30.1)	12.0	(10.6-13.7)
Sex	Men	2,119	25.0	(22.7-27.6)	37.3	(34.5-40.3)	2.3	(1.7-3.2)	26.8	(24.2-29.5)	8.6	(7.0-10.5)
	Women	3,604	22.4	(20.6-24.2)	29.7	(27.6-31.9)	2.9	(2.1-4.1)	30.9	(28.7-33.1)	14.1	(12.6-15.9)
Race/ Ethnicity	Hispanic or Latino	1,825	36.2	(33.2-39.3)	30.1	(27.2-33.3)	2.4	(1.6-3.6)	21.9	(19.3-24.7)	9.4	(7.7-11.4)
	Not Hispanic or Latino:	3,306	18.7	(16.9-20.6)	35.9	(33.6-38.2)	2.4	(1.9-3.0)	31.6	(29.4-33.9)	11.4	(10.1-12.9)
	White											
	Black or African American	51	19.6	(10.6-33.4)	18.6	(7.9-37.5)*	17.7	(3.0-60.0)*	29.2	(13.9-51.4)*	14.9	(6.5-30.6)*
	All Other Races	510	24.4	(19.5-30.0)	34.2	(28.3-40.7)	2.8	(1.3-6.0)*	26.0	(21.2-31.4)	12.6	(8.5-18.4)
Education	<High School	704	43.0	(37.9-48.2)	31.2	(25.9-37.0)	4.7	(2.9-7.7)	12.7	(9.7-16.5)	8.4	(5.1-13.6)
	High School Graduate (or GED)	1,524	27.1	(24.0-30.5)	34.9	(31.3-38.6)	2.3	(1.5-3.5)	26.8	(23.3-30.6)	8.9	(7.2-11.1)
	Some College	1,529	19.3	(16.5-22.4)	35.0	(31.5-38.6)	2.3	(1.4-3.6)	29.6	(26.4-33.0)	13.9	(11.6-16.6)
	College Graduate	1,960	17.1	(15.1-19.3)	32.9	(30.2-35.6)	2.6	(1.6-4.2)	34.8	(32.1-37.6)	12.6	(10.9-14.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,167	28.4 (25.0-32.1)	30.9 (26.9-35.3)	3.3 (2.1-5.1)	25.6 (21.9-29.6)	11.8 (9.6-14.4)				
	\$20,000-\$34,999	1,189	30.3 (26.7-34.1)	29.4 (25.9-33.2)	2.4 (1.6-3.7)	23.9 (20.8-27.3)	14.0 (11.4-17.1)				
	\$35,000-\$49,999	878	20.3 (17.3-23.7)	33.6 (29.4-38.1)	1.9 (1.1-3.2)	32.4 (27.9-37.2)	11.9 (9.2-15.1)				
	\$50,000-\$74,999	865	17.5 (14.5-21.0)	34.3 (29.9-38.9)	1.9 (1.0-3.8)*	33.5 (28.9-38.5)	12.8 (9.9-16.3)				
	\$75,000 and more	1,112	20.8 (17.6-24.5)	37.5 (33.7-41.5)	2.1 (1.0-4.6)*	30.2 (27.0-33.6)	9.3 (7.4-11.6)				
Employment Status	Employed	3,157	23.5 (21.6-25.6)	33.5 (31.3-35.9)	1.7 (1.2-2.3)	29.3 (27.2-31.5)	11.9 (10.4-13.7)				
	Out of work	218	18.5 (12.9-25.9)	35.4 (25.9-46.2)	2.2 (0.9-5.2)*	34.5 (23.5-47.5)	9.4 (6.0-14.4)				
	Homemaker	555	32.7 (27.9-37.8)	28.7 (24.0-33.9)	2.9 (1.6-5.2)*	27.0 (22.6-32.0)	8.7 (6.4-11.6)				
	Student	128	21.9 (13.6-33.5)	43.5 (32.4-55.3)	5.7 (1.5-18.7)*	20.6 (12.6-32.0)	8.3 (3.4-18.8)*				
	Retired	1,303	22.7 (20.1-25.6)	32.6 (29.3-36.0)	4.7 (3.5-6.4)	30.5 (27.2-34.1)	9.5 (7.8-11.6)				
	Unable to work	348	22.4 (17.6-28.2)	28.2 (22.0-35.4)	3.7 (1.5-8.7)*	24.7 (19.0-31.5)	20.9 (15.8-27.3)				
Veteran Status	Yes	793	20.4 (17.4-23.7)	35.3 (31.2-39.7)	2.8 (1.8-4.4)	32.4 (28.1-37.0)	9.1 (6.9-12.0)				
	No	4,929	24.3 (22.6-26.0)	33.1 (31.1-35.1)	2.6 (2.0-3.4)	28.3 (26.4-30.2)	11.8 (10.5-13.1)				
Diabetes	Yes	570	25.2 (21.2-29.6)	28.4 (23.2-34.1)	3.7 (2.2-6.4)	30.3 (24.3-37.0)	12.4 (9.4-16.2)				
	No	5,151	23.5 (21.9-25.2)	33.8 (31.9-35.7)	2.5 (2.0-3.3)	28.7 (27.0-30.6)	11.4 (10.2-12.7)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

New Mexico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	544	25.3	(20.9-30.3)	24.6	(20.0-29.8)	3.5	(2.2-5.5)	31.6	(26.9-36.6)	15.1	(11.6-19.4)
	No	5,118	23.5	(21.9-25.2)	34.1	(32.2-36.1)	2.6	(2.0-3.3)	28.7	(26.9-30.6)	11.1	(9.9-12.5)
Asthma	Yes	779	20.6	(17.1-24.6)	27.9	(23.7-32.5)	2.5	(1.4-4.6)*	35.7	(30.9-40.8)	13.3	(10.2-17.2)
	No	4,932	24.2	(22.6-25.9)	34.4	(32.4-36.4)	2.7	(2.1-3.4)	27.6	(25.8-29.5)	11.1	(9.9-12.4)
Arthritis	Yes	2,002	21.9	(19.7-24.2)	28.9	(26.3-31.8)	2.5	(1.8-3.4)	31.5	(28.6-34.5)	15.2	(13.1-17.5)
	No	3,707	24.6	(22.7-26.6)	35.1	(32.9-37.4)	2.7	(2.0-3.7)	27.7	(25.6-29.8)	9.9	(8.6-11.4)
Frequent Mental Distress	Yes	599	21.6	(16.8-27.3)	24.7	(19.9-30.2)	3.5	(1.2-9.5)*	30.5	(24.7-36.9)	19.7	(16.0-24.0)
	No	5,063	24.0	(22.5-25.7)	34.5	(32.6-36.5)	2.4	(2.0-3.0)	28.7	(26.9-30.5)	10.3	(9.1-11.6)
Serious Psychological Distress	Yes	232	21.5	(15.6-28.9)	21.8	(14.8-31.1)	1.1	(0.3-4.1)*	30.5	(22.0-40.6)	25.0	(18.6-32.6)
	No	5,415	23.7	(22.1-25.3)	34.1	(32.3-36.0)	2.6	(2.0-3.3)	28.9	(27.2-30.7)	10.8	(9.6-12.0)
Receiving Treatment for Mental Health Condition	Yes	669	15.9	(12.7-19.7)	28.7	(23.8-34.2)	1.8	(0.9-3.5)*	32.8	(27.9-38.0)	20.9	(17.2-25.1)
	No	5,049	24.6	(23.0-26.3)	34.1	(32.2-36.0)	2.7	(2.1-3.5)	28.4	(26.6-30.3)	10.2	(9.0-11.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,853	69.3	(67.5-71.1)	23.9	(22.2-25.7)	1.4	(1.1-1.9)	3.9	(3.1-4.7)	1.4	(1.0-2.1)
Age (years)	18-24	167	67.4	(58.1-75.5)	23.6	(16.6-32.5)	2.7	(1.0-7.0)*	6.0	(3.1-11.4)*	0.3	(0.0-1.8)*
	25-34	476	61.1	(55.4-66.5)	31.7	(26.6-37.3)	0.9	(0.4-2.3)*	4.8	(2.7-8.3)	1.5	(0.5-4.1)*
	35-54	1,756	72.2	(69.4-74.7)	22.5	(20.1-25.2)	1.1	(0.7-1.9)	2.4	(1.7-3.4)	1.8	(0.9-3.2)*
	55+	2,426	72.3	(70.0-74.4)	20.5	(18.5-22.6)	1.6	(1.1-2.2)	4.3	(3.2-5.6)	1.4	(0.9-2.2)
Sex	Men	1,751	64.4	(61.4-67.4)	26.8	(24.1-29.7)	1.5	(0.9-2.3)	5.2	(3.9-6.9)	2.0	(1.3-3.3)
	Women	3,102	73.9	(71.8-76.0)	21.2	(19.2-23.2)	1.4	(0.9-2.1)	2.6	(2.0-3.3)	0.9	(0.5-1.5)
Race/Ethnicity	Hispanic or Latino	55	72.7	(54.6-85.5)	20.5	(10.1-37.1)*	4.3	(0.6-24.0)*	1.1	(0.3-4.5)*	1.5	(0.2-10.4)*
	Not Hispanic or Latino:	4,318	70.4	(68.5-72.2)	23.2	(21.5-25.1)	1.4	(1.0-1.9)	3.8	(3.0-4.7)	1.2	(0.8-1.8)
	White											
	Black or African American	311	65.2	(57.5-72.2)	27.0	(20.7-34.3)	1.5	(0.5-3.9)*	5.1	(2.7-9.6)*	1.3	(0.3-5.7)*
	All Other Races	119	59.3	(47.1-70.5)	37.1	(26.1-49.7)	1.2	(0.4-3.4)*	1.5	(0.4-5.0)*	0.8	(0.2-4.0)*
Education	<High School	347	60.6	(52.4-68.2)	29.3	(22.5-37.1)	2.0	(0.8-5.2)*	5.8	(3.5-9.5)	2.3	(1.0-5.4)*
	High School Graduate (or GED)	1,767	64.5	(61.1-67.7)	27.6	(24.6-30.8)	1.8	(1.1-2.8)	4.7	(3.5-6.5)	1.4	(0.8-2.5)
	Some College	1,258	70.2	(66.6-73.5)	23.0	(20.0-26.3)	1.3	(0.6-2.6)*	3.9	(2.6-5.8)	1.6	(0.7-3.8)*
	College Graduate	1,476	75.4	(72.3-78.2)	20.0	(17.3-22.8)	1.1	(0.7-1.7)	2.4	(1.5-4.0)	1.2	(0.5-2.4)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	767	62.8	(57.4-68.0)	28.1	(23.2-33.5)	2.5	(1.4-4.6)*	5.0	(3.1-7.8)	1.6	(0.7-3.4)*
	\$20,000-\$34,999	980	65.4	(61.2-69.4)	25.9	(22.2-29.9)	0.9	(0.5-1.7)*	5.8	(3.9-8.3)	2.0	(1.0-4.1)*
	\$35,000-\$49,999	668	67.4	(62.4-72.1)	25.8	(21.4-30.8)	1.6	(0.8-3.4)*	3.9	(2.2-6.8)	1.2	(0.5-2.7)*
	\$50,000-\$74,999	801	70.4	(66.0-74.4)	22.3	(18.7-26.4)	1.5	(0.7-3.1)*	3.3	(1.7-6.3)*	2.4	(1.3-4.6)*
	\$75,000 and more	1,060	75.2	(71.7-78.5)	21.8	(18.7-25.3)	0.6	(0.3-1.1)*	2.0	(1.2-3.5)	0.3	(0.1-1.2)*
Employment Status	Employed	2,536	70.1	(67.6-72.5)	22.9	(20.7-25.2)	1.2	(0.8-1.8)	4.1	(3.0-5.5)	1.7	(1.0-2.9)
	Out of work	172	65.8	(53.4-76.3)	28.0	(18.7-39.6)	2.0	(0.4-9.0)*	4.2	(1.7-10.3)*	0.0	(0.0-0.0)
	Homemaker	461	69.2	(63.4-74.5)	24.3	(19.2-30.1)	1.4	(0.5-3.5)*	3.0	(1.6-5.4)*	2.2	(0.9-5.2)*
	Student	92	77.1	(66.1-85.3)	19.4	(12.2-29.6)	2.4	(0.3-15.4)*	1.0	(0.3-3.5)*	0.0	(0.0-0.0)
	Retired	1,273	65.9	(62.5-69.2)	27.2	(24.1-30.6)	1.3	(0.8-2.1)	4.2	(3.0-5.9)	1.4	(0.7-2.5)*
	Unable to work	302	66.3	(59.0-72.9)	25.1	(19.0-32.4)	3.0	(1.4-6.3)*	4.6	(2.7-7.8)	1.0	(0.4-2.8)*
Veteran Status	Yes	614	66.1	(61.2-70.7)	28.2	(23.7-33.1)	0.7	(0.3-1.4)*	3.9	(2.4-6.2)	1.1	(0.5-2.6)*
	No	4,239	69.7	(67.7-71.6)	23.3	(21.5-25.2)	1.6	(1.1-2.2)	3.9	(3.1-4.9)	1.5	(1.0-2.3)
Diabetes	Yes	613	73.0	(68.1-77.3)	21.8	(17.7-26.5)	1.1	(0.6-2.1)*	3.1	(1.7-5.7)*	1.0	(0.4-2.5)*
	No	4,236	68.9	(66.9-70.8)	24.1	(22.3-26.0)	1.5	(1.1-2.0)	4.0	(3.2-4.9)	1.5	(1.0-2.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	618	66.5 (61.7-71.1)	24.5 (20.3-29.3)	1.3 (0.7-2.5)*	5.6 (3.6-8.6)	2.1 (0.8-5.1)*				
	No	4,176	69.7 (67.7-71.6)	23.9 (22.1-25.8)	1.4 (1.0-2.0)	3.6 (2.9-4.6)	1.4 (0.9-2.1)				
Asthma	Yes	640	74.8 (69.4-79.6)	19.6 (15.5-24.5)	0.4 (0.2-0.9)*	3.5 (1.9-6.5)*	1.5 (0.7-3.3)*				
	No	4,198	68.4 (66.5-70.3)	24.5 (22.7-26.4)	1.6 (1.2-2.2)	3.9 (3.2-4.9)	1.5 (1.0-2.3)				
Arthritis	Yes	2,014	69.5 (66.7-72.1)	23.0 (20.5-25.6)	1.4 (0.9-2.1)	4.2 (3.1-5.6)	1.9 (1.2-3.1)				
	No	2,814	69.3 (66.9-71.6)	24.3 (22.1-26.6)	1.4 (0.9-2.1)	3.7 (2.8-4.9)	1.2 (0.7-2.1)				
Frequent Mental Distress	Yes	576	67.3 (62.1-72.2)	24.6 (20.1-29.7)	1.4 (0.7-2.8)*	5.1 (3.4-7.5)	1.7 (0.8-3.7)*				
	No	4,196	69.7 (67.7-71.7)	23.7 (21.9-25.6)	1.5 (1.1-2.0)	3.7 (2.9-4.7)	1.4 (0.9-2.1)				
Serious Psychological Distress	Yes	212	57.6 (48.3-66.4)	30.7 (22.6-40.2)	2.9 (1.2-6.6)*	5.9 (3.2-10.6)*	2.9 (1.2-7.0)*				
	No	4,536	70.1 (68.1-71.9)	23.6 (21.8-25.4)	1.3 (1.0-1.9)	3.7 (2.9-4.6)	1.4 (0.9-2.1)				
Receiving Treatment for Mental Health Condition	Yes	726	77.1 (72.9-80.8)	17.4 (14.0-21.5)	1.1 (0.6-2.3)*	3.7 (2.4-5.7)	0.7 (0.3-1.7)*				
	No	4,114	68.3 (66.3-70.3)	24.8 (22.9-26.7)	1.5 (1.1-2.0)	3.9 (3.1-4.9)	1.6 (1.0-2.3)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,866	20.1	(18.6-21.7)	38.0	(36.2-39.9)	3.3	(2.7-4.0)	27.7	(26.1-29.4)	10.8	(9.7-12.0)
Age (years)	18-24	168	20.5	(13.7-29.6)	47.9	(38.9-57.0)	0.3	(0.0-1.8)*	20.9	(14.7-28.8)	10.4	(6.1-17.4)
	25-34	476	14.5	(10.7-19.3)	40.2	(34.9-45.7)	3.7	(2.1-6.4)	31.9	(27.1-37.2)	9.8	(7.1-13.2)
	35-54	1,765	17.4	(15.2-19.8)	37.6	(34.8-40.5)	3.8	(2.8-5.2)	29.2	(26.6-31.8)	12.0	(10.3-14.1)
	55+	2,431	27.1	(25.0-29.3)	34.1	(31.8-36.5)	3.3	(2.5-4.3)	25.0	(22.9-27.2)	10.5	(9.2-12.1)
Sex	Men	1,768	21.0	(18.6-23.6)	42.7	(39.7-45.8)	3.3	(2.4-4.5)	24.3	(21.8-27.0)	8.8	(7.2-10.6)
	Women	3,098	19.4	(17.6-21.3)	33.6	(31.4-35.8)	3.3	(2.5-4.2)	31.0	(28.9-33.2)	12.8	(11.3-14.4)
Race/ Ethnicity	Hispanic or Latino	58	15.2	(8.6-25.7)	38.8	(24.0-56.0)	0.6	(0.1-3.3)*	35.7	(20.6-54.3)	9.7	(2.6-30.1)*
	Not Hispanic or Latino:	4,330	17.9	(16.4-19.4)	39.0	(37.0-41.0)	3.4	(2.7-4.2)	29.1	(27.3-31.0)	10.7	(9.5-11.9)
	White											
	Black or African American	311	19.8	(14.9-25.8)	33.5	(26.6-41.3)	1.8	(0.7-5.0)*	27.2	(21.2-34.3)	17.6	(12.5-24.3)
	All Other Races	118	22.8	(13.9-34.9)	41.4	(30.0-53.8)	7.3	(2.8-17.9)*	20.6	(12.2-32.7)	7.9	(4.4-13.9)
Education	<High School	344	29.6	(23.2-37.1)	40.0	(32.4-48.1)	2.3	(1.0-5.0)*	17.8	(12.8-24.2)	10.3	(6.8-15.4)
	High School Graduate (or GED)	1,796	28.1	(25.2-31.3)	33.6	(30.5-36.8)	3.1	(2.2-4.4)	23.9	(21.3-26.7)	11.2	(9.3-13.5)
	Some College	1,243	13.6	(11.4-16.1)	41.7	(38.0-45.4)	3.5	(2.4-5.0)	30.3	(27.1-33.8)	11.0	(9.1-13.1)
	College Graduate	1,479	13.0	(10.9-15.5)	40.5	(37.4-43.8)	3.5	(2.4-5.1)	32.2	(29.1-35.4)	10.7	(8.9-12.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	773	23.0	(18.6-28.0)	30.8	(25.9-36.1)	2.8	(1.9-4.2)	27.6	(23.1-32.5)	15.8	(12.3-20.1)
	\$20,000-\$34,999	983	22.9	(19.8-26.2)	35.1	(31.1-39.4)	2.5	(1.4-4.3)	27.8	(24.2-31.8)	11.7	(9.3-14.5)
	\$35,000-\$49,999	672	22.4	(18.4-26.9)	35.0	(30.3-39.9)	2.9	(1.9-4.6)	28.5	(24.2-33.3)	11.2	(8.4-14.7)
	\$50,000-\$74,999	800	16.1	(13.1-19.6)	42.1	(37.8-46.5)	3.6	(2.1-6.1)	29.0	(25.3-33.0)	9.2	(7.1-11.9)
	\$75,000 and more	1,066	16.4	(13.7-19.6)	43.0	(39.3-46.8)	3.8	(2.5-5.6)	28.7	(25.5-32.1)	8.1	(6.4-10.2)
Employment Status	Employed	2,539	18.4	(16.4-20.5)	40.9	(38.4-43.4)	3.5	(2.6-4.6)	27.3	(25.2-29.6)	10.0	(8.6-11.6)
	Out of work	171	22.8	(13.7-35.5)	32.1	(23.0-42.8)	3.7	(1.2-10.5)*	26.5	(18.9-35.7)	14.9	(8.9-23.9)
	Homemaker	459	22.4	(17.9-27.7)	28.9	(23.9-34.4)	4.9	(2.9-8.0)	32.5	(27.4-38.1)	11.4	(8.3-15.4)
	Student	91	13.3	(6.7-24.6)*	44.9	(32.8-57.7)	0.7	(0.2-2.6)*	28.4	(18.9-40.2)	12.7	(6.3-23.8)*
	Retired	1,288	24.1	(21.5-27.0)	37.2	(33.8-40.7)	2.9	(2.0-4.3)	26.8	(23.7-30.2)	9.0	(7.2-11.1)
	Unable to work	302	24.1	(17.9-31.7)	21.8	(15.9-29.1)	1.6	(0.8-3.1)*	30.5	(23.7-38.1)	21.9	(16.6-28.4)
Veteran Status	Yes	630	19.8	(16.5-23.7)	36.4	(31.5-41.5)	3.0	(1.6-5.3)*	30.4	(25.5-35.8)	10.4	(7.4-14.6)
	No	4,236	20.2	(18.5-21.9)	38.2	(36.2-40.3)	3.3	(2.7-4.1)	27.4	(25.6-29.2)	10.9	(9.7-12.2)
Diabetes	Yes	622	22.5	(18.8-26.7)	31.5	(26.8-36.6)	3.7	(2.0-6.5)	30.1	(25.3-35.5)	12.2	(9.3-15.9)
	No	4,238	19.9	(18.3-21.6)	38.7	(36.7-40.7)	3.2	(2.6-4.0)	27.5	(25.8-29.3)	10.7	(9.5-11.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Ohio, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	616	22.0 (18.5-25.9)	37.0 (31.8-42.4)	2.8 (1.6-4.8)	25.2 (20.9-30.0)	13.1 (10.2-16.7)				
	No	4,188	19.9 (18.3-21.7)	38.2 (36.2-40.2)	3.3 (2.6-4.1)	28.0 (26.2-29.8)	10.6 (9.4-11.9)				
Asthma	Yes	637	16.2 (12.2-21.1)	34.7 (29.6-40.2)	2.6 (1.4-4.8)*	34.5 (29.5-39.9)	12.0 (9.4-15.1)				
	No	4,214	20.7 (19.1-22.4)	38.6 (36.6-40.6)	3.4 (2.7-4.2)	26.6 (24.9-28.4)	10.7 (9.5-12.0)				
Arthritis	Yes	2,021	20.2 (18.1-22.4)	35.3 (32.5-38.2)	2.6 (1.9-3.7)	30.0 (27.3-32.8)	11.9 (10.3-13.8)				
	No	2,820	20.0 (17.9-22.1)	39.2 (36.8-41.7)	3.6 (2.8-4.6)	26.8 (24.8-29.0)	10.4 (9.0-12.0)				
Frequent Mental Distress	Yes	582	16.5 (13.0-20.8)	28.0 (23.1-33.5)	3.0 (1.6-5.4)*	31.7 (26.9-37.0)	20.7 (16.6-25.7)				
	No	4,202	20.9 (19.2-22.6)	39.1 (37.1-41.1)	3.4 (2.7-4.2)	27.2 (25.4-29.0)	9.5 (8.5-10.7)				
Serious Psychological Distress	Yes	216	13.9 (9.2-20.3)	21.2 (14.1-30.5)	0.9 (0.4-2.4)*	33.3 (24.9-43.0)	30.8 (22.9-40.0)				
	No	4,560	20.5 (18.9-22.2)	39.0 (37.1-41.0)	3.3 (2.7-4.1)	27.4 (25.7-29.2)	9.7 (8.7-10.9)				
Receiving Treatment for Mental Health Condition	Yes	721	14.0 (11.3-17.3)	32.4 (28.0-37.2)	4.1 (2.4-6.8)	32.5 (28.1-37.1)	17.1 (14.0-20.7)				
	No	4,132	21.0 (19.4-22.8)	38.8 (36.7-40.9)	3.2 (2.5-3.9)	27.1 (25.3-28.9)	10.0 (8.8-11.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,487	70.8	(69.4-72.3)	24.1	(22.8-25.5)	0.8	(0.5-1.2)	3.0	(2.5-3.6)	1.2	(0.9-1.6)
Age (years)	18-24	268	60.1	(53.4-66.6)	33.3	(27.2-40.0)	1.7	(0.6-4.6)*	4.3	(2.4-7.6)	0.6	(0.1-2.5)*
	25-34	750	67.2	(63.2-71.0)	28.9	(25.2-32.8)	0.4	(0.1-1.2)*	2.7	(1.7-4.4)	0.8	(0.3-2.0)*
	35-54	2,268	74.8	(72.6-76.8)	20.8	(19.0-22.8)	0.6	(0.3-1.2)*	2.7	(2.0-3.6)	1.1	(0.7-1.9)
	55+	3,181	72.2	(70.4-73.9)	21.8	(20.2-23.4)	0.9	(0.6-1.3)	3.2	(2.5-4.0)	2.0	(1.5-2.6)
Sex	Men	2,292	63.5	(61.0-65.9)	30.2	(27.9-32.7)	1.1	(0.6-1.8)	3.8	(3.0-5.0)	1.4	(0.9-2.0)
	Women	4,195	77.6	(76.1-79.2)	18.4	(17.0-19.9)	0.6	(0.4-0.9)	2.3	(1.8-2.9)	1.1	(0.8-1.5)
Race/ Ethnicity	Hispanic or Latino	275	69.6	(62.9-75.5)	21.9	(16.8-27.9)	1.8	(0.6-5.6)*	5.4	(3.1-9.0)	1.4	(0.6-3.0)*
	Not Hispanic or Latino:	5,000	71.3	(69.6-72.9)	24.6	(23.0-26.2)	0.8	(0.6-1.2)	2.5	(2.0-3.0)	0.8	(0.6-1.2)
	White											
	Black or African American	368	67.4	(61.7-72.7)	25.5	(20.6-31.2)	0.2	(0.1-0.4)*	4.0	(2.2-7.1)	2.9	(1.5-5.6)*
	All Other Races	827	70.2	(66.0-74.1)	24.3	(20.6-28.3)	0.6	(0.2-1.6)*	3.3	(2.0-5.6)	1.7	(0.9-3.1)*
Education	<High School	756	61.3	(56.5-65.9)	29.9	(25.6-34.5)	1.0	(0.5-2.0)*	5.2	(3.7-7.3)	2.7	(1.9-3.8)
	High School Graduate (or GED)	2,089	66.9	(64.1-69.6)	27.3	(24.7-30.0)	1.0	(0.6-1.9)*	3.3	(2.4-4.6)	1.5	(1.0-2.4)
	Some College	1,849	72.3	(69.6-74.9)	23.0	(20.5-25.6)	1.1	(0.6-2.0)*	2.8	(2.0-4.0)	0.8	(0.4-1.7)*
	College Graduate	1,781	76.1	(73.6-78.4)	20.7	(18.5-23.2)	0.4	(0.3-0.7)	2.1	(1.4-3.1)	0.7	(0.3-1.3)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,328	65.4 (61.8-68.8)	27.2 (23.9-30.7)	1.0 (0.5-2.3)*	4.6 (3.3-6.3)	1.8 (1.2-2.6)				
	\$20,000-\$34,999	1,374	73.0 (69.8-76.0)	20.9 (18.2-23.8)	1.4 (0.7-2.7)*	3.5 (2.5-5.0)	1.2 (0.8-2.0)				
	\$35,000-\$49,999	975	72.8 (69.1-76.1)	23.4 (20.2-26.9)	0.5 (0.2-1.4)*	2.3 (1.4-3.8)	1.1 (0.4-2.5)*				
	\$50,000-\$74,999	927	72.1 (68.5-75.4)	24.3 (21.1-27.9)	0.3 (0.1-0.9)*	2.8 (1.6-4.7)	0.5 (0.2-1.5)*				
	\$75,000 and more	1,144	72.2 (68.8-75.4)	23.9 (20.8-27.2)	0.6 (0.2-1.6)*	2.8 (1.6-4.9)	0.5 (0.2-1.7)*				
Employment Status	Employed	3,239	72.1 (70.1-74.0)	23.7 (21.9-25.6)	1.0 (0.6-1.6)	2.5 (1.9-3.3)	0.7 (0.5-1.2)				
	Out of work	223	67.1 (58.8-74.5)	27.3 (20.1-35.8)	0.1 (0.0-0.6)*	4.3 (2.0-9.2)*	1.2 (0.5-2.7)*				
	Homemaker	617	74.3 (69.8-78.3)	21.1 (17.4-25.3)	0.7 (0.2-2.3)*	2.2 (1.1-4.2)*	1.8 (0.8-4.0)*				
	Student	144	72.6 (63.3-80.3)	23.6 (16.5-32.5)	0.0 (0.0-0.0)	2.9 (1.2-7.2)*	0.9 (0.1-5.9)*				
	Retired	1,685	66.1 (63.3-68.8)	27.1 (24.4-29.8)	0.8 (0.5-1.3)	4.0 (3.0-5.3)	2.0 (1.3-3.1)				
	Unable to work	572	68.2 (63.4-72.7)	23.4 (19.3-28.1)	0.9 (0.3-2.2)*	5.0 (3.3-7.5)	2.5 (1.4-4.4)				
Veteran Status	Yes	930	69.6 (66.2-72.9)	23.5 (20.5-26.7)	1.1 (0.6-2.2)*	3.8 (2.7-5.5)	1.9 (1.1-3.4)				
	No	5,553	71.0 (69.4-72.6)	24.2 (22.7-25.8)	0.7 (0.5-1.1)	2.9 (2.4-3.6)	1.1 (0.8-1.5)				
Diabetes	Yes	822	69.7 (65.7-73.4)	23.9 (20.5-27.8)	0.9 (0.5-1.9)*	3.4 (2.2-5.2)	2.1 (1.1-3.9)*				
	No	5,664	70.9 (69.3-72.4)	24.1 (22.6-25.6)	0.9 (0.6-1.2)	3.0 (2.5-3.7)	1.2 (0.9-1.5)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	893	69.4	(65.6-73.0)	24.7	(21.3-28.3)	0.8	(0.4-1.6)*	3.3	(2.2-4.9)	1.8	(0.9-3.5)*
	No	5,529	71.1	(69.5-72.6)	24.0	(22.5-25.5)	0.8	(0.5-1.2)	3.0	(2.5-3.7)	1.1	(0.8-1.5)
Asthma	Yes	863	74.5	(70.3-78.3)	20.1	(16.7-24.0)	0.7	(0.3-1.9)*	2.7	(1.7-4.4)	1.9	(1.0-3.7)*
	No	5,610	70.1	(68.5-71.6)	24.7	(23.2-26.2)	0.9	(0.6-1.3)	3.1	(2.6-3.8)	1.2	(0.9-1.5)
Arthritis	Yes	2,544	72.3	(70.0-74.4)	21.9	(19.9-24.1)	0.7	(0.4-1.1)	3.1	(2.5-4.0)	2.0	(1.4-2.9)
	No	3,918	70.2	(68.3-72.0)	25.0	(23.3-26.8)	0.9	(0.5-1.4)	3.0	(2.4-3.8)	0.9	(0.6-1.4)
Frequent Mental Distress	Yes	805	69.6	(65.4-73.6)	24.4	(20.6-28.6)	0.5	(0.2-1.1)*	3.7	(2.4-5.7)	1.8	(1.0-3.2)*
	No	5,570	71.1	(69.5-72.6)	23.9	(22.5-25.4)	0.9	(0.6-1.3)	3.0	(2.5-3.6)	1.1	(0.8-1.5)
Serious Psychological Distress	Yes	333	65.7	(58.6-72.2)	26.8	(20.8-33.9)	1.0	(0.3-3.2)*	3.9	(2.3-6.6)	2.5	(1.4-4.4)
	No	6,004	71.3	(69.7-72.7)	23.8	(22.4-25.3)	0.9	(0.6-1.3)	3.0	(2.5-3.7)	1.0	(0.8-1.4)
Receiving Treatment for Mental Health Condition	Yes	908	77.0	(73.2-80.5)	19.6	(16.3-23.5)	0.2	(0.1-0.4)*	2.1	(1.3-3.3)	1.1	(0.6-2.0)*
	No	5,564	70.1	(68.5-71.7)	24.6	(23.2-26.2)	0.9	(0.6-1.3)	3.1	(2.6-3.8)	1.2	(0.9-1.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,558	27.3	(25.9-28.7)	34.7	(33.2-36.3)	1.2	(0.9-1.6)	25.1	(23.8-26.5)	11.7	(10.7-12.7)
Age (years)	18-24	270	23.8	(18.7-29.8)	46.0	(39.1-53.1)	2.2	(0.9-5.2)*	22.0	(17.0-28.1)	5.9	(3.5-9.8)
	25-34	759	23.8	(20.6-27.5)	36.2	(32.4-40.2)	1.2	(0.6-2.7)*	25.2	(21.9-28.7)	13.5	(10.9-16.6)
	35-54	2,281	25.5	(23.5-27.6)	32.6	(30.5-34.8)	1.0	(0.6-1.6)	27.1	(25.1-29.2)	13.8	(12.3-15.5)
	55+	3,227	32.9	(31.0-34.8)	32.5	(30.7-34.4)	0.9	(0.6-1.3)	23.8	(22.2-25.6)	9.9	(8.8-11.1)
Sex	Men	2,319	30.1	(27.9-32.4)	38.2	(35.8-40.7)	1.0	(0.6-1.7)	21.5	(19.5-23.5)	9.1	(7.8-10.7)
	Women	4,239	24.7	(23.1-26.3)	31.4	(29.6-33.2)	1.3	(0.9-1.9)	28.6	(27.0-30.4)	14.0	(12.8-15.4)
Race/ Ethnicity	Hispanic or Latino	278	41.6	(35.0-48.5)	26.2	(20.0-33.7)	2.4	(0.9-6.4)*	23.1	(17.7-29.6)	6.6	(4.2-10.2)
	Not Hispanic or Latino:	5,036	24.8	(23.4-26.3)	37.1	(35.3-38.8)	1.2	(0.9-1.7)	25.4	(23.9-26.9)	11.5	(10.5-12.7)
	White											
	Black or African American	376	35.1	(29.4-41.3)	27.8	(22.4-33.9)	0.4	(0.1-1.2)*	24.6	(19.6-30.4)	12.1	(8.8-16.5)
	All Other Races	850	28.5	(24.6-32.8)	30.4	(26.4-34.6)	1.1	(0.5-2.3)*	25.3	(21.8-29.2)	14.7	(12.1-17.7)
Education	<High School	769	34.4	(30.2-38.8)	34.6	(29.9-39.7)	1.7	(0.8-3.5)*	18.8	(15.3-22.9)	10.5	(8.2-13.5)
	High School Graduate (or GED)	2,122	32.7	(30.1-35.4)	34.5	(31.8-37.4)	1.3	(0.8-2.3)	21.0	(18.9-23.2)	10.5	(8.9-12.3)
	Some College	1,866	25.6	(23.1-28.3)	31.8	(29.1-34.6)	1.1	(0.5-2.0)*	27.9	(25.3-30.5)	13.7	(11.9-15.7)
	College Graduate	1,790	21.7	(19.5-24.2)	37.9	(35.2-40.6)	0.9	(0.5-1.6)	28.0	(25.6-30.5)	11.5	(9.9-13.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,368	28.0 (25.0-31.3)	30.4 (27.0-34.0)	2.2 (1.2-4.1)*	24.9 (21.9-28.2)	14.5 (12.1-17.2)				
	\$20,000-\$34,999	1,388	28.7 (25.8-31.9)	33.0 (29.7-36.4)	1.1 (0.6-2.2)*	25.8 (22.9-29.0)	11.4 (9.5-13.5)				
	\$35,000-\$49,999	985	29.1 (25.8-32.8)	35.0 (31.4-38.8)	0.7 (0.3-1.6)*	24.7 (21.6-28.1)	10.4 (8.3-13.0)				
	\$50,000-\$74,999	926	26.7 (23.5-30.2)	34.7 (31.2-38.4)	1.5 (0.7-3.3)*	24.1 (21.1-27.3)	13.0 (10.6-16.0)				
	\$75,000 and more	1,138	24.0 (20.9-27.3)	38.1 (34.7-41.5)	0.4 (0.2-1.0)*	26.7 (23.9-29.6)	10.9 (9.0-13.0)				
Employment Status	Employed	3,238	28.3 (26.4-30.3)	34.3 (32.3-36.4)	1.3 (0.8-1.9)	24.5 (22.8-26.3)	11.6 (10.3-13.0)				
	Out of work	222	25.4 (19.1-33.0)	36.8 (28.5-46.0)	1.0 (0.3-3.5)*	22.1 (15.8-29.9)	14.7 (9.3-22.4)				
	Homemaker	629	24.7 (20.9-28.9)	35.4 (31.0-40.2)	1.3 (0.5-3.4)*	27.0 (23.1-31.3)	11.6 (9.1-14.7)				
	Student	148	22.3 (14.8-32.3)	39.8 (30.5-50.0)	1.5 (0.3-6.2)*	26.2 (18.8-35.2)	10.2 (5.6-17.8)				
	Retired	1,729	30.3 (27.8-32.9)	35.0 (32.3-37.8)	0.9 (0.6-1.4)	25.7 (23.1-28.5)	8.1 (6.8-9.7)				
	Unable to work	585	20.9 (17.3-24.9)	30.1 (25.4-35.4)	1.8 (0.9-3.7)*	27.3 (22.9-32.3)	19.8 (16.1-24.2)				
Veteran Status	Yes	946	28.2 (25.0-31.7)	34.1 (30.6-37.7)	1.5 (0.8-2.7)*	24.4 (21.1-28.2)	11.8 (9.3-14.8)				
	No	5,609	27.1 (25.6-28.7)	34.8 (33.2-36.5)	1.2 (0.9-1.7)	25.2 (23.8-26.7)	11.6 (10.6-12.7)				
Diabetes	Yes	821	25.3 (22.0-29.0)	35.9 (31.9-40.2)	1.1 (0.6-2.1)*	25.7 (22.1-29.7)	11.9 (9.5-14.9)				
	No	5,736	27.5 (26.1-29.0)	34.5 (32.9-36.2)	1.3 (0.9-1.7)	25.0 (23.7-26.4)	11.6 (10.6-12.7)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oklahoma, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	907	28.1	(24.8-31.7)	32.8	(29.0-36.8)	1.3	(0.6-2.5)*	24.0	(20.6-27.6)	13.9	(11.4-16.8)
	No	5,582	27.3	(25.8-28.8)	34.9	(33.2-36.5)	1.2	(0.8-1.6)	25.2	(23.8-26.6)	11.5	(10.5-12.6)
Asthma	Yes	873	25.6	(21.9-29.6)	31.9	(28.0-36.1)	0.5	(0.2-1.1)*	26.1	(22.6-29.8)	16.0	(13.1-19.3)
	No	5,670	27.5	(26.1-29.0)	35.2	(33.6-36.9)	1.2	(0.9-1.7)	25.0	(23.6-26.5)	11.0	(10.1-12.1)
Arthritis	Yes	2,568	26.0	(24.0-28.1)	31.6	(29.3-33.9)	0.8	(0.5-1.3)	27.8	(25.7-30.1)	13.8	(12.1-15.6)
	No	3,966	27.8	(26.1-29.7)	36.0	(34.1-37.9)	1.3	(0.9-1.9)	24.1	(22.5-25.7)	10.8	(9.7-12.0)
Frequent Mental Distress	Yes	817	20.8	(17.4-24.6)	30.6	(26.4-35.2)	1.5	(0.7-3.0)*	28.2	(24.3-32.5)	18.9	(15.9-22.3)
	No	5,629	28.4	(26.9-29.9)	34.9	(33.3-36.5)	1.2	(0.9-1.7)	24.7	(23.4-26.1)	10.8	(9.8-11.9)
Serious Psychological Distress	Yes	341	17.8	(13.1-23.6)	28.9	(22.6-36.1)	1.8	(0.6-4.8)*	31.4	(24.9-38.6)	20.2	(15.3-26.2)
	No	6,064	28.0	(26.5-29.4)	34.9	(33.3-36.5)	1.1	(0.8-1.6)	24.8	(23.5-26.2)	11.2	(10.3-12.2)
Receiving Treatment for Mental Health Condition	Yes	910	19.8	(16.9-23.0)	27.3	(23.7-31.2)	1.2	(0.6-2.3)*	33.7	(29.8-37.9)	18.0	(15.1-21.4)
	No	5,632	28.3	(26.8-29.8)	35.7	(34.1-37.4)	1.2	(0.8-1.6)	24.0	(22.7-25.4)	10.8	(9.8-11.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		1,833	72.8	(70.1-75.4)	22.1	(19.7-24.8)	1.4	(0.8-2.5)	2.5	(1.7-3.8)	1.1	(0.6-1.9)
Age (years)	18-24	61	67.5	(54.0-78.7)	27.2	(17.1-40.3)	2.5	(0.4-16.0)*	2.7	(0.9-8.3)*	0.0	(0.0-0.0)
	25-34	180	65.5	(57.5-72.7)	28.7	(21.7-36.7)	1.3	(0.4-3.6)*	3.9	(1.4-10.0)*	0.7	(0.1-5.0)*
	35-54	625	75.5	(71.5-79.1)	20.1	(16.8-23.9)	1.8	(0.8-3.8)*	1.1	(0.5-2.5)*	1.6	(0.7-3.4)*
	55+	967	76.4	(73.4-79.2)	18.3	(15.8-21.1)	0.8	(0.4-1.5)*	3.1	(2.0-4.8)	1.4	(0.7-2.9)*
Sex	Men	698	68.9	(64.4-73.1)	24.8	(20.9-29.2)	1.5	(0.8-3.0)*	3.3	(1.8-5.9)	1.4	(0.6-3.1)*
	Women	1,135	76.4	(73.2-79.4)	19.6	(16.9-22.7)	1.3	(0.5-3.3)*	1.8	(1.1-2.9)	0.8	(0.4-1.6)*
Race/ Ethnicity	Hispanic or Latino	58	64.4	(49.5-77.0)	32.0	(19.8-47.1)	3.4	(0.9-12.7)*	0.1	(0.1-0.2)	0.1	(0.0-0.1)
	Not Hispanic or Latino:	1,664	73.5	(70.6-76.1)	21.8	(19.3-24.5)	1.3	(0.7-2.5)*	2.5	(1.6-3.8)	0.9	(0.5-1.7)*
	White											
	Black or African American	8	--	--	--	--	--	--	--	--	--	--
	All Other Races	90	79.0	(66.6-87.6)	14.4	(7.7-25.6)*	0.8	(0.1-5.1)*	3.9	(1.3-11.1)*	1.9	(0.5-7.8)*
Education	<High School	118	62.9	(51.3-73.2)	27.1	(18.0-38.6)	6.4	(3.1-12.8)*	2.8	(1.0-7.5)*	0.7	(0.1-5.2)*
	High School Graduate (or GED)	456	68.2	(62.0-73.7)	23.9	(19.0-29.7)	1.2	(0.4-3.4)*	4.7	(2.4-9.1)*	2.0	(0.9-4.7)*
	Some College	569	69.1	(63.9-73.8)	26.9	(22.4-32.0)	1.8	(0.6-5.0)*	1.6	(0.8-3.2)*	0.6	(0.2-1.6)*
	College Graduate	688	82.2	(78.3-85.4)	14.7	(11.6-18.4)	0.4	(0.2-0.7)*	1.7	(0.8-3.5)*	1.0	(0.4-2.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	198	69.6	(60.4-77.5)	24.0	(16.7-33.2)	1.6	(0.5-4.5)*	3.5	(1.5-8.0)*	1.4	(0.4-4.8)*
	\$20,000-\$34,999	421	67.5	(60.8-73.5)	25.9	(20.3-32.3)	1.4	(0.6-3.1)*	4.3	(2.1-8.7)*	0.9	(0.3-3.1)*
	\$35,000-\$49,999	291	74.1	(67.3-79.9)	22.5	(17.0-29.1)	1.1	(0.3-4.5)*	1.1	(0.3-4.2)*	1.3	(0.2-7.0)*
	\$50,000-\$74,999	328	73.1	(67.0-78.4)	23.2	(18.2-29.2)	1.3	(0.3-5.2)*	1.9	(0.8-4.8)*	0.5	(0.1-2.1)*
	\$75,000 and more	395	81.4	(75.9-85.9)	15.7	(11.5-21.1)	1.0	(0.3-3.2)*	1.6	(0.6-4.7)*	0.3	(0.0-1.8)*
Employment Status	Employed	934	74.5	(70.7-77.9)	20.7	(17.6-24.2)	1.7	(0.8-3.6)*	2.1	(1.0-4.3)*	1.0	(0.4-2.4)*
	Out of work	58	72.1	(55.8-84.0)	23.0	(11.9-39.7)*	3.3	(0.8-12.7)*	1.1	(0.2-5.2)*	0.5	(0.1-4.0)*
	Homemaker	153	70.8	(61.1-78.9)	22.9	(15.8-32.1)	0.0	(0.0-0.0)	4.1	(1.5-10.7)*	2.2	(0.7-6.9)*
	Student	35	--	--	--	--	--	--	--	--	--	--
	Retired	558	71.4	(67.0-75.4)	22.7	(19.0-26.9)	1.2	(0.6-2.3)*	4.0	(2.4-6.8)	0.6	(0.1-2.4)*
	Unable to work	94	56.1	(43.9-67.6)	35.0	(24.0-47.9)	0.0	(0.0-0.0)	5.3	(1.8-14.5)*	3.6	(1.1-10.7)*
Veteran Status	Yes	267	72.8	(66.3-78.5)	21.3	(16.0-27.8)	0.8	(0.2-2.8)*	2.8	(1.3-6.0)*	2.2	(0.6-7.8)*
	No	1,566	72.7	(69.7-75.6)	22.3	(19.6-25.2)	1.5	(0.8-2.8)*	2.5	(1.6-4.0)	0.9	(0.5-1.6)
Diabetes	Yes	166	78.4	(71.0-84.3)	16.0	(11.1-22.6)	0.8	(0.2-3.4)*	4.1	(1.7-9.9)*	0.7	(0.1-3.7)*
	No	1,666	72.3	(69.4-75.0)	22.6	(20.0-25.4)	1.5	(0.8-2.6)	2.5	(1.6-3.8)	1.2	(0.7-2.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	168	70.6	(62.0-78.0)	22.5	(15.9-31.0)	2.2	(0.8-5.8)*	4.6	(1.8-11.4)*	0.0	(0.0-0.0)
	No	1,646	73.0	(70.1-75.7)	22.1	(19.5-24.9)	1.4	(0.7-2.6)*	2.4	(1.5-3.7)	1.2	(0.7-2.1)
Asthma	Yes	295	77.0	(69.9-82.8)	15.4	(10.6-21.7)	1.3	(0.5-3.3)*	4.4	(1.7-10.7)*	2.0	(0.7-5.7)*
	No	1,531	71.8	(68.8-74.6)	23.6	(21.0-26.5)	1.5	(0.8-2.8)*	2.1	(1.4-3.2)	1.0	(0.5-1.8)*
Arthritis	Yes	651	72.3	(67.9-76.3)	21.0	(17.4-25.2)	1.1	(0.5-2.4)*	2.0	(1.1-3.5)	3.6	(1.9-6.8)*
	No	1,174	73.0	(69.6-76.2)	22.3	(19.4-25.6)	1.6	(0.8-3.1)*	2.7	(1.6-4.5)	0.3	(0.1-0.7)*
Frequent Mental Distress	Yes	173	79.0	(70.0-85.8)	17.6	(11.3-26.3)	0.9	(0.2-4.1)*	2.0	(0.6-6.1)*	0.5	(0.1-2.4)*
	No	1,643	72.3	(69.4-75.0)	22.7	(20.1-25.5)	1.3	(0.7-2.4)*	2.6	(1.7-4.1)	1.0	(0.6-1.9)*
Serious Psychological Distress	Yes	64	56.5	(43.7-68.6)	29.4	(18.5-43.3)	3.7	(1.0-12.1)*	7.1	(2.4-19.0)*	3.3	(0.7-14.5)*
	No	1,740	73.2	(70.4-75.9)	21.9	(19.4-24.6)	1.3	(0.7-2.4)*	2.5	(1.6-3.8)	1.1	(0.6-1.9)
Receiving Treatment for Mental Health Condition	Yes	292	79.5	(73.1-84.6)	17.1	(12.3-23.3)	0.1	(0.1-0.1)	2.4	(0.9-6.1)*	1.0	(0.3-3.3)*
	No	1,536	72.1	(69.1-74.9)	22.6	(20.0-25.5)	1.6	(0.9-2.8)	2.6	(1.6-4.0)	1.1	(0.6-2.0)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		1,822	17.7	(15.5-20.1)	32.8	(30.0-35.6)	2.1	(1.4-3.2)	31.4	(28.7-34.2)	16.1	(14.1-18.2)
Age (years)	18-24	62	12.1	(5.9-23.3)*	39.3	(27.3-52.9)	2.7	(0.4-16.3)*	36.3	(24.4-50.2)	9.6	(4.5-19.5)*
	25-34	179	13.1	(8.2-20.2)	30.9	(23.8-39.0)	1.9	(0.7-4.9)*	37.7	(30.2-45.9)	16.4	(11.3-23.2)
	35-54	622	17.4	(14.0-21.3)	32.9	(28.9-37.2)	1.6	(0.8-3.2)*	29.9	(26.2-34.0)	18.2	(15.0-21.8)
	55+	959	23.3	(20.4-26.5)	31.1	(28.0-34.5)	2.7	(1.7-4.3)	27.6	(24.6-30.7)	15.2	(13.0-17.8)
Sex	Men	697	18.8	(15.4-22.8)	35.1	(30.7-39.7)	2.1	(1.2-3.7)	30.9	(26.8-35.4)	13.0	(10.2-16.5)
	Women	1,125	16.6	(14.2-19.5)	30.6	(27.4-34.0)	2.1	(1.2-3.8)	31.7	(28.5-35.1)	18.9	(16.4-21.7)
Race/ Ethnicity	Hispanic or Latino	58	14.7	(7.5-27.0)*	40.2	(25.8-56.6)	0.2	(0.2-0.3)	34.0	(21.0-50.0)	10.8	(5.2-21.1)*
	Not Hispanic or Latino:	1,654	15.8	(13.8-18.2)	33.3	(30.4-36.3)	2.1	(1.4-3.3)	33.0	(30.1-36.0)	15.8	(13.8-18.0)
	White											
	Black or African American	8	--	--	--	--	--	--	--	--	--	--
	All Other Races	90	19.7	(10.6-33.9)*	30.5	(20.3-43.0)	4.1	(1.3-12.8)*	20.7	(12.8-31.7)	24.9	(14.7-39.0)
Education	<High School	117	26.9	(17.2-39.5)	33.6	(23.7-45.2)	0.5	(0.1-3.7)*	22.4	(14.2-33.4)	16.6	(10.3-25.6)
	High School Graduate (or GED)	452	20.4	(16.1-25.5)	33.1	(27.4-39.2)	1.3	(0.5-3.0)*	29.6	(24.1-35.9)	15.6	(11.5-20.8)
	Some College	571	15.6	(12.3-19.5)	33.3	(28.5-38.5)	3.3	(1.8-6.2)*	30.8	(26.2-35.8)	17.0	(13.7-21.0)
	College Graduate	680	13.4	(10.4-17.1)	31.9	(27.9-36.3)	2.1	(1.1-3.9)*	36.4	(32.1-40.9)	16.2	(13.2-19.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	200	18.7 (11.8-28.3)	27.2 (19.9-36.0)	1.0 (0.3-3.2)*	31.8 (24.0-40.8)	21.3 (15.3-28.9)				
	\$20,000-\$34,999	418	17.3 (12.9-22.8)	32.9 (26.8-39.6)	2.0 (0.9-4.1)*	28.5 (22.7-35.0)	19.3 (14.4-25.4)				
	\$35,000-\$49,999	289	19.1 (14.4-25.0)	36.1 (29.4-43.5)	0.5 (0.1-2.0)*	33.7 (27.4-40.6)	10.6 (7.4-14.9)				
	\$50,000-\$74,999	326	18.3 (13.6-24.2)	33.3 (27.6-39.6)	2.2 (0.9-4.9)*	28.3 (22.9-34.4)	17.9 (13.6-23.2)				
	\$75,000 and more	394	13.0 (9.6-17.5)	35.3 (29.9-41.1)	2.5 (1.2-4.9)*	36.6 (31.1-42.5)	12.5 (9.4-16.5)				
Employment Status	Employed	929	17.7 (14.7-21.1)	32.2 (28.6-36.0)	2.5 (1.5-4.3)	31.6 (28.0-35.4)	16.0 (13.3-19.0)				
	Out of work	57	10.4 (2.7-32.3)*	35.8 (21.6-53.0)	4.1 (1.2-12.7)*	34.8 (21.0-51.7)	14.9 (6.4-31.2)*				
	Homemaker	151	20.8 (13.8-30.2)	33.5 (25.2-43.0)	1.3 (0.4-3.8)*	30.3 (22.5-39.4)	14.1 (9.2-21.0)				
	Student	35	--	--	--	--	--	--			
	Retired	555	21.2 (17.8-25.0)	32.1 (27.9-36.6)	1.7 (0.9-3.2)*	31.9 (27.8-36.4)	13.0 (10.3-16.3)				
	Unable to work	94	15.3 (7.8-27.8)*	27.0 (17.2-39.6)	0.5 (0.1-4.5)*	24.8 (15.7-36.9)	32.4 (22.4-44.5)				
Veteran Status	Yes	264	19.6 (14.8-25.5)	36.4 (29.8-43.6)	2.2 (0.8-5.6)*	28.4 (22.5-35.1)	13.4 (9.4-18.7)				
	No	1,557	17.4 (15.1-20.1)	32.1 (29.2-35.2)	2.1 (1.4-3.3)	31.9 (29.0-34.9)	16.5 (14.4-18.9)				
Diabetes	Yes	162	26.2 (18.9-35.1)	29.9 (22.3-38.9)	2.1 (0.6-7.6)*	25.8 (18.7-34.3)	16.0 (10.2-24.1)				
	No	1,659	17.0 (14.8-19.5)	33.1 (30.2-36.0)	2.1 (1.3-3.2)	31.8 (29.0-34.7)	16.1 (14.1-18.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Oregon, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	165	20.0	(14.2-27.4)	28.6	(21.7-36.8)	2.5	(1.1-5.4)*	24.9	(18.2-33.0)	24.0	(16.7-33.3)
	No	1,637	17.5	(15.2-20.1)	33.1	(30.3-36.1)	2.1	(1.4-3.3)	31.8	(29.1-34.8)	15.4	(13.4-17.6)
Asthma	Yes	291	18.3	(12.8-25.5)	26.4	(20.3-33.5)	2.1	(0.9-4.7)*	31.5	(25.2-38.5)	21.7	(16.0-28.8)
	No	1,524	17.5	(15.2-20.0)	34.0	(31.0-37.1)	2.1	(1.3-3.3)	31.4	(28.6-34.5)	15.0	(13.0-17.2)
Arthritis	Yes	653	15.6	(12.9-18.9)	34.5	(30.2-39.1)	1.9	(0.9-3.7)*	29.3	(25.4-33.5)	18.7	(15.4-22.6)
	No	1,160	18.5	(15.7-21.6)	32.1	(28.8-35.6)	2.2	(1.3-3.6)	32.2	(28.9-35.7)	15.0	(12.7-17.6)
Frequent Mental Distress	Yes	173	10.0	(5.3-18.0)*	27.9	(20.4-36.8)	0.5	(0.1-2.0)*	31.2	(22.6-41.4)	30.5	(22.7-39.5)
	No	1,630	18.7	(16.4-21.3)	33.1	(30.2-36.1)	2.3	(1.5-3.5)	31.6	(28.8-34.5)	14.3	(12.4-16.5)
Serious Psychological Distress	Yes	64	9.9	(5.0-18.9)*	28.8	(18.1-42.5)	0.0	(0.0-0.0)	19.8	(11.2-32.7)	41.4	(28.9-55.2)
	No	1,728	17.8	(15.6-20.3)	33.0	(30.2-36.0)	2.2	(1.4-3.3)	31.8	(29.1-34.7)	15.2	(13.2-17.4)
Receiving Treatment for Mental Health Condition	Yes	291	14.7	(10.2-20.8)	34.2	(27.9-41.2)	0.5	(0.2-1.6)*	28.5	(22.7-35.2)	22.0	(16.9-28.2)
	No	1,526	18.1	(15.7-20.7)	32.5	(29.5-35.6)	2.4	(1.6-3.6)	31.9	(29.0-34.9)	15.2	(13.1-17.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,717	29.4	(26.9-32.1)	57.2	(54.0-60.2)	8.9	(7.0-11.3)	3.7	(2.9-4.8)	0.8	(0.5-1.2)
Age (years)	18-24	252	31.4	(25.1-38.4)	50.1	(43.1-57.1)	12.6	(8.5-18.3)	4.9	(2.7-8.6)	1.0	(0.4-2.6)*
	25-34	322	22.7	(17.5-28.9)	62.4	(55.7-68.7)	10.3	(6.9-15.3)	2.6	(1.3-5.0)*	1.9	(1.0-3.8)*
	35-54	1,199	32.4	(28.9-36.0)	55.3	(51.3-59.2)	8.2	(6.1-11.0)	3.5	(2.5-5.1)	0.5	(0.3-1.0)*
	55+	1,929	27.6	(24.8-30.6)	62.3	(58.9-65.5)	5.9	(4.4-7.8)	3.9	(2.9-5.1)	0.4	(0.2-0.7)
Sex	Men	1,303	28.3	(24.9-32.0)	58.4	(54.2-62.4)	8.9	(6.6-11.9)	3.5	(2.4-4.9)	1.0	(0.6-1.8)
	Women	2,414	30.8	(27.9-34.0)	55.8	(52.3-59.3)	8.8	(6.8-11.4)	3.9	(2.9-5.1)	0.6	(0.4-1.0)
Race/ Ethnicity	Hispanic or Latino	3,691	29.0	(27.0-31.0)	55.8	(53.5-58.0)	10.2	(8.9-11.6)	3.9	(3.2-4.6)	1.3	(0.9-1.8)
	Not Hispanic or Latino:	16	--	--	--	--	--	--	--	--	--	--
	White											
	Black or African American	4	--	--	--	--	--	--	--	--	--	--
	All Other Races	6	--	--	--	--	--	--	--	--	--	--
Education	<High School	1,027	21.5	(18.0-25.4)	64.7	(60.0-69.0)	7.9	(5.6-10.9)	5.2	(3.7-7.2)	0.8	(0.4-1.8)*
	High School Graduate (or GED)	931	20.9	(17.2-25.2)	54.8	(49.7-59.9)	17.0	(13.1-21.7)	6.0	(4.4-8.3)	1.2	(0.8-2.0)
	Some College	752	35.0	(30.1-40.2)	44.9	(39.8-50.1)	16.2	(12.7-20.5)	3.1	(2.0-4.7)	0.8	(0.4-1.6)*
	College Graduate	1,000	31.6	(27.7-35.8)	50.2	(45.6-54.8)	15.3	(12.0-19.3)	2.1	(1.2-3.6)	0.8	(0.4-1.6)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,913	24.9 (21.9-28.2)	62.6 (58.9-66.1)	7.4 (5.5-9.8)	4.3 (3.2-5.7)	0.8 (0.4-1.4)				
	\$20,000-\$34,999	724	29.5 (25.3-33.9)	56.1 (51.1-61.0)	9.5 (6.7-13.3)	4.1 (2.6-6.5)	0.8 (0.3-2.1)*				
	\$35,000-\$49,999	271	34.9 (28.2-42.4)	52.9 (45.2-60.5)	9.7 (5.7-16.0)	1.3 (0.4-4.1)*	1.1 (0.4-3.4)*				
	\$50,000-\$74,999	162	25.6 (18.9-33.7)	59.6 (50.1-68.4)	11.3 (6.2-19.8)	3.2 (0.7-13.1)*	0.3 (0.0-1.9)*				
	\$75,000 and more	141	33.3 (25.2-42.5)	47.1 (37.0-57.5)	17.9 (10.2-29.4)	0.8 (0.1-5.7)*	0.9 (0.2-4.0)*				
Employment Status	Employed	1,384	30.0 (26.7-33.6)	57.1 (53.1-61.0)	8.7 (6.5-11.6)	3.2 (2.2-4.6)	0.9 (0.5-1.7)				
	Out of work	148	30.8 (21.9-41.3)	57.3 (47.1-66.9)	6.7 (3.3-13.1)*	4.0 (1.8-8.6)*	1.3 (0.4-4.6)*				
	Homemaker	830	23.8 (20.0-28.0)	64.6 (59.9-69.1)	7.1 (4.9-10.2)	3.9 (2.7-5.6)	0.6 (0.3-1.4)*				
	Student	163	37.9 (29.0-47.7)	40.3 (32.0-49.1)	17.8 (11.6-26.4)	3.0 (1.4-6.4)*	1.0 (0.3-3.1)*				
	Retired	861	29.7 (25.9-33.8)	61.4 (57.0-65.6)	4.8 (3.3-6.9)	3.6 (2.4-5.5)	0.5 (0.2-1.0)*				
	Unable to work	327	26.2 (20.4-32.9)	56.0 (48.8-63.1)	10.0 (6.3-15.4)	7.3 (4.3-12.1)	0.5 (0.2-1.6)*				
Veteran Status	Yes	267	35.2 (27.4-43.9)	51.8 (43.6-60.0)	6.7 (4.0-11.1)	4.4 (2.0-9.1)*	1.8 (0.7-5.0)*				
	No	3,450	29.3 (26.7-32.0)	57.5 (54.3-60.6)	9.0 (7.0-11.4)	3.6 (2.8-4.6)	0.7 (0.5-1.1)				
Diabetes	Yes	666	21.4 (17.4-26.0)	64.0 (58.7-68.9)	8.5 (6.1-11.7)	5.7 (3.8-8.5)	0.4 (0.2-0.9)*				
	No	3,043	29.4 (26.7-32.3)	53.0 (49.6-56.3)	12.9 (10.4-15.9)	3.8 (2.9-4.9)	0.9 (0.6-1.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	505	29.8 (24.5-35.8)	56.8 (50.5-62.8)	6.8 (4.4-10.3)	5.7 (3.4-9.3)	0.9 (0.4-2.3)*				
	No	3,192	29.4 (26.8-32.2)	57.1 (53.9-60.3)	9.2 (7.2-11.8)	3.5 (2.7-4.5)	0.8 (0.5-1.2)				
Asthma	Yes	568	24.7 (20.0-30.2)	53.4 (47.5-59.3)	15.7 (11.9-20.3)	4.7 (3.1-6.9)	1.5 (0.7-3.0)*				
	No	3,142	28.4 (25.7-31.2)	53.2 (49.8-56.6)	13.5 (10.9-16.6)	4.1 (3.2-5.3)	0.8 (0.5-1.2)				
Arthritis	Yes	1,210	28.7 (25.2-32.5)	60.2 (56.1-64.2)	6.7 (4.8-9.3)	3.8 (2.7-5.3)	0.6 (0.3-1.3)*				
	No	2,489	29.5 (26.6-32.5)	56.3 (52.8-59.6)	9.8 (7.6-12.5)	3.7 (2.8-4.8)	0.9 (0.5-1.4)				
Frequent Mental Distress	Yes	461	28.2 (22.6-34.5)	53.6 (47.2-59.8)	12.3 (8.6-17.2)	5.2 (3.3-8.1)	0.8 (0.3-2.1)*				
	No	3,170	30.0 (27.3-32.8)	57.4 (54.1-60.5)	8.6 (6.6-11.0)	3.3 (2.5-4.4)	0.8 (0.5-1.2)				
Serious Psychological Distress	Yes	268	19.0 (12.9-27.1)	70.0 (60.9-77.7)	5.8 (2.1-14.8)*	3.1 (0.9-10.3)*	2.1 (1.1-4.1)*				
	No	3,374	28.3 (25.6-31.0)	53.6 (50.3-56.9)	13.3 (10.7-16.4)	3.9 (3.0-5.0)	0.9 (0.6-1.4)				
Receiving Treatment for Mental Health Condition	Yes	446	31.3 (25.9-37.3)	51.0 (44.7-57.3)	11.3 (7.6-16.4)	5.3 (3.4-8.1)	1.1 (0.4-3.3)*				
	No	3,269	28.9 (26.2-31.7)	57.7 (54.5-60.9)	9.1 (7.1-11.6)	3.5 (2.7-4.6)	0.8 (0.5-1.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,684	4.4	(3.7-5.1)	30.9	(28.1-33.8)	17.3	(14.1-21.1)	40.8	(37.4-44.4)	6.6	(5.4-8.0)
Age (years)	18-24	254	5.3	(3.3-8.6)	30.3	(24.3-37.0)	22.1	(16.1-29.7)	33.5	(26.7-40.9)	8.8	(5.5-13.7)
	25-34	328	3.0	(1.8-5.1)	29.7	(24.1-35.9)	14.1	(9.9-19.6)	46.4	(39.8-53.1)	6.9	(4.4-10.7)
	35-54	1,190	3.6	(2.7-4.7)	26.0	(22.8-29.6)	17.5	(13.8-22.0)	45.2	(40.9-49.6)	7.6	(5.9-9.7)
	55+	1,899	5.8	(4.9-6.9)	38.8	(35.3-42.3)	15.6	(12.5-19.3)	35.7	(32.2-39.5)	4.0	(3.1-5.3)
Sex	Men	1,285	4.2	(3.3-5.3)	33.6	(29.8-37.5)	17.3	(13.7-21.7)	39.1	(34.8-43.6)	5.9	(4.4-7.9)
	Women	2,399	4.5	(3.7-5.5)	28.2	(25.3-31.3)	17.3	(13.9-21.3)	42.7	(38.9-46.6)	7.3	(5.8-9.2)
Race/ Ethnicity	Hispanic or Latino	3,658	5.5	(4.6-6.4)	35.6	(33.6-37.7)	15.3	(13.7-17.1)	37.7	(35.6-39.8)	5.9	(5.0-7.0)
	Not Hispanic or Latino:	16	--	--	--	--	--	--	--	--	--	--
	White											
	Black or African American	4	--	--	--	--	--	--	--	--	--	--
	All Other Races	6	--	--	--	--	--	--	--	--	--	--
Education	<High School	997	7.1	(5.6-9.0)	55.0	(50.1-59.7)	10.8	(8.0-14.5)	24.6	(20.6-29.1)	2.5	(1.5-4.2)
	High School Graduate (or GED)	929	4.0	(3.0-5.4)	33.5	(29.1-38.3)	18.2	(13.9-23.4)	36.8	(31.9-42.0)	7.5	(5.4-10.2)
	Some College	753	5.3	(3.9-7.1)	27.0	(22.6-31.8)	22.3	(17.3-28.4)	37.9	(32.6-43.5)	7.5	(5.3-10.6)
	College Graduate	999	2.7	(1.8-4.0)	20.0	(16.8-23.6)	18.8	(14.5-24.0)	51.6	(46.6-56.6)	6.9	(5.2-9.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,881	5.3 (4.4-6.4)	34.8 (31.2-38.5)	16.9 (13.2-21.4)	36.8 (32.9-40.9)	6.2 (4.7-8.1)				
	\$20,000-\$34,999	722	4.7 (3.3-6.8)	25.8 (21.8-30.2)	20.4 (15.6-26.3)	41.1 (35.9-46.6)	8.0 (5.7-11.0)				
	\$35,000-\$49,999	273	4.7 (2.4-8.8)*	22.5 (16.9-29.3)	20.0 (13.9-27.9)	46.1 (38.5-53.9)	6.7 (3.7-11.9)				
	\$50,000-\$74,999	161	3.0 (1.3-6.8)*	23.5 (16.5-32.3)	13.8 (8.2-22.2)	54.9 (45.3-64.1)	4.9 (2.5-9.4)*				
	\$75,000 and more	142	2.7 (0.7-10.1)*	13.5 (8.2-21.3)	21.2 (13.4-31.8)	50.7 (40.8-60.6)	11.9 (7.5-18.5)				
Employment Status	Employed	1,375	2.0 (1.3-3.2)	31.8 (28.3-35.4)	15.4 (12.0-19.4)	43.8 (39.6-48.2)	7.0 (5.4-9.0)				
	Out of work	149	2.4 (0.7-8.4)*	43.0 (33.9-52.6)	12.1 (6.5-21.6)*	31.8 (23.5-41.4)	10.7 (5.3-20.3)*				
	Homemaker	817	3.3 (2.2-4.8)	37.9 (33.3-42.7)	17.1 (13.3-21.7)	35.7 (31.1-40.7)	6.0 (4.1-8.6)				
	Student	164	1.0 (0.0-30.2)*	38.7 (30.5-47.7)	29.7 (20.8-40.6)	29.4 (21.0-39.4)	1.1 (0.0-50.4)*				
	Retired	848	5.1 (4.0-6.6)	39.1 (34.6-43.9)	16.5 (12.9-21.0)	36.0 (31.3-41.0)	3.2 (2.0-5.1)				
	Unable to work	327	1.8 (0.7-4.4)*	37.8 (31.6-44.4)	10.6 (6.6-16.7)	41.3 (34.4-48.6)	8.5 (5.6-12.8)				
Veteran Status	Yes	271	2.8 (1.4-5.6)*	36.1 (29.1-43.7)	10.2 (6.0-16.7)	44.8 (36.7-53.2)	6.1 (2.9-12.2)*				
	No	3,413	2.4 (1.8-3.1)	37.0 (34.0-40.1)	13.5 (10.8-16.8)	41.1 (37.6-44.5)	6.0 (4.9-7.5)				
Diabetes	Yes	646	2.3 (1.5-3.5)	41.0 (35.8-46.4)	16.4 (12.6-21.0)	36.0 (30.5-41.8)	4.4 (2.6-7.3)				
	No	3,029	2.5 (1.8-3.3)	36.7 (33.7-39.8)	11.7 (9.2-14.8)	42.5 (39.0-46.0)	6.7 (5.4-8.2)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Puerto Rico, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	499	4.8	(3.4-6.6)	29.2	(24.1-34.9)	16.9	(12.3-22.7)	41.7	(35.2-48.5)	7.5	(4.7-11.7)
	No	3,166	4.3	(3.6-5.1)	31.0	(28.1-34.0)	17.5	(14.2-21.3)	40.8	(37.3-44.5)	6.4	(5.2-7.9)
Asthma	Yes	566	3.6	(2.3-5.5)	29.1	(24.2-34.4)	13.7	(9.9-18.6)	45.4	(39.3-51.6)	8.3	(5.6-12.0)
	No	3,112	4.5	(3.8-5.4)	31.3	(28.4-34.4)	18.1	(14.8-22.1)	39.8	(36.3-43.5)	6.2	(5.0-7.7)
Arthritis	Yes	1,208	4.9	(3.9-6.1)	34.9	(31.0-39.0)	15.9	(12.3-20.2)	37.8	(33.6-42.3)	6.5	(4.8-8.6)
	No	2,459	4.2	(3.5-5.2)	29.8	(26.8-33.0)	18.1	(14.6-22.3)	41.3	(37.5-45.2)	6.5	(5.2-8.1)
Frequent Mental Distress	Yes	466	2.2	(1.0-4.8)*	24.9	(20.4-29.9)	11.1	(7.0-17.0)	46.4	(40.3-52.6)	15.5	(11.9-19.9)
	No	3,132	2.5	(1.9-3.3)	38.4	(35.3-41.6)	13.2	(10.5-16.5)	40.7	(37.2-44.2)	5.3	(4.1-6.7)
Serious Psychological Distress	Yes	271	6.0	(3.5-10.0)	24.6	(18.4-32.0)	12.4	(8.2-18.4)	45.8	(37.7-54.1)	11.2	(7.3-16.9)
	No	3,339	4.2	(3.6-5.0)	30.6	(27.8-33.5)	17.8	(14.6-21.5)	40.9	(37.4-44.6)	6.5	(5.2-8.0)
Receiving Treatment for Mental Health Condition	Yes	444	5.5	(3.8-7.8)	31.6	(26.1-37.7)	13.5	(9.5-18.9)	39.2	(33.1-45.6)	10.2	(6.8-15.0)
	No	3,239	4.3	(3.6-5.1)	31.0	(28.1-34.1)	17.7	(14.4-21.6)	40.9	(37.4-44.6)	6.0	(4.9-7.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,820	61.3	(59.1-63.5)	29.9	(27.7-32.1)	3.0	(2.3-4.0)	4.1	(3.2-5.2)	1.7	(1.1-2.5)
Age (years)	18-24	106	49.9	(39.2-60.7)	42.9	(32.6-53.8)	3.0	(1.0-8.7)*	2.2	(0.7-6.7)*	2.0	(0.4-8.9)*
	25-34	317	51.2	(44.3-58.0)	34.0	(27.6-41.1)	4.9	(2.6-8.9)*	7.9	(4.6-13.3)	2.0	(0.6-6.3)*
	35-54	1,446	63.0	(60.0-65.9)	28.9	(26.1-31.8)	2.6	(1.6-4.0)	4.1	(3.0-5.6)	1.5	(0.9-2.6)
	55+	1,918	69.0	(66.6-71.4)	23.8	(21.7-26.1)	2.9	(2.1-4.0)	2.9	(2.1-3.8)	1.4	(0.9-2.2)
Sex	Men	1,319	55.5	(51.7-59.2)	33.7	(30.2-37.4)	3.4	(2.3-5.1)	5.4	(3.9-7.5)	1.9	(1.1-3.5)*
	Women	2,501	66.8	(64.1-69.4)	26.2	(23.7-28.8)	2.7	(1.9-3.8)	2.9	(2.2-4.0)	1.4	(0.8-2.5)*
Race/ Ethnicity	Hispanic or Latino	254	59.4	(51.6-66.8)	29.2	(22.4-37.0)	0.9	(0.3-2.7)*	6.8	(3.7-12.2)*	3.7	(1.5-9.1)*
	Not Hispanic or Latino:	3,331	62.6	(60.2-65.0)	29.8	(27.5-32.2)	3.1	(2.4-4.1)	3.4	(2.6-4.4)	1.0	(0.6-1.8)
	White											
	Black or African American	92	43.4	(31.5-56.1)	35.6	(23.9-49.4)	9.7	(2.4-32.0)*	5.0	(1.3-16.7)*	6.2	(2.2-16.2)*
	All Other Races	117	67.3	(53.7-78.5)	23.7	(14.4-36.4)	1.7	(0.2-11.1)*	6.6	(2.7-15.2)*	0.7	(0.2-2.8)*
Education	<High School	335	53.2	(46.2-60.1)	33.3	(27.0-40.4)	4.6	(2.1-10.1)*	7.4	(4.3-12.6)	1.4	(0.6-3.0)*
	High School Graduate (or GED)	1,044	59.2	(55.0-63.2)	29.1	(25.3-33.2)	3.9	(2.5-6.0)	5.0	(3.2-7.8)	2.8	(1.5-5.1)*
	Some College	941	61.6	(56.5-66.4)	32.8	(28.1-37.8)	1.8	(1.0-3.1)	3.2	(2.0-5.1)	0.7	(0.3-1.4)*
	College Graduate	1,498	65.3	(61.9-68.6)	27.2	(24.1-30.5)	2.8	(1.8-4.5)	3.0	(2.1-4.4)	1.6	(0.7-3.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	531	61.8	(55.6-67.6)	28.6	(23.2-34.8)	2.6	(1.3-5.3)*	4.6	(2.7-7.8)	2.4	(1.3-4.1)
	\$20,000-\$34,999	611	59.1	(53.5-64.6)	30.3	(25.3-35.8)	2.6	(1.6-4.3)	5.9	(3.8-9.0)	2.1	(0.6-7.0)*
	\$35,000-\$49,999	510	65.7	(59.6-71.4)	24.4	(19.2-30.5)	2.9	(1.6-5.4)*	4.9	(2.5-9.4)*	2.0	(1.0-4.1)*
	\$50,000-\$74,999	605	61.9	(56.6-66.8)	30.6	(25.9-35.9)	4.3	(2.4-7.3)	2.7	(1.6-4.7)	0.5	(0.2-1.6)*
	\$75,000 and more	1,083	63.6	(59.5-67.5)	30.0	(26.3-34.1)	2.1	(1.1-4.1)*	2.7	(1.7-4.3)	1.6	(0.6-4.2)*
Employment Status	Employed	2,088	60.4	(57.4-63.4)	30.5	(27.6-33.4)	3.5	(2.4-5.0)	4.1	(2.9-5.7)	1.5	(0.8-2.7)*
	Out of work	164	56.9	(47.3-66.0)	34.0	(25.1-44.1)	2.6	(1.0-7.1)*	6.1	(2.5-14.3)*	0.4	(0.1-1.6)*
	Homemaker	243	66.2	(58.8-72.8)	25.1	(19.1-32.1)	2.8	(1.3-6.0)*	5.8	(2.8-11.7)*	0.2	(0.0-1.4)*
	Student	49	68.2	(51.1-81.5)	26.5	(14.7-43.0)	1.4	(0.2-7.6)*	0.0	(0.0-0.0)	3.9	(0.5-23.4)*
	Retired	1,028	65.5	(62.0-68.8)	26.0	(23.0-29.4)	2.8	(1.8-4.3)	3.5	(2.3-5.2)	2.2	(1.3-3.7)
	Unable to work	239	51.7	(44.5-58.9)	37.8	(30.4-45.8)	1.5	(0.5-4.8)*	6.5	(3.7-11.3)	2.4	(1.1-5.6)*
Veteran Status	Yes	525	63.8	(58.8-68.6)	28.0	(23.5-32.9)	3.0	(1.8-5.1)	3.6	(2.2-5.8)	1.6	(0.7-3.5)*
	No	3,294	60.9	(58.5-63.4)	30.2	(27.8-32.6)	3.0	(2.3-4.1)	4.2	(3.2-5.4)	1.7	(1.0-2.7)
Diabetes	Yes	385	60.6	(54.4-66.4)	29.3	(24.0-35.2)	2.8	(1.4-5.3)*	5.0	(2.9-8.5)	2.4	(1.1-4.9)*
	No	3,433	61.3	(58.9-63.7)	29.9	(27.7-32.3)	3.1	(2.3-4.1)	4.0	(3.1-5.2)	1.6	(1.0-2.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	409	65.9	(60.5-71.0)	28.6	(23.8-34.0)	2.0	(0.9-4.1)*	1.7	(0.9-3.5)*	1.7	(0.7-4.0)*
	No	3,378	61.0	(58.6-63.4)	29.8	(27.6-32.2)	3.2	(2.4-4.2)	4.3	(3.4-5.5)	1.6	(1.0-2.6)
Asthma	Yes	581	61.3	(55.7-66.6)	29.4	(24.3-35.0)	2.9	(1.4-6.1)*	4.5	(2.5-8.1)*	1.9	(0.5-6.9)*
	No	3,230	61.3	(58.8-63.7)	29.9	(27.6-32.3)	3.1	(2.3-4.1)	4.1	(3.2-5.2)	1.7	(1.1-2.5)
Arthritis	Yes	1,509	66.3	(63.2-69.3)	26.7	(23.9-29.7)	2.4	(1.7-3.4)	3.1	(2.2-4.2)	1.5	(0.9-2.5)
	No	2,298	59.2	(56.3-62.1)	31.3	(28.5-34.2)	3.2	(2.3-4.5)	4.5	(3.4-6.0)	1.7	(1.0-2.9)
Frequent Mental Distress	Yes	410	57.0	(50.0-63.8)	33.2	(26.9-40.3)	3.3	(1.7-6.4)*	5.1	(2.7-9.5)*	1.3	(0.6-2.9)*
	No	3,364	62.0	(59.6-64.3)	29.4	(27.2-31.8)	2.9	(2.1-3.9)	4.0	(3.1-5.1)	1.7	(1.1-2.7)
Serious Psychological Distress	Yes	199	45.6	(35.8-55.9)	36.1	(26.9-46.4)	5.1	(2.1-11.6)*	10.8	(5.9-19.0)*	2.4	(1.2-4.8)*
	No	3,561	62.2	(59.9-64.5)	29.5	(27.3-31.8)	3.0	(2.2-3.9)	3.7	(2.9-4.8)	1.5	(1.0-2.5)
Receiving Treatment for Mental Health Condition	Yes	597	63.6	(58.3-68.7)	27.3	(22.4-32.9)	2.3	(1.1-5.0)*	4.9	(3.0-7.9)	1.9	(0.9-3.7)*
	No	3,211	61.1	(58.6-63.5)	30.2	(27.9-32.6)	3.1	(2.4-4.2)	4.0	(3.0-5.1)	1.6	(1.0-2.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		3,823	24.9	(22.9-26.9)	38.6	(36.5-40.8)	4.4	(3.7-5.3)	22.4	(20.7-24.3)	9.7	(8.6-10.9)
Age (years)	18-24	108	25.0	(16.8-35.6)	41.1	(31.1-51.8)	3.3	(1.2-9.1)*	25.9	(17.9-36.0)	4.7	(2.0-10.5)*
	25-34	318	22.4	(17.3-28.4)	40.8	(34.5-47.4)	4.6	(2.7-7.9)	23.8	(18.9-29.5)	8.4	(5.4-12.8)
	35-54	1,449	21.3	(18.8-23.9)	38.4	(35.6-41.4)	4.6	(3.5-5.9)	23.2	(20.8-25.8)	12.5	(10.7-14.6)
	55+	1,915	31.1	(28.7-33.6)	36.6	(34.1-39.2)	4.6	(3.6-5.9)	19.0	(17.2-21.1)	8.6	(7.4-10.1)
Sex	Men	1,322	27.2	(24.1-30.6)	40.4	(36.9-43.9)	3.6	(2.7-4.8)	20.9	(18.2-23.9)	7.9	(6.2-10.0)
	Women	2,501	22.8	(20.6-25.1)	36.8	(34.2-39.5)	5.1	(4.0-6.4)	24.0	(21.8-26.4)	11.3	(9.9-12.9)
Race/ Ethnicity	Hispanic or Latino	253	38.4	(30.9-46.6)	27.5	(20.8-35.3)	3.3	(1.2-8.3)*	22.7	(15.8-31.4)	8.1	(5.0-13.0)
	Not Hispanic or Latino:	3,333	21.1	(19.2-23.1)	41.0	(38.7-43.4)	4.7	(3.9-5.7)	23.6	(21.7-25.7)	9.5	(8.3-10.8)
	White											
	Black or African American	91	32.0	(20.8-45.8)	24.4	(14.4-38.3)	3.9	(0.9-14.6)*	25.9	(14.9-41.1)	13.8	(7.7-23.5)
	All Other Races	118	31.5	(18.2-48.8)	37.3	(25.3-51.2)	1.7	(0.5-5.9)*	17.2	(10.2-27.7)	12.3	(7.4-19.7)
Education	<High School	337	37.6	(31.0-44.8)	31.0	(24.7-38.2)	2.6	(1.4-4.8)*	20.6	(15.2-27.3)	8.1	(5.4-11.9)
	High School Graduate (or GED)	1,044	28.7	(25.1-32.6)	37.5	(33.5-41.7)	4.3	(3.1-5.9)	19.6	(16.5-23.2)	9.9	(7.7-12.7)
	Some College	945	22.9	(18.7-27.9)	39.1	(34.4-44.0)	4.4	(2.9-6.7)	23.0	(19.4-27.1)	10.6	(8.5-13.1)
	College Graduate	1,493	18.6	(16.2-21.3)	41.2	(38.0-44.5)	4.9	(3.7-6.5)	25.5	(22.7-28.6)	9.7	(8.0-11.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	531	29.5 (24.1-35.6)	30.1 (24.8-36.1)	4.3 (2.5-7.3)	23.0 (17.4-29.8)	13.0 (10.0-16.7)				
	\$20,000-\$34,999	613	27.5 (22.7-32.8)	35.5 (30.3-41.0)	4.5 (2.9-6.9)	22.5 (18.5-27.2)	10.0 (7.0-14.1)				
	\$35,000-\$49,999	509	23.5 (19.0-28.7)	36.1 (30.3-42.4)	4.5 (2.7-7.4)	24.1 (19.2-29.8)	11.7 (8.7-15.6)				
	\$50,000-\$74,999	605	21.9 (18.1-26.2)	45.1 (40.2-50.2)	4.4 (3.0-6.3)	20.8 (17.1-25.1)	7.8 (5.6-10.8)				
	\$75,000 and more	1,082	21.2 (18.0-24.8)	42.2 (38.3-46.1)	3.7 (2.6-5.1)	23.9 (20.9-27.2)	9.2 (7.3-11.4)				
Employment Status	Employed	2,088	25.3 (22.6-28.2)	39.3 (36.4-42.2)	3.8 (3.0-4.8)	21.7 (19.5-24.0)	10.0 (8.5-11.7)				
	Out of work	165	20.9 (13.3-31.2)	40.6 (30.9-51.0)	3.3 (1.2-8.6)*	25.5 (17.6-35.3)	9.8 (6.3-14.8)				
	Homemaker	247	25.4 (19.7-32.1)	40.5 (33.6-47.7)	6.3 (3.6-10.6)	20.5 (15.4-26.6)	7.4 (4.5-11.8)				
	Student	49	12.9 (4.1-34.0)*	33.2 (19.5-50.4)	16.0 (7.7-30.4)*	30.3 (17.4-47.3)	7.6 (2.2-23.5)*				
	Retired	1,026	29.0 (26.1-32.2)	37.8 (34.4-41.4)	3.9 (2.8-5.5)	22.1 (19.2-25.3)	7.1 (5.6-9.0)				
	Unable to work	240	19.4 (14.2-25.9)	34.2 (26.4-43.0)	6.8 (3.8-11.9)	23.6 (16.3-32.8)	16.0 (11.6-21.7)				
Veteran Status	Yes	520	24.7 (21.0-28.9)	34.1 (29.5-39.2)	3.4 (2.1-5.3)	25.1 (20.8-30.0)	12.7 (9.1-17.5)				
	No	3,302	24.8 (22.7-27.1)	39.2 (36.9-41.6)	4.6 (3.8-5.6)	22.1 (20.2-24.1)	9.3 (8.2-10.6)				
Diabetes	Yes	381	28.3 (22.7-34.7)	32.8 (27.0-39.3)	5.4 (3.1-9.1)	21.6 (17.1-27.0)	11.9 (8.6-16.1)				
	No	3,439	24.5 (22.5-26.7)	39.1 (36.8-41.4)	4.4 (3.6-5.3)	22.5 (20.7-24.5)	9.5 (8.3-10.8)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Rhode Island, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illnesst

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	406	24.0	(19.5-29.1)	36.8	(31.4-42.4)	4.5	(2.8-7.3)	22.6	(18.2-27.8)	12.1	(8.7-16.6)
	No	3,385	25.0	(23.0-27.2)	38.9	(36.6-41.2)	4.4	(3.6-5.4)	22.3	(20.5-24.3)	9.3	(8.2-10.6)
Asthma	Yes	582	17.9	(14.2-22.4)	36.7	(31.4-42.4)	5.9	(4.0-8.7)	25.8	(20.9-31.4)	13.6	(10.3-17.8)
	No	3,232	26.0	(23.9-28.3)	39.0	(36.7-41.4)	4.1	(3.3-5.1)	21.9	(20.0-23.8)	8.9	(7.8-10.2)
Arthritis	Yes	1,506	21.8	(19.5-24.3)	37.0	(34.0-40.2)	4.7	(3.6-6.1)	23.4	(20.8-26.3)	13.0	(11.1-15.3)
	No	2,302	26.2	(23.6-28.9)	39.3	(36.5-42.1)	4.2	(3.2-5.4)	22.2	(19.9-24.5)	8.3	(7.0-9.8)
Frequent Mental Distress	Yes	410	17.6	(12.8-23.8)	33.8	(27.4-40.9)	4.7	(2.8-7.9)	26.8	(21.0-33.6)	17.0	(12.8-22.3)
	No	3,367	25.8	(23.8-28.0)	39.3	(37.0-41.6)	4.4	(3.6-5.3)	21.8	(19.9-23.7)	8.8	(7.7-10.0)
Serious Psychological Distress	Yes	200	14.2	(9.8-20.0)	18.3	(12.3-26.3)	9.5	(4.6-18.4)*	33.4	(23.7-44.7)	24.7	(18.1-32.7)
	No	3,564	25.4	(23.4-27.6)	39.6	(37.4-41.9)	4.2	(3.4-5.1)	22.0	(20.2-23.9)	8.8	(7.7-10.0)
Receiving Treatment for Mental Health Condition	Yes	592	20.3	(16.3-25.0)	34.7	(29.6-40.2)	5.7	(3.5-9.2)	24.1	(19.6-29.3)	15.1	(12.0-18.9)
	No	3,219	25.6	(23.5-27.9)	39.2	(36.9-41.6)	4.2	(3.5-5.2)	22.2	(20.3-24.2)	8.7	(7.6-10.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		8,761	68.1	(66.2-69.9)	25.6	(23.9-27.5)	1.1	(0.7-1.8)	3.8	(3.1-4.6)	1.4	(1.0-2.1)
Age (years)	18-24	251	58.5	(49.0-67.5)	33.7	(25.7-42.7)	0.4	(0.1-2.1)*	6.7	(3.7-11.7)	0.7	(0.3-1.5)*
	25-34	632	62.7	(57.3-67.7)	30.6	(25.8-35.8)	1.8	(0.6-5.5)*	2.9	(1.7-4.7)	2.1	(0.8-5.3)*
	35-54	2,804	68.6	(65.8-71.2)	24.9	(22.4-27.6)	1.1	(0.6-2.0)*	3.5	(2.6-4.6)	1.9	(1.1-3.2)
	55+	5,003	73.6	(71.6-75.5)	20.6	(18.8-22.5)	0.9	(0.5-1.5)	3.6	(2.8-4.5)	1.3	(0.9-1.9)
Sex	Men	3,316	62.0	(58.9-65.0)	29.5	(26.7-32.5)	1.4	(0.8-2.7)*	4.9	(3.8-6.4)	2.1	(1.3-3.4)
	Women	5,445	73.7	(71.6-75.8)	22.0	(20.0-24.1)	0.7	(0.4-1.4)*	2.7	(2.1-3.5)	0.9	(0.6-1.2)
Race/ Ethnicity	Hispanic or Latino	136	49.7	(37.4-62.1)	31.6	(20.2-45.7)	8.6	(4.1-17.1)*	3.0	(1.1-7.9)*	7.2	(3.2-15.4)*
	Not Hispanic or Latino:	5,820	68.1	(66.1-70.1)	25.7	(23.8-27.7)	1.1	(0.7-1.6)	3.6	(2.8-4.6)	1.4	(1.0-2.1)
	White											
	Black or African American	2,323	65.8	(61.6-69.7)	25.4	(21.8-29.4)	1.0	(0.3-2.9)*	5.4	(4.3-6.8)	2.5	(1.7-3.6)
	All Other Races	278	66.5	(57.1-74.7)	22.9	(16.0-31.6)	1.5	(0.4-4.8)*	7.4	(3.7-14.3)*	1.8	(0.7-4.5)*
Education	<High School	1,101	58.2	(52.5-63.6)	33.2	(27.9-39.0)	1.0	(0.4-2.2)*	6.3	(4.4-8.8)	1.4	(0.7-2.9)*
	High School Graduate (or GED)	2,699	64.9	(61.3-68.3)	27.4	(24.3-30.8)	0.6	(0.3-1.3)*	4.7	(3.5-6.4)	2.3	(1.4-3.7)
	Some College	2,255	67.9	(64.4-71.3)	26.5	(23.3-30.0)	1.1	(0.5-2.4)*	2.7	(1.7-4.3)	1.8	(0.9-3.6)*
	College Graduate	2,693	73.7	(70.4-76.7)	21.5	(18.7-24.6)	2.0	(0.8-4.7)*	2.6	(1.6-4.2)	0.3	(0.1-0.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,696	63.2	(58.4-67.8)	27.6	(23.5-32.2)	1.0	(0.4-2.4)*	6.6	(4.6-9.4)	1.6	(1.0-2.4)
	\$20,000-\$34,999	1,766	67.5	(63.5-71.3)	25.7	(22.2-29.7)	1.4	(0.6-3.4)*	3.2	(2.2-4.5)	2.2	(1.0-4.4)*
	\$35,000-\$49,999	1,109	71.0	(66.1-75.4)	23.8	(19.6-28.6)	1.0	(0.4-2.2)*	2.3	(1.0-5.1)*	2.0	(0.9-4.3)*
	\$50,000-\$74,999	1,178	69.4	(64.9-73.5)	26.6	(22.6-31.0)	0.9	(0.2-3.3)*	2.2	(1.3-3.6)	0.9	(0.4-2.2)*
	\$75,000 and more	1,726	69.3	(65.4-73.0)	26.5	(22.9-30.4)	1.3	(0.5-3.7)*	2.5	(1.5-4.0)	0.4	(0.2-1.0)*
Employment Status	Employed	4,055	66.9	(64.3-69.4)	26.3	(23.9-28.8)	1.3	(0.7-2.5)*	3.4	(2.6-4.6)	2.0	(1.2-3.3)
	Out of work	587	70.7	(64.4-76.3)	24.4	(19.2-30.5)	0.4	(0.2-1.1)*	3.7	(2.2-6.0)	0.8	(0.3-2.5)*
	Homemaker	548	69.6	(62.7-75.6)	25.0	(19.3-31.6)	1.6	(0.3-8.4)*	3.2	(1.8-5.9)*	0.6	(0.2-1.9)*
	Student	129	74.6	(62.7-83.7)	19.7	(12.1-30.4)	0.0	(0.0-0.0)	5.5	(2.0-14.2)*	0.2	(0.0-1.0)*
	Retired	2,572	70.3	(67.3-73.2)	24.2	(21.4-27.2)	1.1	(0.5-2.2)*	3.4	(2.5-4.7)	0.9	(0.5-1.6)
	Unable to work	845	62.5	(56.8-67.8)	29.0	(23.5-35.0)	0.7	(0.2-2.4)*	6.2	(4.1-9.4)	1.6	(0.9-2.8)
Veteran Status	Yes	1,310	66.7	(62.3-70.8)	26.2	(22.3-30.4)	2.0	(0.6-6.1)*	3.7	(2.3-5.9)	1.4	(0.4-4.5)*
	No	7,443	68.2	(66.1-70.2)	25.6	(23.7-27.6)	0.9	(0.6-1.5)	3.8	(3.0-4.7)	1.5	(1.0-2.1)
Diabetes	Yes	1,287	67.2	(62.5-71.7)	28.5	(24.1-33.5)	0.4	(0.1-1.2)*	2.6	(1.7-3.9)	1.2	(0.7-2.2)*
	No	7,466	68.2	(66.1-70.2)	25.3	(23.5-27.2)	1.2	(0.7-1.9)	3.9	(3.1-4.8)	1.5	(1.0-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	1,107	66.9	(62.3-71.3)	27.5	(23.2-32.2)	0.9	(0.3-2.2)*	3.9	(2.7-5.7)	0.9	(0.5-1.6)*
	No	7,550	68.3	(66.2-70.3)	25.4	(23.5-27.3)	1.1	(0.7-1.8)	3.7	(3.0-4.6)	1.5	(1.0-2.3)
Asthma	Yes	1,095	65.8	(60.1-71.0)	27.1	(22.1-32.7)	1.3	(0.4-3.5)*	5.1	(3.4-7.7)	0.8	(0.4-1.7)*
	No	7,644	68.4	(66.4-70.4)	25.5	(23.6-27.4)	1.0	(0.6-1.7)	3.6	(2.9-4.5)	1.5	(1.0-2.2)
Arthritis	Yes	3,624	68.3	(65.4-71.0)	24.7	(22.0-27.5)	1.5	(0.8-2.9)*	4.1	(3.0-5.6)	1.4	(0.8-2.5)
	No	5,104	68.0	(65.6-70.3)	26.1	(23.9-28.3)	0.9	(0.5-1.7)*	3.6	(2.8-4.6)	1.5	(0.9-2.3)
Frequent Mental Distress	Yes	984	65.9	(61.1-70.5)	26.7	(22.4-31.5)	1.1	(0.3-3.8)*	3.5	(2.3-5.4)	2.8	(1.4-5.4)*
	No	7,573	68.2	(66.1-70.2)	25.7	(23.9-27.7)	1.1	(0.6-1.8)	3.8	(3.0-4.7)	1.3	(0.8-1.9)
Serious Psychological Distress	Yes	415	57.3	(49.3-64.9)	30.0	(22.6-38.7)	1.6	(0.5-4.8)*	7.2	(4.3-11.7)	3.9	(1.9-7.8)*
	No	8,027	69.0	(67.0-70.9)	25.3	(23.5-27.2)	1.0	(0.6-1.5)	3.5	(2.8-4.4)	1.2	(0.8-1.7)
Receiving Treatment for Mental Health Condition	Yes	1,199	69.4	(64.7-73.7)	22.0	(18.0-26.7)	1.6	(0.5-4.8)*	5.7	(3.7-8.7)	1.3	(0.7-2.4)*
	No	7,527	67.9	(65.9-69.9)	26.0	(24.1-28.0)	1.0	(0.6-1.7)	3.6	(2.9-4.4)	1.5	(1.0-2.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		8,775	24.9	(23.3-26.5)	37.2	(35.3-39.2)	1.1	(0.8-1.5)	26.4	(24.7-28.1)	10.4	(9.4-11.6)
Age (years)	18-24	251	18.6	(13.4-25.1)	46.8	(37.8-56.0)	0.3	(0.1-1.0)*	25.0	(17.4-34.6)	9.3	(5.4-15.6)
	25-34	629	21.0	(16.8-25.8)	39.8	(34.6-45.3)	1.1	(0.3-3.6)*	29.4	(24.7-34.6)	8.7	(6.3-11.9)
	35-54	2,803	22.8	(20.5-25.2)	37.1	(34.3-40.0)	1.1	(0.7-1.7)	27.2	(24.8-29.7)	11.9	(10.2-13.8)
	55+	5,024	32.0	(30.0-34.1)	32.9	(30.8-35.1)	1.3	(0.9-2.0)	23.6	(21.8-25.5)	10.2	(9.0-11.5)
Sex	Men	3,332	26.9	(24.4-29.5)	42.2	(39.2-45.2)	1.1	(0.6-1.9)	22.3	(19.9-24.9)	7.5	(6.2-9.1)
	Women	5,443	23.1	(21.3-24.9)	32.6	(30.4-34.9)	1.1	(0.8-1.5)	30.2	(28.1-32.5)	13.0	(11.6-14.6)
Race/ Ethnicity	Hispanic or Latino	135	27.6	(17.7-40.5)	40.9	(27.8-55.3)	1.3	(0.3-5.7)*	20.8	(11.3-35.1)	9.4	(5.0-16.7)*
	Not Hispanic or Latino:	5,848	23.6	(21.9-25.3)	37.1	(35.0-39.2)	1.4	(1.1-1.8)	27.2	(25.3-29.1)	10.8	(9.6-12.1)
	White											
	Black or African American	2,317	31.1	(27.7-34.7)	34.0	(30.0-38.3)	1.3	(0.7-2.5)*	22.6	(19.1-26.4)	11.0	(8.8-13.7)
	All Other Races	278	22.6	(16.0-30.9)	36.3	(27.4-46.3)	1.6	(0.7-3.4)*	27.5	(20.4-36.0)	12.0	(7.2-19.5)
Education	<High School	1,072	33.3	(28.3-38.6)	38.9	(33.1-45.0)	1.0	(0.5-2.0)*	18.1	(14.6-22.2)	8.8	(6.5-11.7)
	High School Graduate (or GED)	2,734	29.7	(26.7-32.8)	37.7	(34.2-41.3)	0.5	(0.3-0.9)	23.5	(20.4-26.9)	8.6	(7.0-10.5)
	Some College	2,262	24.9	(22.0-27.9)	35.4	(32.0-38.9)	1.3	(0.8-2.2)	28.1	(25.0-31.4)	10.4	(8.6-12.5)
	College Graduate	2,692	18.2	(15.9-20.8)	37.8	(34.6-41.1)	1.6	(1.0-2.8)	29.3	(26.6-32.3)	13.0	(10.9-15.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,694	25.6	(22.3-29.3)	34.1	(29.4-39.2)	0.6	(0.3-1.2)*	28.0	(23.6-32.9)	11.6	(9.2-14.6)
	\$20,000-\$34,999	1,773	27.2	(23.8-30.8)	37.0	(32.8-41.3)	1.2	(0.6-2.3)*	24.7	(21.0-28.7)	10.0	(8.1-12.4)
	\$35,000-\$49,999	1,113	25.2	(21.3-29.6)	36.9	(32.0-42.0)	1.1	(0.5-2.3)*	26.3	(22.3-30.8)	10.5	(7.6-14.2)
	\$50,000-\$74,999	1,174	21.3	(18.1-25.0)	38.6	(34.2-43.1)	0.7	(0.3-1.9)*	28.6	(24.8-32.8)	10.7	(8.4-13.5)
	\$75,000 and more	1,723	21.6	(18.6-25.1)	39.6	(35.8-43.4)	1.4	(0.7-2.7)*	26.5	(23.5-29.8)	10.9	(8.6-13.6)
Employment Status	Employed	4,053	24.4	(22.2-26.7)	37.8	(35.3-40.4)	1.1	(0.6-1.7)	26.7	(24.6-29.0)	10.0	(8.6-11.7)
	Out of work	584	22.9	(18.0-28.7)	35.9	(29.4-43.0)	0.7	(0.3-1.5)*	28.5	(21.6-36.5)	12.0	(8.6-16.7)
	Homemaker	546	23.7	(18.9-29.1)	40.6	(34.0-47.5)	1.4	(0.5-3.8)*	24.6	(19.4-30.6)	9.9	(6.8-14.0)
	Student	127	21.4	(13.0-33.2)	44.7	(32.2-58.0)	0.4	(0.1-2.9)*	25.5	(15.7-38.8)	7.9	(3.2-18.1)*
	Retired	2,587	30.9	(28.1-33.9)	35.5	(32.5-38.7)	1.1	(0.7-1.9)	23.1	(20.6-25.8)	9.3	(7.7-11.2)
	Unable to work	852	21.4	(17.5-25.9)	29.3	(23.4-36.0)	2.0	(1.0-3.9)*	30.8	(25.5-36.6)	16.5	(12.8-21.0)
Veteran Status	Yes	1,308	24.4	(21.0-28.2)	38.6	(34.1-43.3)	1.1	(0.5-2.4)*	26.9	(22.9-31.2)	9.0	(6.6-12.2)
	No	7,457	25.0	(23.3-26.8)	37.0	(34.9-39.1)	1.1	(0.8-1.5)	26.3	(24.5-28.2)	10.6	(9.5-11.9)
Diabetes	Yes	1,275	24.2	(20.8-28.0)	39.1	(34.0-44.5)	1.3	(0.6-2.8)*	24.7	(20.9-28.9)	10.8	(8.4-13.7)
	No	7,492	25.0	(23.3-26.8)	37.0	(35.0-39.1)	1.1	(0.8-1.5)	26.5	(24.8-28.4)	10.4	(9.3-11.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

South Carolina, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	1,118	25.5	(22.2-29.2)	33.1	(28.5-38.0)	0.9	(0.4-2.0)*	24.3	(20.4-28.8)	16.2	(12.7-20.5)
	No	7,550	24.8	(23.1-26.5)	37.6	(35.6-39.6)	1.1	(0.8-1.5)	26.6	(24.9-28.5)	9.9	(8.8-11.1)
Asthma	Yes	1,093	24.2	(20.3-28.6)	33.7	(28.5-39.3)	1.2	(0.6-2.5)*	27.7	(23.0-32.9)	13.2	(10.1-17.1)
	No	7,661	24.9	(23.3-26.7)	37.8	(35.8-39.8)	1.1	(0.8-1.5)	26.3	(24.5-28.1)	10.0	(8.9-11.2)
Arthritis	Yes	3,639	24.5	(22.4-26.8)	35.4	(32.4-38.4)	1.2	(0.8-1.8)	27.0	(24.4-29.8)	11.9	(10.2-13.9)
	No	5,101	25.0	(23.0-27.1)	38.0	(35.6-40.4)	1.0	(0.7-1.6)	26.1	(24.1-28.3)	9.8	(8.5-11.2)
Frequent Mental Distress	Yes	989	21.0	(17.1-25.4)	27.6	(22.9-32.9)	1.1	(0.5-2.3)*	31.8	(27.0-37.0)	18.7	(15.0-22.9)
	No	7,585	25.3	(23.6-27.1)	38.5	(36.5-40.6)	1.1	(0.8-1.5)	25.7	(24.0-27.5)	9.4	(8.3-10.6)
Serious Psychological Distress	Yes	414	14.8	(10.7-20.1)	25.8	(18.1-35.4)	1.7	(0.5-4.9)*	33.7	(25.8-42.7)	24.0	(18.0-31.1)
	No	8,053	25.3	(23.7-27.1)	37.8	(35.8-39.8)	1.1	(0.8-1.5)	26.1	(24.4-27.9)	9.7	(8.6-10.8)
Receiving Treatment for Mental Health Condition	Yes	1,187	18.0	(14.8-21.7)	28.6	(24.1-33.6)	1.3	(0.7-2.4)*	34.5	(29.5-39.8)	17.7	(14.4-21.5)
	No	7,556	26.2	(24.5-27.9)	37.8	(35.8-39.9)	1.3	(1.0-1.7)	25.3	(23.5-27.1)	9.5	(8.4-10.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,124	75.5	(72.5-78.4)	17.2	(14.7-20.1)	3.0	(1.9-4.8)	2.9	(1.9-4.5)	1.3	(0.6-2.5)*
Age (years)												
18-24		63	69.9	(54.6-81.8)	20.8	(11.4-35.0)	0.5	(0.1-3.6)*	8.7	(2.9-23.2)*	0.1	(0.0-0.1)
25-34		175	71.0	(61.5-79.0)	11.1	(6.4-18.6)	11.0	(6.0-19.3)	3.3	(1.3-7.9)*	3.5	(0.8-14.6)*
35-54		685	76.4	(71.5-80.7)	18.9	(14.8-23.9)	1.7	(1.0-3.1)	1.9	(1.0-3.5)*	1.1	(0.5-2.5)*
55+		1,191	78.3	(75.2-81.2)	15.2	(12.6-18.1)	3.0	(2.0-4.5)	2.3	(1.6-3.5)	1.1	(0.6-2.3)*
Sex												
Men		672	71.2	(65.8-76.0)	19.0	(14.8-24.0)	3.9	(2.1-7.4)*	4.2	(2.3-7.4)	1.7	(0.6-4.8)*
Women		1,452	79.6	(76.4-82.4)	15.6	(13.0-18.7)	2.2	(1.4-3.5)	1.7	(1.1-2.6)	0.9	(0.5-1.6)*
Race/Ethnicity												
Hispanic or Latino		10	--	--	--	--	--	--	--	--	--	--
Not Hispanic or Latino:												
White		1,791	74.6	(71.5-77.5)	17.4	(14.9-20.4)	4.2	(3.0-5.9)	2.6	(1.8-3.7)	1.1	(0.6-2.2)*
Black or African American		277	72.5	(62.0-81.0)	20.9	(13.1-31.6)	2.4	(1.0-5.7)*	3.5	(1.7-7.0)*	0.7	(0.3-1.5)*
All Other Races		41	--	--	--	--	--	--	--	--	--	--
Education												
<High School		285	66.6	(57.0-75.1)	22.8	(15.0-33.0)	3.3	(1.7-6.2)*	3.2	(1.7-6.1)*	4.1	(1.1-14.3)*
High School Graduate (or GED)		747	72.0	(66.3-77.0)	19.2	(14.8-24.4)	3.5	(1.7-6.9)*	4.2	(2.1-8.2)*	1.2	(0.5-3.0)*
Some College		648	78.0	(72.4-82.8)	18.2	(13.6-23.8)	1.3	(0.6-3.0)*	1.8	(0.9-3.6)*	0.7	(0.3-1.8)*
College Graduate		442	78.3	(71.7-83.8)	12.9	(8.9-18.3)	5.5	(2.3-12.6)*	2.7	(1.0-7.0)*	0.6	(0.1-2.5)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	439	72.6 (65.3-78.9)	18.0 (12.6-25.0)	2.1 (1.0-4.2)*	3.5 (1.9-6.3)*	3.7 (1.3-10.3)*				
	\$20,000-\$34,999	555	76.0 (69.5-81.4)	15.3 (10.7-21.3)	3.8 (1.5-9.3)*	4.5 (2.0-10.1)*	0.4 (0.1-2.5)*				
	\$35,000-\$49,999	263	71.1 (61.9-78.8)	22.4 (15.1-31.8)	4.1 (1.8-9.3)*	2.1 (0.7-6.7)*	0.3 (0.0-2.3)*				
	\$50,000-\$74,999	185	82.1 (72.0-89.1)	13.6 (8.2-21.8)	4.0 (0.8-18.2)*	0.3 (0.0-1.8)*	0.0 (0.0-0.0)				
	\$75,000 and more	263	80.8 (72.6-87.0)	14.2 (8.8-22.3)	0.7 (0.2-2.9)*	1.9 (0.5-7.2)*	2.3 (0.8-6.9)*				
Employment Status	Employed	840	76.7 (72.1-80.7)	17.1 (13.5-21.5)	2.9 (1.5-5.4)*	2.3 (1.2-4.6)*	1.0 (0.4-2.3)*				
	Out of work	120	74.7 (62.6-83.9)	10.1 (5.4-18.2)*	5.1 (1.2-18.9)*	6.0 (2.0-16.6)*	4.1 (0.9-17.2)*				
	Homemaker	204	77.3 (67.6-84.7)	18.3 (11.2-28.3)	0.3 (0.0-2.4)*	3.2 (1.1-8.9)*	1.0 (0.2-5.2)*				
	Student	34	--	--	--	--	--	--			
	Retired	671	73.3 (68.4-77.6)	19.4 (15.4-24.2)	3.9 (2.4-6.3)	3.0 (1.8-5.1)	0.5 (0.2-1.3)*				
	Unable to work	247	71.9 (61.8-80.2)	21.6 (13.6-32.5)	2.4 (1.0-5.4)*	2.4 (1.1-5.2)*	1.8 (0.7-4.4)*				
Veteran Status	Yes	212	82.1 (73.7-88.2)	12.4 (7.0-21.1)	3.5 (1.7-7.1)*	1.1 (0.4-2.8)*	0.9 (0.2-4.1)*				
	No	1,909	74.7 (71.4-77.8)	17.9 (15.2-21.0)	3.0 (1.8-4.9)	3.1 (2.0-4.9)	1.3 (0.6-2.8)*				
Diabetes	Yes	268	66.7 (57.7-74.7)	25.1 (17.5-34.8)	4.4 (2.1-9.1)*	1.7 (0.8-3.8)*	1.9 (0.7-5.3)*				
	No	1,854	76.4 (73.1-79.3)	16.5 (13.9-19.5)	2.9 (1.8-4.8)	3.0 (1.9-4.7)	1.2 (0.5-2.6)*				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	284	74.5	(67.4-80.4)	17.7	(12.2-25.0)	2.9	(1.2-7.0)*	4.4	(2.6-7.4)	0.5	(0.1-2.4)*
	No	1,834	75.6	(72.3-78.7)	17.2	(14.5-20.2)	3.0	(1.9-5.0)	2.8	(1.7-4.5)	1.4	(0.7-2.8)*
Asthma	Yes	254	76.8	(67.7-84.0)	15.6	(9.5-24.5)	1.7	(0.6-4.4)*	4.6	(2.0-10.3)*	1.3	(0.3-5.2)*
	No	1,869	75.4	(72.1-78.4)	17.5	(14.8-20.5)	3.2	(2.0-5.3)	2.6	(1.6-4.4)	1.3	(0.6-2.7)*
Arthritis	Yes	722	74.0	(68.1-79.1)	16.8	(12.4-22.5)	4.5	(1.9-10.3)*	2.7	(1.5-4.8)	2.0	(0.6-7.0)*
	No	1,394	76.1	(72.5-79.4)	17.4	(14.5-20.8)	2.5	(1.5-4.2)	3.0	(1.7-5.0)	1.0	(0.5-1.9)*
Frequent Mental Distress	Yes	266	66.8	(56.9-75.4)	22.7	(15.0-33.0)	3.0	(1.2-7.0)*	2.3	(1.0-4.9)*	5.2	(1.9-13.4)*
	No	1,828	77.2	(73.9-80.1)	16.1	(13.5-19.1)	3.1	(1.8-5.1)	3.0	(1.9-4.9)	0.6	(0.3-1.2)*
Serious Psychological Distress	Yes	189	59.6	(48.8-69.6)	30.1	(20.6-41.7)	1.5	(0.6-3.8)*	3.2	(1.6-6.5)*	5.6	(1.7-16.6)*
	No	1,885	77.7	(74.5-80.6)	15.8	(13.2-18.8)	2.9	(1.8-4.5)	2.9	(1.8-4.7)	0.7	(0.4-1.3)*
Receiving Treatment for Mental Health Condition	Yes	402	78.4	(71.5-84.0)	14.7	(9.9-21.3)	1.7	(0.7-4.1)*	2.2	(1.2-4.0)*	2.9	(0.8-10.5)*
	No	1,718	75.2	(71.8-78.3)	17.6	(14.8-20.8)	3.3	(2.0-5.3)	3.0	(1.8-4.9)	0.9	(0.5-1.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,100	35.9	(32.8-39.1)	27.9	(25.0-31.0)	7.1	(5.3-9.4)	21.4	(18.8-24.2)	7.7	(6.3-9.4)
Age (years)	18-24	65	25.1	(14.3-40.1)	38.5	(24.9-54.1)	11.1	(3.6-29.2)*	15.3	(7.2-29.6)*	10.1	(3.7-24.7)*
	25-34	173	34.0	(25.4-43.8)	35.2	(26.7-44.8)	8.0	(3.2-18.5)*	19.0	(13.0-26.8)	3.8	(2.0-7.2)*
	35-54	678	35.7	(31.0-40.8)	26.3	(21.9-31.2)	5.5	(3.7-7.9)	23.3	(19.2-28.0)	9.2	(6.8-12.3)
	55+	1,174	40.8	(37.2-44.5)	23.6	(20.5-27.1)	7.6	(6.0-9.7)	20.5	(17.5-23.9)	7.4	(5.8-9.5)
Sex	Men	667	38.3	(33.3-43.6)	29.4	(24.7-34.7)	6.9	(4.2-11.0)	19.0	(15.1-23.7)	6.4	(4.3-9.3)
	Women	1,433	33.7	(30.3-37.2)	26.4	(23.1-29.9)	7.3	(5.6-9.6)	23.7	(20.6-27.0)	8.9	(7.1-11.1)
Race/ Ethnicity	Hispanic or Latino	11	--	--	--	--	--	--	--	--	--	--
	Not Hispanic or Latino:	1,776	32.5	(29.3-35.8)	26.6	(23.4-30.1)	13.1	(11.4-15.0)	20.5	(17.6-23.6)	7.3	(5.8-9.3)
	White											
	Black or African American	268	48.1	(38.5-57.9)	23.4	(15.5-33.6)	2.9	(1.4-6.0)*	19.2	(12.8-27.8)	6.4	(3.5-11.3)*
	All Other Races	40	--	--	--	--	--	--	--	--	--	--
Education	<High School	276	44.7	(35.2-54.5)	26.4	(18.4-36.3)	4.6	(2.6-7.8)	19.0	(12.4-28.0)	5.3	(2.7-10.3)*
	High School Graduate (or GED)	745	38.3	(33.1-43.8)	29.3	(24.3-34.9)	8.5	(5.1-13.7)	16.3	(12.8-20.5)	7.6	(5.1-11.2)
	Some College	639	30.8	(25.7-36.4)	26.7	(21.7-32.3)	9.4	(6.8-13.0)	23.9	(19.3-29.2)	9.2	(6.9-12.3)
	College Graduate	438	41.0	(34.7-47.6)	26.5	(21.3-32.5)	2.8	(1.6-4.8)	22.8	(17.6-29.0)	6.8	(4.3-10.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	434	40.7	(33.8-47.9)	24.6	(18.7-31.6)	4.1	(2.4-6.8)	21.4	(15.8-28.3)	9.4	(6.1-14.0)
	\$20,000-\$34,999	546	28.4	(23.1-34.5)	24.2	(19.1-30.3)	16.9	(11.3-24.7)	23.1	(18.1-28.9)	7.3	(4.7-11.2)
	\$35,000-\$49,999	260	43.3	(34.5-52.4)	30.4	(22.6-39.5)	4.7	(2.3-9.3)*	15.5	(10.4-22.6)	6.1	(3.1-11.8)*
	\$50,000-\$74,999	185	34.7	(25.8-44.8)	24.3	(16.9-33.7)	3.2	(1.0-9.4)*	30.4	(22.4-39.8)	7.4	(4.4-12.3)
	\$75,000 and more	263	38.4	(31.0-46.3)	32.2	(24.7-40.7)	1.8	(0.8-4.0)*	19.5	(13.4-27.4)	8.2	(5.1-13.0)
Employment Status	Employed	830	37.6	(33.0-42.4)	27.6	(23.5-32.2)	4.3	(2.8-6.4)	22.5	(18.7-26.8)	8.0	(5.9-10.7)
	Out of work	120	26.6	(17.3-38.5)	29.7	(19.1-43.2)	19.3	(8.3-38.5)*	17.1	(8.9-30.4)*	7.3	(2.8-17.8)*
	Homemaker	200	36.7	(28.8-45.4)	27.7	(19.8-37.3)	6.6	(3.5-12.3)*	22.0	(15.8-29.7)	6.9	(3.5-13.1)*
	Student	34	--	--	--	--	--	--	--	--	--	--
	Retired	663	37.7	(33.0-42.6)	27.2	(22.6-32.4)	9.7	(7.1-13.3)	20.1	(16.2-24.5)	5.3	(3.5-7.9)
	Unable to work	244	34.3	(26.0-43.7)	19.7	(13.4-28.2)	7.2	(3.8-13.0)*	24.7	(17.5-33.7)	14.0	(8.9-21.5)
Veteran Status	Yes	211	30.6	(24.0-38.1)	34.2	(25.2-44.6)	3.3	(1.7-6.2)*	22.8	(15.8-31.7)	9.1	(5.4-15.0)
	No	1,886	36.5	(33.2-40.0)	27.1	(24.1-30.4)	7.6	(5.7-10.1)	21.2	(18.5-24.2)	7.6	(6.1-9.4)
Diabetes	Yes	263	35.4	(28.1-43.4)	25.6	(18.6-34.2)	6.2	(3.5-10.6)	21.9	(15.6-29.9)	10.9	(5.8-19.5)*
	No	1,835	36.1	(32.8-39.5)	28.0	(24.9-31.3)	7.2	(5.3-9.6)	21.4	(18.6-24.3)	7.4	(6.0-9.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Tennessee, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	276	30.2	(24.1-37.2)	31.0	(23.3-39.8)	8.6	(4.9-14.8)	22.8	(16.1-31.2)	7.4	(4.8-11.3)
	No	1,818	36.5	(33.2-39.9)	27.6	(24.5-30.9)	7.0	(5.1-9.4)	21.2	(18.5-24.2)	7.7	(6.2-9.6)
Asthma	Yes	253	29.2	(21.6-38.0)	25.1	(17.0-35.4)	5.0	(2.6-9.5)*	28.9	(20.5-39.0)	11.9	(6.9-19.6)
	No	1,846	36.9	(33.6-40.3)	28.3	(25.2-31.6)	7.4	(5.5-10.0)	20.3	(17.7-23.2)	7.1	(5.7-8.8)
Arthritis	Yes	712	35.4	(30.3-40.9)	24.2	(19.3-29.8)	6.9	(4.1-11.4)	23.1	(18.5-28.4)	10.4	(7.6-14.1)
	No	1,380	36.0	(32.3-39.8)	29.1	(25.6-32.9)	7.2	(5.2-9.9)	20.8	(17.8-24.2)	6.9	(5.3-9.0)
Frequent Mental Distress	Yes	266	20.6	(14.5-28.5)	29.1	(20.5-39.6)	4.2	(2.4-7.3)	27.3	(19.8-36.4)	18.8	(12.8-26.6)
	No	1,806	37.9	(34.6-41.3)	27.6	(24.5-30.9)	7.5	(5.6-10.1)	20.7	(18.0-23.7)	6.3	(4.9-8.0)
Serious Psychological Distress	Yes	188	25.3	(17.2-35.5)	20.8	(13.6-30.6)	4.0	(1.9-8.2)*	27.7	(18.7-38.9)	22.2	(14.3-32.8)
	No	1,866	37.0	(33.7-40.4)	28.3	(25.2-31.6)	7.3	(5.4-9.8)	20.9	(18.2-23.8)	6.5	(5.2-8.2)
Receiving Treatment for Mental Health Condition	Yes	398	27.7	(22.0-34.2)	26.5	(20.0-34.1)	8.1	(5.4-12.0)	27.2	(20.8-34.8)	10.5	(7.1-15.4)
	No	1,698	37.0	(33.6-40.6)	26.3	(23.1-29.7)	11.0	(8.9-13.4)	18.9	(16.2-22.0)	6.8	(5.3-8.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		7,045	63.6	(61.9-65.3)	26.4	(24.7-28.0)	4.3	(3.6-5.1)	4.2	(3.5-5.0)	1.5	(1.2-1.9)
Age (years)	18-24	250	51.0	(42.9-59.1)	34.9	(27.0-43.6)	5.8	(3.3-10.2)	7.4	(4.3-12.4)	0.8	(0.3-2.5)*
	25-34	872	58.0	(53.4-62.5)	32.6	(28.3-37.2)	4.2	(2.7-6.4)	4.4	(2.8-6.6)	0.8	(0.4-2.0)*
	35-54	2,560	66.9	(64.2-69.5)	23.5	(21.1-25.9)	3.9	(2.9-5.2)	3.6	(2.7-5.0)	2.1	(1.5-3.0)
	55+	3,305	67.6	(65.4-69.7)	23.5	(21.6-25.5)	4.2	(3.4-5.1)	3.3	(2.6-4.3)	1.4	(1.0-2.0)
Sex	Men	2,363	57.8	(54.9-60.7)	30.9	(28.2-33.8)	5.2	(4.1-6.7)	4.3	(3.2-5.8)	1.7	(1.2-2.6)
	Women	4,682	69.1	(67.1-71.0)	22.1	(20.4-23.9)	3.5	(2.8-4.2)	4.1	(3.3-5.1)	1.3	(0.9-1.8)
Race/ Ethnicity	Hispanic or Latino	2,100	62.5	(59.0-65.9)	24.7	(21.8-27.9)	5.2	(3.9-6.9)	5.1	(3.9-6.7)	2.5	(1.7-3.8)
	Not Hispanic or Latino:	4,189	65.9	(63.8-67.9)	26.6	(24.7-28.6)	3.7	(3.0-4.6)	2.6	(2.0-3.4)	1.2	(0.8-1.7)
	White											
	Black or African American	432	57.9	(51.6-64.1)	27.7	(21.8-34.4)	5.3	(3.3-8.6)	7.3	(4.5-11.7)	1.7	(0.8-3.5)*
	All Other Races	255	56.8	(48.1-65.1)	31.9	(23.9-41.0)	3.9	(1.7-8.5)*	6.8	(2.9-14.9)*	0.7	(0.3-1.7)*
Education	<High School	1,067	58.2	(53.5-62.9)	24.7	(20.7-29.2)	9.0	(6.5-12.2)	5.8	(3.8-8.8)	2.2	(1.4-3.7)
	High School Graduate (or GED)	1,862	59.9	(56.3-63.5)	29.6	(26.3-33.2)	4.4	(3.3-5.8)	4.8	(3.5-6.5)	1.2	(0.8-2.0)
	Some College	1,887	64.4	(61.1-67.6)	25.9	(22.9-29.2)	4.6	(3.4-6.3)	3.7	(2.7-5.1)	1.3	(0.8-2.2)
	College Graduate	2,208	67.7	(64.7-70.6)	24.6	(21.9-27.4)	3.2	(2.3-4.4)	3.0	(1.9-4.9)	1.5	(0.8-2.8)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,416	60.6 (56.0-65.0)	27.1 (22.9-31.7)	4.9 (3.4-7.2)	5.5 (3.8-8.0)	1.9 (1.2-2.9)				
	\$20,000-\$34,999	1,303	61.4 (57.4-65.4)	27.4 (23.9-31.3)	4.6 (2.9-7.0)	4.9 (3.6-6.8)	1.6 (0.9-3.0)*				
	\$35,000-\$49,999	889	68.5 (63.7-72.9)	22.5 (18.6-27.0)	5.3 (3.5-8.0)	2.7 (1.6-4.4)	0.9 (0.4-2.0)*				
	\$50,000-\$74,999	927	64.1 (59.5-68.5)	25.9 (21.8-30.4)	5.0 (3.3-7.5)	4.7 (2.8-7.9)	0.3 (0.1-1.0)*				
	\$75,000 and more	1,533	65.7 (62.3-69.0)	27.0 (23.8-30.4)	3.0 (2.0-4.5)	2.6 (1.6-4.2)	1.6 (0.8-3.2)*				
Employment Status	Employed	3,506	64.1 (61.8-66.3)	26.5 (24.4-28.7)	4.3 (3.4-5.4)	3.6 (2.8-4.7)	1.6 (1.1-2.3)				
	Out of work	286	62.0 (52.8-70.4)	28.7 (20.7-38.3)	4.4 (2.0-9.4)*	4.5 (1.9-10.6)*	0.4 (0.1-1.6)*				
	Homemaker	826	60.0 (54.9-64.8)	29.9 (25.2-35.1)	3.0 (1.9-4.8)	5.9 (3.7-9.1)	1.3 (0.6-2.7)*				
	Student	130	64.0 (51.9-74.5)	23.0 (14.4-34.8)	7.0 (2.8-16.5)*	5.0 (2.0-12.0)*	1.0 (0.1-6.4)*				
	Retired	1,766	64.7 (61.6-67.6)	25.9 (23.1-28.8)	4.6 (3.6-6.0)	3.3 (2.3-4.8)	1.5 (0.9-2.6)				
	Unable to work	502	65.4 (58.5-71.7)	21.1 (16.1-27.3)	3.9 (2.2-6.7)	7.0 (4.5-10.8)	2.6 (1.5-4.3)				
Veteran Status	Yes	941	60.4 (56.0-64.7)	29.8 (25.7-34.2)	3.3 (2.2-5.0)	4.4 (2.5-7.5)	2.1 (1.1-3.9)*				
	No	6,100	64.1 (62.2-66.0)	25.8 (24.1-27.7)	4.4 (3.7-5.3)	4.2 (3.4-5.0)	1.4 (1.1-1.9)				
Diabetes	Yes	933	64.0 (59.1-68.6)	25.6 (21.5-30.2)	2.8 (1.7-4.4)	5.0 (3.3-7.4)	2.7 (1.4-4.8)*				
	No	6,104	63.5 (61.6-65.3)	26.4 (24.6-28.2)	4.7 (4.0-5.5)	4.1 (3.4-5.0)	1.4 (1.1-1.8)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	852	64.7	(59.8-69.3)	27.6	(23.2-32.5)	1.8	(1.2-2.8)	4.5	(2.9-6.9)	1.3	(0.7-2.5)*
	No	6,126	63.5	(61.6-65.3)	26.2	(24.5-28.0)	4.6	(3.9-5.5)	4.2	(3.4-5.1)	1.5	(1.2-2.0)
Asthma	Yes	924	65.0	(60.1-69.5)	24.9	(20.5-30.0)	2.4	(1.5-3.8)	6.5	(4.2-9.8)	1.2	(0.7-2.3)*
	No	6,111	63.2	(61.3-65.0)	26.5	(24.8-28.3)	4.9	(4.2-5.8)	3.8	(3.1-4.7)	1.6	(1.2-2.1)
Arthritis	Yes	2,316	65.1	(62.2-67.9)	26.0	(23.2-28.9)	2.8	(2.1-3.7)	4.3	(3.2-5.8)	1.8	(1.2-2.7)
	No	4,689	63.1	(61.0-65.2)	26.6	(24.6-28.6)	4.7	(3.9-5.7)	4.1	(3.3-5.2)	1.4	(1.0-2.0)
Frequent Mental Distress	Yes	746	59.7	(54.6-64.6)	26.3	(21.8-31.4)	3.3	(1.8-6.2)*	8.4	(5.8-12.2)	2.2	(1.4-3.5)
	No	6,190	64.0	(62.2-65.8)	26.5	(24.8-28.3)	4.4	(3.7-5.3)	3.7	(3.0-4.5)	1.4	(1.0-1.9)
Serious Psychological Distress	Yes	337	58.0	(49.9-65.6)	23.0	(16.4-31.2)	3.5	(1.0-11.3)*	13.1	(8.8-19.1)	2.4	(1.3-4.4)*
	No	6,541	63.9	(62.1-65.7)	26.2	(24.5-28.0)	4.6	(3.9-5.4)	3.9	(3.2-4.7)	1.4	(1.1-1.9)
Receiving Treatment for Mental Health Condition	Yes	883	71.7	(67.2-75.9)	19.9	(16.0-24.3)	1.0	(0.5-1.8)*	5.8	(3.6-9.1)	1.7	(0.9-3.0)*
	No	6,141	62.6	(60.7-64.4)	27.2	(25.4-29.0)	4.8	(4.0-5.6)	4.0	(3.3-4.9)	1.5	(1.1-2.0)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		7,032	25.1	(23.5-26.7)	35.9	(34.2-37.7)	5.8	(5.1-6.7)	21.9	(20.5-23.5)	11.2	(10.2-12.3)
Age (years)	18-24	246	24.8	(18.8-31.9)	41.2	(32.7-50.3)	6.3	(3.3-11.7)*	22.2	(15.8-30.2)	5.5	(3.1-9.8)
	25-34	865	20.4	(17.2-24.2)	38.5	(34.0-43.2)	6.5	(4.7-8.8)	24.6	(20.8-28.7)	10.0	(7.6-13.2)
	35-54	2,567	25.1	(22.6-27.8)	33.9	(31.3-36.5)	5.4	(4.3-6.8)	22.4	(20.2-24.8)	13.1	(11.4-15.1)
	55+	3,294	28.4	(26.4-30.4)	34.8	(32.7-37.1)	5.9	(4.9-7.0)	19.9	(18.2-21.7)	11.1	(9.7-12.6)
Sex	Men	2,380	27.3	(24.6-30.1)	38.4	(35.5-41.4)	6.0	(4.8-7.5)	18.1	(15.9-20.4)	10.2	(8.6-12.1)
	Women	4,652	23.0	(21.4-24.7)	33.6	(31.6-35.6)	5.7	(4.8-6.6)	25.7	(23.9-27.6)	12.1	(10.8-13.4)
Race/ Ethnicity	Hispanic or Latino	2,085	34.0	(30.8-37.4)	33.3	(29.9-36.9)	5.5	(4.0-7.4)	18.3	(15.6-21.3)	8.9	(7.1-11.1)
	Not Hispanic or Latino:	4,195	21.1	(19.5-22.9)	37.6	(35.5-39.6)	5.8	(4.9-6.9)	23.8	(22.1-25.6)	11.7	(10.4-13.1)
	White											
	Black or African American	433	28.0	(21.7-35.2)	35.0	(28.6-42.1)	5.5	(3.3-8.8)	18.1	(13.5-23.9)	13.4	(9.8-18.2)
	All Other Races	252	36.1	(27.5-45.9)	31.2	(22.7-41.1)	5.8	(3.3-10.0)	16.4	(10.6-24.6)	10.5	(6.9-15.6)
Education	<High School	1,057	35.8	(31.3-40.5)	30.9	(26.3-36.0)	5.5	(3.8-7.8)	17.8	(13.8-22.7)	10.0	(7.3-13.6)
	High School Graduate (or GED)	1,863	29.0	(25.9-32.3)	34.5	(31.0-38.2)	6.4	(4.8-8.4)	19.2	(16.7-22.0)	10.8	(8.8-13.2)
	Some College	1,879	22.2	(19.5-25.2)	37.1	(33.7-40.5)	6.6	(5.0-8.6)	22.3	(19.7-25.2)	11.8	(10.0-14.0)
	College Graduate	2,211	20.6	(17.9-23.6)	38.3	(35.4-41.3)	4.9	(3.9-6.2)	24.8	(22.3-27.6)	11.4	(9.6-13.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	1,407	28.2 (24.0-32.8)	33.7 (29.0-38.8)	6.6 (4.8-9.1)	19.4 (15.8-23.5)	12.1 (9.8-14.9)				
	\$20,000-\$34,999	1,315	26.4 (23.1-30.0)	33.6 (29.8-37.7)	4.6 (3.2-6.6)	22.7 (19.2-26.6)	12.7 (10.1-15.7)				
	\$35,000-\$49,999	881	28.5 (24.0-33.5)	37.1 (32.3-42.3)	5.2 (3.6-7.5)	20.2 (16.6-24.4)	8.9 (6.6-11.8)				
	\$50,000-\$74,999	930	22.9 (19.3-26.9)	37.0 (32.6-41.6)	5.3 (3.7-7.4)	22.4 (19.0-26.1)	12.5 (9.6-16.1)				
	\$75,000 and more	1,531	21.6 (18.8-24.8)	37.8 (34.5-41.2)	6.1 (4.6-8.0)	22.8 (20.1-25.8)	11.6 (9.6-14.0)				
Employment Status	Employed	3,507	25.5 (23.5-27.6)	35.7 (33.5-38.1)	5.7 (4.8-6.8)	22.2 (20.3-24.3)	10.9 (9.5-12.4)				
	Out of work	279	18.7 (13.2-26.0)	45.5 (35.6-55.7)	5.8 (3.0-10.9)*	20.9 (14.6-28.8)	9.1 (4.8-16.6)*				
	Homemaker	822	27.4 (23.4-31.8)	32.8 (28.1-37.9)	5.0 (3.3-7.3)	24.6 (20.3-29.5)	10.2 (7.7-13.4)				
	Student	132	20.7 (12.8-31.7)	34.5 (23.6-47.4)	19.2 (11.3-30.8)	17.2 (10.5-27.0)	8.4 (4.5-15.1)*				
	Retired	1,766	25.3 (22.8-27.9)	37.7 (34.6-40.9)	5.8 (4.4-7.6)	20.8 (18.3-23.5)	10.5 (8.7-12.5)				
	Unable to work	495	25.2 (18.1-33.8)	28.8 (22.7-35.7)	5.2 (3.2-8.6)	20.0 (14.8-26.5)	20.8 (15.9-26.7)				
Veteran Status	Yes	950	25.8 (21.0-31.2)	36.3 (32.0-40.9)	4.5 (3.1-6.4)	17.6 (14.5-21.2)	15.8 (12.1-20.4)				
	No	6,076	25.0 (23.4-26.7)	35.8 (33.9-37.8)	6.2 (5.3-7.1)	22.5 (20.9-24.2)	10.6 (9.5-11.7)				
Diabetes	Yes	928	24.1 (20.4-28.1)	31.9 (27.4-36.7)	4.3 (3.0-6.2)	24.3 (20.5-28.7)	15.4 (12.2-19.2)				
	No	6,097	25.2 (23.5-26.9)	36.3 (34.4-38.2)	6.1 (5.3-7.0)	21.7 (20.2-23.3)	10.7 (9.6-11.9)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Texas, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	849	24.2 (20.5-28.4)	32.7 (28.4-37.4)	4.6 (3.1-6.8)	23.7 (19.5-28.5)	14.7 (11.4-18.9)				
	No	6,116	25.3 (23.6-27.0)	36.2 (34.3-38.1)	6.0 (5.2-6.9)	21.7 (20.2-23.3)	10.9 (9.8-12.1)				
Asthma	Yes	917	25.1 (21.0-29.8)	30.2 (25.4-35.6)	4.5 (2.7-7.6)	24.6 (20.9-28.9)	15.4 (12.4-19.1)				
	No	6,103	25.1 (23.5-26.9)	36.9 (35.0-38.8)	6.0 (5.3-6.9)	21.5 (20.0-23.1)	10.4 (9.4-11.6)				
Arthritis	Yes	2,303	21.6 (19.4-23.9)	35.6 (32.6-38.6)	4.7 (3.7-6.1)	23.0 (20.7-25.6)	15.1 (13.0-17.4)				
	No	4,691	26.3 (24.4-28.3)	36.1 (34.0-38.2)	6.2 (5.3-7.3)	21.6 (19.9-23.5)	9.8 (8.6-11.1)				
Frequent Mental Distress	Yes	758	18.9 (14.0-24.9)	29.2 (24.3-34.6)	6.0 (3.9-9.2)	25.0 (20.6-29.9)	20.9 (16.7-26.0)				
	No	6,161	25.8 (24.2-27.5)	36.9 (35.0-38.8)	5.9 (5.1-6.8)	21.5 (20.0-23.1)	9.8 (8.8-10.9)				
Serious Psychological Distress	Yes	334	19.6 (11.8-30.8)	24.0 (16.5-33.6)	4.2 (2.2-7.8)*	26.0 (18.0-36.1)	26.2 (18.7-35.3)				
	No	6,534	25.3 (23.8-26.9)	36.5 (34.7-38.3)	5.9 (5.1-6.8)	21.8 (20.4-23.4)	10.5 (9.5-11.6)				
Receiving Treatment for Mental Health Condition	Yes	869	19.0 (15.7-22.9)	28.3 (23.9-33.1)	6.3 (4.4-9.0)	26.0 (22.3-30.2)	20.4 (16.7-24.6)				
	No	6,142	25.9 (24.2-27.6)	36.9 (35.1-38.9)	5.8 (5.0-6.7)	21.4 (19.9-23.1)	9.9 (8.9-11.1)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,363	66.8	(63.9-69.6)	26.9	(24.3-29.7)	1.0	(0.6-1.6)	4.1	(3.0-5.6)	1.2	(0.7-2.0)
Age (years)	18-24	95	56.7	(44.1-68.5)	40.0	(28.5-52.6)	0.7	(0.1-4.6)*	1.8	(0.6-5.1)*	0.9	(0.1-5.5)*
	25-34	319	66.4	(59.9-72.4)	27.9	(22.4-34.3)	0.2	(0.1-0.7)*	4.6	(2.1-9.6)*	0.9	(0.2-3.0)*
	35-54	896	67.1	(63.3-70.8)	26.0	(22.6-29.7)	1.2	(0.6-2.4)*	4.6	(3.0-7.0)	1.1	(0.5-2.7)*
	55+	1,042	68.0	(64.4-71.3)	22.9	(20.0-26.1)	2.1	(1.1-3.9)*	5.1	(3.6-7.3)	1.9	(1.0-3.6)*
Sex	Men	1,022	61.4	(56.8-65.8)	30.2	(26.2-34.6)	1.1	(0.6-2.0)	5.3	(3.6-7.8)	2.0	(1.0-3.8)*
	Women	1,341	71.8	(68.2-75.1)	23.8	(20.6-27.3)	0.9	(0.4-1.9)*	3.0	(1.9-4.7)	0.5	(0.2-1.2)*
Race/ Ethnicity	Hispanic or Latino	114	56.8	(43.7-69.0)	34.3	(23.2-47.4)	1.5	(0.4-5.3)*	4.8	(1.7-12.8)*	2.7	(0.9-7.7)*
	Not Hispanic or Latino:	2,164	67.8	(64.8-70.7)	26.2	(23.5-29.0)	1.0	(0.7-1.6)	3.9	(2.9-5.3)	1.1	(0.8-1.5)
	White											
	Black or African American	6	--	--	--	--	--	--	--	--	--	--
	All Other Races	66	54.5	(40.4-67.9)	31.3	(18.8-47.3)	2.7	(0.6-10.9)*	5.7	(1.3-22.1)*	5.8	(2.0-15.8)*
Education	<High School	102	60.8	(45.5-74.2)	24.9	(14.6-39.1)	2.5	(0.5-10.6)*	7.6	(3.6-15.3)*	4.2	(1.5-11.4)*
	High School Graduate (or GED)	614	63.9	(58.0-69.4)	28.2	(23.2-33.8)	0.5	(0.2-1.2)*	5.9	(3.8-9.1)	1.5	(0.5-4.0)*
	Some College	812	68.2	(63.2-72.9)	27.7	(23.3-32.6)	1.2	(0.5-2.5)*	2.3	(1.2-4.4)*	0.6	(0.2-1.7)*
	College Graduate	832	69.3	(64.7-73.6)	25.1	(21.1-29.6)	1.2	(0.5-2.9)*	3.7	(1.9-7.2)*	0.6	(0.2-1.7)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	237	60.9	(49.6-71.2)	27.6	(19.1-38.0)	1.2	(0.5-3.0)*	9.1	(4.0-19.3)*	1.2	(0.4-3.9)*
	\$20,000-\$34,999	364	72.8	(66.0-78.7)	20.6	(15.5-26.8)	1.0	(0.3-3.2)*	4.6	(2.4-8.6)*	1.0	(0.2-4.0)*
	\$35,000-\$49,999	342	64.6	(57.7-70.9)	30.5	(24.5-37.3)	0.7	(0.2-2.2)*	3.1	(1.3-7.0)*	1.2	(0.4-3.4)*
	\$50,000-\$74,999	422	62.8	(56.4-68.8)	31.4	(25.5-38.0)	0.6	(0.2-2.2)*	4.4	(2.6-7.6)	0.8	(0.2-2.8)*
	\$75,000 and more	718	73.1	(68.2-77.5)	24.1	(19.9-28.9)	0.2	(0.0-0.9)*	2.5	(1.2-4.9)*	0.1	(0.0-0.5)*
Employment Status	Employed	1,339	67.4	(63.6-70.9)	26.8	(23.5-30.4)	0.5	(0.2-1.1)*	4.1	(2.8-5.8)	1.2	(0.6-2.4)*
	Out of work	89	67.4	(49.1-81.6)	28.3	(15.2-46.4)	1.6	(0.4-6.6)*	1.6	(0.5-5.1)*	1.2	(0.2-5.2)*
	Homemaker	316	71.6	(64.6-77.8)	25.9	(19.9-32.9)	1.0	(0.1-6.8)*	1.5	(0.4-5.2)*	0.0	(0.0-0.0)
	Student	33	--	--	--	--	--	--	--	--	--	--
	Retired	499	65.9	(60.6-70.9)	25.0	(20.5-30.1)	2.8	(1.4-5.4)*	5.5	(3.3-9.1)	0.7	(0.2-2.2)*
	Unable to work	81	57.5	(44.2-69.8)	29.3	(17.7-44.4)	2.2	(0.5-9.1)*	4.7	(1.4-14.5)*	6.4	(1.7-20.9)*
Veteran Status	Yes	269	72.0	(64.6-78.4)	20.8	(15.1-27.8)	2.5	(1.0-6.0)*	4.5	(2.4-8.1)*	0.3	(0.1-1.1)*
	No	2,087	66.2	(63.1-69.2)	27.5	(24.7-30.5)	0.8	(0.5-1.5)	4.1	(2.9-5.7)	1.3	(0.7-2.3)
Diabetes	Yes	197	63.4	(55.0-71.1)	25.2	(18.4-33.6)	2.8	(0.7-10.3)*	5.6	(2.6-11.5)*	2.9	(0.6-12.6)*
	No	2,165	66.9	(63.9-69.8)	27.1	(24.4-30.1)	0.9	(0.6-1.5)	4.0	(2.9-5.5)	1.0	(0.6-1.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	197	69.9	(61.5-77.1)	18.4	(12.9-25.5)	2.4	(0.6-8.9)*	7.8	(3.9-14.9)*	1.6	(0.4-6.1)*
	No	2,140	66.6	(63.6-69.5)	27.4	(24.7-30.4)	0.9	(0.6-1.6)	3.9	(2.8-5.4)	1.1	(0.6-2.0)*
Asthma	Yes	331	69.0	(61.1-75.9)	22.5	(16.5-29.9)	1.6	(0.5-4.6)*	5.4	(2.4-11.9)*	1.5	(0.4-5.2)*
	No	2,031	66.4	(63.3-69.4)	27.7	(24.8-30.7)	0.9	(0.5-1.5)	3.9	(2.9-5.3)	1.1	(0.6-2.0)*
Arthritis	Yes	752	64.7	(59.7-69.5)	28.9	(24.4-33.9)	1.5	(0.6-3.6)*	3.9	(2.5-6.1)	0.9	(0.4-2.2)*
	No	1,599	67.4	(63.9-70.6)	26.5	(23.4-29.7)	0.9	(0.5-1.5)	4.1	(2.8-5.8)	1.3	(0.7-2.4)*
Frequent Mental Distress	Yes	215	73.1	(64.2-80.5)	19.5	(13.1-28.0)	1.2	(0.2-8.0)*	3.8	(1.8-8.0)*	2.4	(0.9-6.0)*
	No	2,120	66.0	(62.9-69.0)	27.8	(25.0-30.8)	1.0	(0.6-1.6)	4.2	(3.0-5.7)	1.0	(0.5-1.9)*
Serious Psychological Distress	Yes	63	56.4	(40.7-70.9)	33.3	(20.1-49.8)	1.9	(0.3-12.1)*	5.7	(1.6-18.6)*	2.7	(0.4-17.2)*
	No	2,258	67.1	(64.1-70.0)	26.9	(24.2-29.7)	0.9	(0.6-1.5)	4.0	(2.9-5.5)	1.1	(0.6-1.9)*
Receiving Treatment for Mental Health Condition	Yes	331	79.6	(72.0-85.6)	16.0	(10.5-23.7)	2.1	(0.6-6.7)*	1.5	(0.6-3.6)*	0.8	(0.2-2.9)*
	No	2,027	65.0	(61.8-68.0)	28.5	(25.6-31.5)	0.9	(0.5-1.5)	4.5	(3.3-6.1)	1.2	(0.7-2.2)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		2,366	23.2	(20.9-25.7)	42.7	(39.8-45.6)	0.9	(0.6-1.6)	26.2	(23.7-28.8)	7.0	(5.7-8.5)
Age (years)	18-24	95	19.1	(11.8-29.5)	50.3	(38.0-62.6)	0.0	(0.0-0.0)	26.3	(16.2-39.7)	4.2	(1.3-13.0)*
	25-34	322	15.5	(11.4-20.7)	50.6	(44.1-57.1)	1.0	(0.3-3.5)*	25.3	(20.4-31.0)	7.6	(4.8-11.8)
	35-54	905	21.4	(18.2-25.0)	40.6	(36.8-44.5)	1.4	(0.7-2.8)*	30.5	(27.1-34.1)	6.2	(4.7-8.0)
	55+	1,035	33.1	(29.7-36.8)	34.6	(31.3-38.1)	1.0	(0.5-1.9)*	22.9	(20.0-26.0)	8.4	(6.6-10.7)
Sex	Men	1,022	25.8	(22.3-29.6)	43.7	(39.4-48.2)	0.8	(0.4-1.6)*	23.8	(20.2-27.8)	5.9	(4.1-8.4)
	Women	1,344	20.8	(17.9-24.0)	41.8	(38.0-45.6)	1.1	(0.5-2.2)*	28.4	(25.2-31.7)	8.0	(6.3-10.0)
Race/ Ethnicity	Hispanic or Latino	117	38.8	(28.4-50.4)	32.6	(21.9-45.5)	1.2	(0.3-4.3)*	18.8	(11.1-30.1)	8.6	(4.1-17.0)*
	Not Hispanic or Latino:	2,167	20.9	(18.7-23.4)	42.8	(39.7-45.9)	1.3	(0.9-1.9)	27.7	(25.0-30.5)	7.3	(6.0-8.9)
	White											
	Black or African American	5	--	--	--	--	--	--	--	--	--	--
	All Other Races	66	21.4	(10.3-39.4)*	41.8	(27.1-58.0)	2.5	(0.5-11.6)*	19.5	(10.1-34.5)*	14.8	(7.5-27.1)*
Education	<High School	101	32.3	(21.4-45.6)	38.1	(25.8-52.3)	0.0	(0.0-0.0)	20.4	(11.1-34.4)	9.2	(3.6-21.3)*
	High School Graduate (or GED)	609	27.9	(23.0-33.4)	42.3	(36.2-48.5)	0.4	(0.2-1.1)*	22.8	(18.2-28.1)	6.7	(4.3-10.1)
	Some College	820	19.8	(16.2-23.9)	42.2	(37.3-47.2)	1.6	(0.7-3.4)*	28.6	(24.2-33.4)	8.0	(5.7-11.1)
	College Graduate	833	19.7	(16.4-23.4)	44.3	(39.9-48.9)	1.0	(0.4-2.1)*	28.5	(24.6-32.6)	6.6	(4.9-8.8)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	
Annual Household Income	< \$20,000	237	19.6	(14.5-25.9)	45.8	(35.7-56.3)	0.1	(0.0-0.9)*	26.8	(18.4-37.3)	7.7	(4.7-12.3)
	\$20,000-\$34,999	363	25.1	(19.7-31.5)	44.8	(37.3-52.5)	0.5	(0.1-1.9)*	22.2	(17.2-28.2)	7.5	(4.6-11.8)
	\$35,000-\$49,999	342	21.6	(17.0-27.1)	45.3	(38.5-52.2)	0.8	(0.2-3.0)*	23.3	(17.9-29.6)	9.1	(5.7-14.0)
	\$50,000-\$74,999	423	21.1	(16.5-26.7)	42.0	(35.6-48.6)	0.8	(0.2-3.1)*	31.0	(25.2-37.5)	5.1	(3.1-8.3)
	\$75,000 and more	718	22.8	(18.4-27.8)	42.8	(37.8-47.9)	1.0	(0.5-2.4)*	27.4	(23.4-31.7)	6.0	(3.8-9.3)
Employment Status	Employed	1,343	22.4	(19.3-25.7)	43.6	(39.9-47.4)	1.0	(0.5-2.1)*	26.7	(23.5-30.1)	6.3	(4.8-8.3)
	Out of work	94	18.7	(10.7-30.8)	37.6	(23.2-54.5)	0.9	(0.1-6.5)*	27.4	(15.7-43.2)	15.3	(6.7-31.4)*
	Homemaker	320	25.7	(20.0-32.3)	41.9	(35.1-49.1)	0.6	(0.1-2.5)*	24.4	(19.1-30.5)	7.5	(4.6-12.0)
	Student	33	--	--	--	--	--	--	--	--	--	--
	Retired	489	32.8	(27.9-38.1)	41.1	(35.7-46.7)	0.9	(0.4-2.0)*	20.8	(16.8-25.4)	4.5	(2.9-6.9)
	Unable to work	82	14.7	(6.4-30.2)*	28.8	(17.9-42.9)	2.0	(0.4-8.8)*	41.0	(28.7-54.6)	13.5	(7.7-22.5)
Veteran Status	Yes	266	30.0	(23.7-37.2)	34.5	(27.2-42.6)	1.2	(0.4-3.4)*	26.9	(20.5-34.5)	7.4	(4.2-12.5)
	No	2,093	22.5	(20.0-25.1)	43.4	(40.3-46.6)	0.9	(0.5-1.6)	26.2	(23.6-29.0)	7.0	(5.6-8.6)
Diabetes	Yes	202	18.1	(13.1-24.5)	41.4	(33.3-50.0)	0.2	(0.0-1.1)*	27.9	(20.7-36.4)	12.5	(7.8-19.4)
	No	2,163	23.6	(21.1-26.2)	42.7	(39.7-45.8)	1.0	(0.6-1.7)	26.1	(23.6-28.8)	6.6	(5.3-8.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Utah, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	200	27.8	(21.1-35.6)	37.3	(29.4-46.1)	1.1	(0.3-3.4)*	22.2	(16.1-29.8)	11.6	(7.2-18.1)
	No	2,142	22.9	(20.5-25.6)	43.2	(40.2-46.3)	0.9	(0.5-1.6)	26.2	(23.6-28.9)	6.7	(5.4-8.3)
Asthma	Yes	332	13.6	(9.2-19.6)	51.4	(43.4-59.2)	0.4	(0.1-1.9)*	24.1	(18.4-31.0)	10.5	(6.8-15.8)
	No	2,033	24.8	(22.3-27.6)	41.1	(38.1-44.3)	1.0	(0.6-1.8)	26.6	(23.9-29.4)	6.4	(5.1-8.0)
Arthritis	Yes	751	21.0	(17.7-24.6)	38.3	(33.7-43.1)	0.5	(0.2-1.0)*	29.4	(25.4-33.7)	10.9	(8.5-14.0)
	No	1,601	24.0	(21.2-27.1)	43.9	(40.4-47.5)	0.9	(0.5-1.8)*	25.3	(22.5-28.4)	5.8	(4.4-7.6)
Frequent Mental Distress	Yes	217	20.7	(14.5-28.6)	34.9	(26.4-44.5)	0.5	(0.1-3.5)*	28.3	(21.3-36.6)	15.6	(9.5-24.5)
	No	2,121	23.7	(21.2-26.3)	43.6	(40.5-46.7)	1.0	(0.6-1.7)	25.8	(23.3-28.6)	5.9	(4.8-7.3)
Serious Psychological Distress	Yes	69	18.5	(9.0-34.3)*	29.9	(18.0-45.2)	0.0	(0.0-0.0)	31.5	(19.5-46.6)	20.2	(11.5-32.9)
	No	2,258	23.4	(21.1-26.0)	42.9	(40.0-46.0)	1.0	(0.6-1.6)	26.1	(23.6-28.8)	6.5	(5.3-8.1)
Receiving Treatment for Mental Health Condition	Yes	333	15.5	(11.0-21.6)	37.0	(29.5-45.1)	1.8	(0.6-4.9)*	32.3	(25.9-39.5)	13.4	(9.3-18.9)
	No	2,028	24.5	(22.0-27.2)	42.8	(39.7-45.9)	1.1	(0.7-1.7)	25.1	(22.5-27.9)	6.5	(5.2-8.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,096	69.0	(67.2-70.6)	25.2	(23.6-26.9)	1.5	(1.1-2.1)	3.1	(2.4-4.0)	1.2	(0.8-1.7)
Age (years)	18-24	157	53.7	(45.5-61.7)	39.1	(31.3-47.5)	1.2	(0.4-3.8)*	5.3	(2.6-10.6)*	0.6	(0.1-4.0)*
	25-34	431	63.6	(58.6-68.3)	30.1	(25.6-35.1)	1.4	(0.5-3.9)*	2.8	(1.4-5.2)*	2.1	(0.8-5.3)*
	35-54	2,204	71.3	(69.0-73.4)	22.9	(20.9-25.1)	1.7	(1.1-2.7)	3.0	(2.2-4.1)	1.1	(0.6-1.9)
	55+	3,269	74.0	(72.2-75.7)	20.6	(19.0-22.3)	1.6	(1.1-2.2)	2.5	(1.9-3.3)	1.3	(0.9-1.9)
Sex	Men	2,410	62.1	(59.5-64.7)	30.7	(28.2-33.3)	1.9	(1.3-2.8)	3.6	(2.6-4.8)	1.7	(1.0-2.8)
	Women	3,686	75.3	(73.3-77.3)	20.1	(18.2-22.0)	1.2	(0.8-1.8)	2.8	(1.9-4.0)	0.7	(0.4-1.1)
Race/ Ethnicity	Hispanic or Latino	68	53.5	(40.8-65.8)	43.5	(31.0-56.8)	0.5	(0.3-0.8)	1.3	(0.8-2.0)	1.2	(0.2-8.0)*
	Not Hispanic or Latino:	5,798	69.1	(67.5-70.7)	25.2	(23.7-26.9)	1.4	(1.1-1.9)	3.1	(2.5-3.8)	1.1	(0.8-1.4)
	White											
	Black or African American	19	--	--	--	--	--	--	--	--	--	--
	All Other Races	143	70.1	(59.8-78.7)	21.6	(14.0-31.9)	1.1	(0.3-3.3)*	3.0	(1.2-6.9)*	4.2	(2.0-8.7)*
Education	<High School	318	56.8	(49.0-64.2)	34.3	(27.1-42.3)	1.8	(0.6-5.7)*	3.8	(2.2-6.5)	3.3	(1.2-8.9)*
	High School Graduate (or GED)	1,820	61.9	(58.7-65.0)	30.0	(27.0-33.2)	2.0	(1.3-3.2)	4.4	(3.0-6.4)	1.7	(1.0-2.9)
	Some College	1,376	70.3	(66.6-73.7)	24.0	(20.8-27.5)	1.3	(0.7-2.2)	3.6	(2.3-5.8)	0.9	(0.4-1.8)*
	College Graduate	2,576	76.4	(74.1-78.6)	20.4	(18.3-22.7)	1.3	(0.7-2.3)*	1.5	(1.0-2.3)	0.4	(0.2-0.8)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	852	59.8 (54.4-65.0)	28.9 (24.1-34.3)	1.8 (0.8-4.3)*	6.3 (3.9-9.8)	3.2 (1.7-5.8)*				
	\$20,000-\$34,999	1,086	66.8 (62.8-70.5)	28.0 (24.3-32.0)	1.3 (0.7-2.4)*	2.8 (1.9-4.3)	1.1 (0.4-3.3)*				
	\$35,000-\$49,999	900	70.6 (66.5-74.3)	24.5 (20.9-28.4)	2.2 (1.1-4.4)*	2.5 (1.3-4.4)*	0.4 (0.1-1.2)*				
	\$50,000-\$74,999	1,026	74.2 (70.6-77.5)	22.0 (18.9-25.5)	1.6 (0.9-2.9)*	1.5 (0.8-2.8)*	0.7 (0.3-1.6)*				
	\$75,000 and more	1,595	74.9 (71.9-77.6)	21.7 (19.0-24.6)	0.9 (0.5-1.6)	2.1 (1.2-3.7)	0.4 (0.1-1.1)*				
Employment Status	Employed	3,685	69.4 (67.3-71.5)	25.8 (23.8-28.0)	1.5 (1.0-2.3)	2.5 (1.8-3.6)	0.7 (0.4-1.2)				
	Out of work	312	66.4 (59.4-72.7)	22.2 (16.6-29.0)	2.7 (1.0-7.0)*	6.1 (3.3-11.2)*	2.6 (0.9-7.7)*				
	Homemaker	291	66.2 (58.6-73.1)	26.6 (20.0-34.5)	2.6 (0.9-6.9)*	4.3 (1.5-11.4)*	0.3 (0.0-2.2)*				
	Student	52	74.4 (58.9-85.5)	19.7 (10.2-34.4)*	0.0 (0.0-0.0)	6.0 (1.5-21.0)*	0.0 (0.0-0.0)				
	Retired	1,417	69.5 (66.5-72.4)	25.3 (22.6-28.3)	1.4 (0.8-2.4)	2.9 (2.0-4.3)	0.8 (0.5-1.4)				
	Unable to work	313	65.4 (58.9-71.4)	25.9 (20.3-32.4)	1.1 (0.4-3.1)*	3.3 (1.9-5.9)	4.2 (1.9-9.0)*				
Veteran Status	Yes	782	66.2 (62.2-70.0)	26.9 (23.3-30.8)	1.6 (0.7-4.0)*	3.2 (2.0-5.0)	2.0 (1.1-3.7)*				
	No	5,298	69.3 (67.4-71.1)	25.1 (23.4-27.0)	1.5 (1.1-2.1)	3.1 (2.3-4.1)	1.0 (0.6-1.6)				
Diabetes	Yes	529	71.8 (67.0-76.1)	21.7 (17.6-26.3)	1.5 (0.7-3.0)*	3.8 (2.3-6.1)	1.3 (0.6-2.6)*				
	No	5,564	68.7 (66.9-70.5)	25.5 (23.8-27.3)	1.5 (1.1-2.2)	3.1 (2.4-4.0)	1.1 (0.7-1.7)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	553	68.6	(63.8-73.1)	22.9	(18.7-27.6)	2.7	(1.3-5.3)*	4.2	(2.6-6.6)	1.7	(0.8-3.4)*
	No	5,503	69.1	(67.3-70.9)	25.3	(23.6-27.1)	1.4	(1.0-2.0)	3.1	(2.3-4.0)	1.1	(0.7-1.7)
Asthma	Yes	855	68.1	(63.7-72.3)	23.8	(19.8-28.2)	1.6	(0.9-3.0)*	4.1	(2.5-6.5)	2.4	(1.0-5.6)*
	No	5,221	69.2	(67.4-71.0)	25.4	(23.6-27.2)	1.5	(1.1-2.1)	3.0	(2.2-4.0)	0.9	(0.6-1.3)
Arthritis	Yes	2,274	70.2	(67.7-72.6)	23.2	(20.8-25.7)	1.8	(1.1-2.8)	3.4	(2.5-4.7)	1.4	(0.8-2.5)*
	No	3,792	68.5	(66.4-70.6)	25.9	(23.9-28.1)	1.4	(1.0-2.1)	3.0	(2.2-4.2)	1.0	(0.6-1.7)
Frequent Mental Distress	Yes	556	61.6	(56.3-66.6)	27.7	(22.8-33.2)	1.7	(0.9-3.3)*	5.4	(3.2-9.0)	3.6	(1.7-7.4)*
	No	5,466	69.9	(68.1-71.7)	25.1	(23.4-26.8)	1.5	(1.1-2.2)	2.6	(2.0-3.4)	0.9	(0.6-1.3)
Serious Psychological Distress	Yes	186	53.3	(43.5-62.8)	31.0	(22.0-41.7)	2.1	(0.8-5.7)*	7.3	(4.3-12.1)	6.3	(2.4-15.5)*
	No	5,826	69.8	(68.0-71.5)	25.0	(23.4-26.8)	1.4	(1.0-1.9)	2.9	(2.2-3.8)	0.9	(0.6-1.3)
Receiving Treatment for Mental Health Condition	Yes	899	75.5	(71.1-79.5)	17.8	(14.1-22.2)	0.7	(0.3-1.7)*	3.6	(1.9-6.8)*	2.4	(1.0-5.8)*
	No	5,179	68.0	(66.1-69.8)	26.3	(24.6-28.2)	1.7	(1.2-2.4)	3.1	(2.4-4.0)	0.9	(0.6-1.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,094	24.5	(23.0-26.1)	38.9	(37.2-40.6)	2.7	(2.2-3.4)	25.5	(24.1-27.0)	8.3	(7.4-9.4)
Age (years)												
	18-24	157	24.1	(17.6-32.0)	41.5	(33.5-50.0)	3.8	(1.6-8.8)*	24.7	(18.2-32.6)	5.9	(2.6-12.8)*
	25-34	433	18.2	(14.4-22.8)	46.0	(40.9-51.1)	1.6	(0.8-3.5)*	27.3	(23.1-31.9)	6.9	(4.8-9.9)
	35-54	2,205	21.1	(19.1-23.2)	39.8	(37.5-42.1)	2.3	(1.7-3.2)	27.4	(25.5-29.5)	9.3	(8.1-10.8)
	55+	3,264	31.6	(29.8-33.5)	34.0	(32.1-35.8)	3.2	(2.6-4.1)	22.4	(20.8-24.0)	8.8	(7.8-9.9)
Sex												
	Men	2,428	25.3	(23.1-27.8)	42.4	(39.7-45.0)	2.7	(1.9-3.7)	22.5	(20.5-24.7)	7.1	(5.8-8.7)
	Women	3,666	23.8	(21.9-25.8)	35.6	(33.5-37.7)	2.8	(2.0-3.8)	28.4	(26.5-30.4)	9.5	(8.3-10.8)
Race/Ethnicity												
	Hispanic or Latino	68	34.1	(21.1-49.9)	23.5	(13.5-37.8)	0.8	(0.5-1.1)	33.6	(18.7-52.8)	8.0	(3.3-18.2)*
	Not Hispanic or Latino:	5,797	23.0	(21.6-24.5)	39.0	(37.3-40.7)	2.8	(2.3-3.5)	26.7	(25.2-28.2)	8.5	(7.6-9.5)
	White											
	Black or African American	19	--	--	--	--	--	--	--	--	--	--
	All Other Races	141	20.4	(13.1-30.3)	42.4	(32.1-53.5)	1.2	(0.4-3.1)*	22.3	(15.1-31.7)	13.7	(9.0-20.3)
Education												
	<High School	325	32.1	(26.1-38.7)	41.0	(33.4-49.1)	0.9	(0.4-2.4)*	18.8	(13.7-25.2)	7.2	(4.5-11.3)
	High School Graduate (or GED)	1,835	32.4	(29.4-35.6)	33.0	(30.0-36.2)	2.8	(1.7-4.4)	23.5	(20.9-26.4)	8.2	(6.6-10.3)
	Some College	1,376	22.0	(19.0-25.3)	38.8	(35.2-42.6)	3.2	(2.1-4.7)	25.8	(22.9-29.1)	10.2	(8.0-12.9)
	College Graduate	2,553	17.3	(15.4-19.4)	43.2	(40.8-45.7)	2.9	(2.2-3.9)	28.7	(26.5-30.9)	7.8	(6.7-9.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	865	29.5	(24.6-35.0)	32.0	(27.1-37.2)	1.8	(1.1-3.1)	25.7	(21.5-30.4)	11.0	(8.9-13.6)
	\$20,000-\$34,999	1,091	24.2	(21.1-27.6)	36.6	(32.7-40.6)	3.2	(1.8-5.7)	24.5	(20.9-28.4)	11.5	(9.1-14.5)
	\$35,000-\$49,999	903	23.7	(20.3-27.4)	40.7	(36.6-44.9)	1.6	(1.0-2.7)	26.8	(23.3-30.6)	7.2	(5.6-9.3)
	\$50,000-\$74,999	1,024	22.7	(19.5-26.2)	41.2	(37.5-45.0)	3.1	(1.9-4.8)	26.3	(23.2-29.6)	6.7	(5.2-8.7)
	\$75,000 and more	1,586	22.4	(19.7-25.3)	41.7	(38.7-44.7)	3.3	(2.2-4.7)	25.7	(23.3-28.2)	7.0	(5.6-8.7)
Employment Status	Employed	3,688	23.2	(21.3-25.3)	40.1	(37.9-42.3)	2.8	(2.1-3.8)	26.1	(24.3-28.0)	7.8	(6.6-9.0)
	Out of work	308	25.1	(18.9-32.5)	35.3	(28.3-43.0)	1.8	(0.7-5.0)*	23.7	(17.6-31.0)	14.1	(10.0-19.4)
	Homemaker	284	28.2	(22.5-34.7)	28.8	(22.6-35.9)	4.9	(2.6-9.0)*	27.2	(21.9-33.3)	10.9	(6.1-18.7)
	Student	53	27.6	(14.5-46.1)	49.1	(33.0-65.4)	3.2	(0.4-20.2)*	16.6	(8.0-31.3)*	3.5	(0.8-14.9)*
	Retired	1,414	30.2	(27.5-33.1)	39.3	(36.2-42.5)	2.6	(1.8-3.8)	21.2	(18.8-23.8)	6.7	(5.3-8.3)
	Unable to work	319	18.7	(14.6-23.7)	30.1	(24.0-36.9)	2.3	(1.1-4.7)*	35.1	(28.5-42.2)	13.8	(10.3-18.3)
Veteran Status	Yes	787	21.8	(18.8-25.2)	38.9	(34.8-43.2)	3.7	(2.5-5.4)	25.5	(21.6-29.9)	10.1	(7.8-12.9)
	No	5,291	24.9	(23.2-26.6)	38.8	(36.9-40.7)	2.6	(2.0-3.4)	25.6	(24.0-27.2)	8.2	(7.2-9.3)
Diabetes	Yes	530	26.3	(22.5-30.5)	31.1	(26.7-36.0)	2.0	(1.1-3.9)*	29.7	(25.1-34.6)	10.9	(8.2-14.3)
	No	5,561	24.4	(22.8-26.1)	39.3	(37.5-41.2)	2.8	(2.2-3.5)	25.3	(23.8-26.9)	8.2	(7.2-9.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Vermont, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	553	26.3	(22.4-30.7)	34.7	(30.2-39.6)	2.2	(1.2-3.9)	24.3	(20.0-29.1)	12.5	(9.5-16.2)
	No	5,501	24.4	(22.8-26.1)	39.1	(37.3-41.0)	2.8	(2.2-3.5)	25.6	(24.1-27.2)	8.1	(7.1-9.1)
Asthma	Yes	852	20.1	(16.9-23.7)	39.4	(34.9-44.0)	2.7	(1.5-4.8)	28.2	(24.4-32.4)	9.6	(6.7-13.4)
	No	5,222	25.3	(23.6-27.1)	38.7	(36.9-40.6)	2.8	(2.2-3.5)	25.1	(23.5-26.7)	8.1	(7.2-9.1)
Arthritis	Yes	2,267	24.6	(22.5-26.7)	35.5	(33.0-38.1)	2.3	(1.7-3.1)	26.7	(24.4-29.1)	11.0	(9.2-13.1)
	No	3,798	24.4	(22.5-26.5)	40.1	(37.9-42.3)	2.9	(2.2-3.9)	25.2	(23.5-27.1)	7.4	(6.3-8.5)
Frequent Mental Distress	Yes	556	61.6	(56.3-66.6)	27.7	(22.8-33.2)	1.7	(0.9-3.3)*	5.4	(3.2-9.0)	3.6	(1.7-7.4)*
	No	5,466	69.9	(68.1-71.7)	25.1	(23.4-26.8)	1.5	(1.1-2.2)	2.6	(2.0-3.4)	0.9	(0.6-1.3)
Serious Psychological Distress	Yes	186	53.3	(43.5-62.8)	31.0	(22.0-41.7)	2.1	(0.8-5.7)*	7.3	(4.3-12.1)	6.3	(2.4-15.5)*
	No	5,826	69.8	(68.0-71.5)	25.0	(23.4-26.8)	1.4	(1.0-1.9)	2.9	(2.2-3.8)	0.9	(0.6-1.3)
Receiving Treatment for Mental Health Condition	Yes	899	75.5	(71.1-79.5)	17.8	(14.1-22.2)	0.7	(0.3-1.7)*	3.6	(1.9-6.8)*	2.4	(1.0-5.8)*
	No	5,179	68.0	(66.1-69.8)	26.3	(24.6-28.2)	1.7	(1.2-2.4)	3.1	(2.4-4.0)	0.9	(0.6-1.3)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,159	70.3	(67.9-72.5)	22.0	(20.2-24.0)	1.9	(1.4-2.5)	3.2	(2.5-4.0)	2.7	(1.4-5.1)*
Age (years)	18-24	148	62.0	(51.2-71.8)	27.7	(19.3-38.0)	0.8	(0.2-2.5)*	3.7	(1.5-9.0)*	5.8	(2.5-12.9)*
	25-34	504	65.1	(57.3-72.2)	25.1	(19.8-31.2)	1.7	(0.7-4.0)*	2.4	(1.0-5.6)*	5.7	(1.5-19.2)*
	35-54	1,941	74.3	(71.4-77.0)	19.7	(17.3-22.3)	2.3	(1.5-3.5)	2.6	(1.8-3.9)	1.2	(0.5-2.5)*
	55+	2,512	72.5	(70.1-74.8)	20.9	(18.8-23.2)	1.9	(1.3-2.7)	3.6	(2.8-4.7)	1.1	(0.7-1.7)
Sex	Men	1,887	65.7	(61.4-69.7)	23.9	(20.7-27.3)	2.6	(1.8-3.7)	3.2	(2.2-4.6)	4.7	(2.2-9.9)*
	Women	3,272	74.5	(72.3-76.6)	20.3	(18.4-22.4)	1.3	(0.9-1.9)	3.1	(2.3-4.2)	0.8	(0.5-1.4)
Race/ Ethnicity	Hispanic or Latino	155	66.4	(54.7-76.4)	22.9	(14.7-33.8)	1.0	(0.3-3.5)*	1.4	(0.5-4.1)*	8.4	(3.2-20.0)*
	Not Hispanic or Latino:	4,201	72.1	(69.8-74.3)	21.8	(19.8-23.8)	1.8	(1.3-2.5)	3.3	(2.5-4.3)	1.1	(0.7-1.7)
	White											
	Black or African American	543	69.6	(63.6-75.1)	22.2	(17.4-27.8)	2.9	(1.6-5.4)*	3.4	(2.1-5.5)	1.9	(0.9-3.9)*
	All Other Races	189	56.8	(40.3-72.0)	16.8	(7.9-32.0)*	1.8	(0.5-5.6)*	3.8	(1.7-8.0)*	20.9	(7.9-44.7)*
Education	<High School	479	69.1	(62.0-75.4)	20.5	(15.5-26.6)	0.5	(0.2-1.0)*	4.6	(3.0-7.0)	5.3	(2.9-9.5)*
	High School Graduate (or GED)	1,342	64.4	(60.2-68.4)	25.2	(21.7-28.9)	2.7	(1.8-4.0)	4.8	(3.2-7.0)	3.0	(1.6-5.6)*
	Some College	1,184	70.3	(65.9-74.4)	24.7	(20.8-29.0)	1.8	(1.1-2.9)	2.4	(1.5-3.9)	0.8	(0.3-1.9)*
	College Graduate	2,143	75.0	(70.7-78.8)	18.1	(15.4-21.2)	1.6	(0.9-2.7)	2.0	(1.1-3.6)	3.3	(0.9-10.9)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	606	70.2 (62.5-76.8)	22.1 (16.7-28.7)	1.3 (0.6-2.6)*	5.0 (2.7-8.9)*	1.5 (0.6-3.8)*				
	\$20,000-\$34,999	784	68.5 (62.8-73.7)	22.3 (17.7-27.8)	2.0 (0.9-4.1)*	4.8 (3.0-7.8)	2.4 (0.8-6.8)*				
	\$35,000-\$49,999	679	70.5 (64.9-75.5)	22.2 (17.9-27.3)	2.0 (0.9-4.5)*	2.7 (1.5-4.7)	2.6 (1.0-6.5)*				
	\$50,000-\$74,999	839	72.0 (67.3-76.2)	21.7 (18.0-26.0)	1.6 (0.8-3.2)*	3.9 (2.0-7.7)*	0.8 (0.3-2.0)*				
	\$75,000 and more	1,670	75.2 (70.7-79.3)	18.4 (15.4-21.9)	1.5 (0.8-2.6)	1.1 (0.7-1.9)	3.7 (1.2-11.0)*				
Employment Status	Employed	2,802	70.5 (67.2-73.5)	21.4 (19.0-24.1)	2.1 (1.5-3.0)	2.6 (1.8-3.9)	3.3 (1.5-7.3)*				
	Out of work	157	71.7 (60.8-80.5)	20.5 (13.2-30.6)	3.8 (1.3-10.7)*	1.8 (0.4-6.8)*	2.2 (0.6-8.4)*				
	Homemaker	445	69.6 (64.0-74.8)	23.2 (18.6-28.6)	1.0 (0.4-2.9)*	5.3 (2.4-11.5)*	0.8 (0.2-3.4)*				
	Student	90	83.5 (71.9-90.9)	12.5 (6.3-23.2)*	0.0 (0.0-0.0)	2.1 (0.6-6.8)*	1.9 (0.4-8.4)*				
	Retired	1,344	65.4 (61.7-68.9)	26.8 (23.4-30.5)	1.7 (1.0-2.8)	4.5 (3.2-6.3)	1.6 (0.9-2.9)*				
	Unable to work	302	66.0 (58.9-72.4)	27.0 (20.8-34.1)	1.5 (0.6-3.5)*	4.8 (2.9-7.8)	0.8 (0.3-2.4)*				
Veteran Status	Yes	887	67.5 (62.7-71.9)	25.1 (21.0-29.6)	2.6 (1.5-4.2)	2.9 (1.9-4.5)	2.0 (0.9-4.5)*				
	No	4,272	70.8 (68.1-73.3)	21.4 (19.4-23.7)	1.7 (1.3-2.4)	3.2 (2.5-4.2)	2.8 (1.3-5.9)*				
Diabetes	Yes	603	67.0 (61.7-71.9)	23.3 (19.0-28.1)	1.6 (0.8-3.0)*	5.2 (3.5-7.7)	2.9 (0.8-9.7)*				
	No	4,550	70.5 (68.0-72.9)	21.9 (20.0-24.1)	1.9 (1.4-2.6)	3.0 (2.3-3.9)	2.7 (1.3-5.4)*				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	553	67.1	(61.2-72.6)	24.4	(19.4-30.1)	1.8	(0.8-3.6)*	5.7	(3.7-8.8)	1.0	(0.4-2.4)*
	No	4,571	70.5	(68.0-72.9)	21.8	(19.8-23.9)	1.9	(1.4-2.6)	2.9	(2.2-3.9)	2.8	(1.4-5.5)*
Asthma	Yes	662	71.7	(65.2-77.5)	22.4	(17.3-28.5)	1.7	(0.9-3.2)*	3.3	(2.0-5.5)	0.9	(0.4-2.0)*
	No	4,487	69.9	(67.4-72.3)	21.9	(20.0-24.0)	1.9	(1.4-2.6)	3.2	(2.4-4.1)	3.1	(1.6-6.0)*
Arthritis	Yes	1,906	70.1	(67.2-73.0)	23.1	(20.4-26.0)	1.7	(1.1-2.6)	4.1	(2.9-5.8)	1.0	(0.6-1.6)
	No	3,225	70.3	(67.3-73.2)	21.6	(19.3-24.1)	2.0	(1.4-2.8)	2.8	(2.1-3.9)	3.3	(1.6-6.6)*
Frequent Mental Distress	Yes	450	67.4	(59.6-74.4)	24.2	(17.8-32.1)	1.9	(1.0-3.7)*	3.4	(1.8-6.4)*	3.1	(1.0-8.8)*
	No	4,647	70.6	(68.1-73.0)	21.8	(19.9-23.9)	1.8	(1.4-2.5)	3.1	(2.4-4.1)	2.6	(1.2-5.4)*
Serious Psychological Distress	Yes	180	54.9	(44.9-64.6)	34.6	(25.6-44.8)	1.6	(0.5-5.2)*	4.3	(1.7-10.9)*	4.6	(2.0-9.9)*
	No	4,905	71.0	(68.5-73.3)	21.5	(19.6-23.5)	1.9	(1.5-2.6)	3.0	(2.3-3.9)	2.6	(1.3-5.2)*
Receiving Treatment for Mental Health Condition	Yes	698	72.9	(67.9-77.3)	22.1	(17.8-27.0)	0.7	(0.3-2.0)*	3.3	(2.1-5.2)	1.1	(0.5-2.2)*
	No	4,451	70.0	(67.4-72.4)	22.0	(20.0-24.2)	2.0	(1.5-2.7)	3.1	(2.4-4.1)	2.8	(1.4-5.6)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,130	21.5	(19.5-23.6)	35.9	(33.6-38.2)	4.2	(3.5-5.0)	27.3	(25.2-29.4)	11.2	(10.0-12.5)
Age (years)												
	18-24	146	8.9	(4.7-16.3)*	48.2	(37.3-59.4)	6.1	(3.1-11.6)*	28.6	(19.6-39.7)	8.2	(3.9-16.4)*
	25-34	504	21.5	(15.3-29.3)	32.5	(26.3-39.5)	4.2	(2.7-6.7)	29.6	(23.5-36.4)	12.2	(9.1-16.2)
	35-54	1,933	20.0	(17.3-23.0)	35.7	(32.6-38.9)	4.3	(3.2-5.7)	28.9	(25.9-32.0)	11.2	(9.5-13.1)
	55+	2,491	27.6	(25.3-30.0)	34.1	(31.4-36.9)	3.6	(2.9-4.5)	23.6	(21.3-26.1)	11.1	(9.6-12.8)
Sex												
	Men	1,865	23.7	(20.2-27.6)	37.8	(33.9-41.8)	4.0	(3.0-5.4)	25.6	(22.2-29.3)	8.9	(7.1-11.1)
	Women	3,265	19.4	(17.5-21.4)	34.1	(31.8-36.5)	4.3	(3.4-5.4)	28.9	(26.6-31.2)	13.3	(11.9-14.9)
Race/Ethnicity												
	Hispanic or Latino	150	42.0	(29.8-55.2)	25.6	(17.4-35.9)	4.9	(1.9-12.3)*	23.9	(14.0-37.6)	3.7	(1.6-8.1)*
	Not Hispanic or Latino:	4,176	17.0	(15.4-18.8)	37.1	(34.7-39.6)	4.5	(3.7-5.5)	29.4	(27.0-31.9)	11.9	(10.6-13.5)
	White											
	Black or African American	538	24.2	(19.5-29.7)	32.9	(26.5-40.0)	2.8	(1.5-5.0)*	26.8	(21.2-33.3)	13.2	(9.9-17.5)
	All Other Races	193	37.1	(20.7-57.3)	36.6	(23.3-52.4)	2.8	(1.1-7.4)*	13.6	(8.2-21.8)	9.8	(5.1-18.2)*
Education												
	<High School	480	36.3	(30.0-43.0)	31.9	(24.4-40.4)	4.3	(1.9-9.2)*	17.4	(13.1-22.9)	10.1	(6.4-15.6)
	High School Graduate (or GED)	1,333	24.5	(21.5-27.8)	40.1	(36.0-44.2)	3.6	(2.4-5.6)	21.0	(18.0-24.4)	10.7	(8.8-13.1)
	Some College	1,183	17.0	(14.0-20.4)	35.0	(30.7-39.6)	4.2	(3.0-5.8)	31.8	(27.4-36.6)	12.0	(9.5-15.0)
	College Graduate	2,122	16.8	(13.3-20.9)	35.7	(32.0-39.5)	4.5	(3.4-5.9)	31.3	(27.8-35.1)	11.7	(9.8-13.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	604	24.8	(20.2-30.0)	31.7	(24.3-40.2)	2.2	(1.1-4.3)*	25.3	(19.3-32.5)	16.0	(11.6-21.7)
	\$20,000-\$34,999	786	24.6	(19.9-30.0)	34.6	(29.1-40.4)	2.5	(1.6-4.0)	28.8	(23.4-35.0)	9.5	(7.4-12.1)
	\$35,000-\$49,999	679	23.2	(18.8-28.2)	37.8	(32.1-43.8)	4.1	(2.5-6.6)	23.5	(19.4-28.1)	11.5	(8.7-15.0)
	\$50,000-\$74,999	840	21.1	(17.6-25.0)	35.2	(30.9-39.7)	3.3	(2.0-5.3)	30.3	(26.0-35.0)	10.1	(7.8-13.0)
	\$75,000 and more	1,652	17.6	(13.7-22.5)	39.2	(35.0-43.6)	4.7	(3.5-6.3)	28.7	(24.9-32.8)	9.7	(7.9-11.9)
Employment Status	Employed	2,791	21.3	(18.3-24.5)	36.5	(33.5-39.7)	4.1	(3.2-5.3)	27.9	(25.1-30.9)	10.2	(8.8-11.8)
	Out of work	155	28.2	(20.4-37.4)	26.0	(18.1-35.9)	6.6	(2.7-15.5)*	19.6	(12.7-29.0)	19.6	(11.8-30.8)
	Homemaker	439	21.1	(16.9-26.0)	34.9	(28.9-41.5)	7.2	(4.6-11.1)	24.8	(19.9-30.5)	12.0	(8.8-16.2)
	Student	90	5.7	(2.8-11.4)*	50.7	(36.1-65.2)	4.2	(1.2-14.2)*	26.7	(16.1-40.8)	12.7	(5.1-28.1)*
	Retired	1,335	25.8	(22.9-28.9)	33.3	(29.8-37.1)	3.6	(2.6-4.9)	26.8	(23.6-30.2)	10.5	(8.3-13.2)
	Unable to work	301	20.3	(15.6-25.9)	30.3	(23.7-37.8)	1.3	(0.5-3.4)*	32.2	(24.6-40.8)	16.0	(12.0-21.1)
Veteran Status	Yes	878	18.7	(15.6-22.3)	40.6	(35.6-45.8)	3.9	(2.6-5.7)	25.6	(21.4-30.3)	11.2	(8.2-15.2)
	No	4,252	22.1	(19.8-24.6)	34.9	(32.4-37.5)	4.2	(3.4-5.2)	27.6	(25.2-30.0)	11.2	(10.0-12.6)
Diabetes	Yes	595	23.3	(19.4-27.6)	29.6	(24.8-34.9)	2.5	(1.5-4.2)	33.2	(27.6-39.2)	11.4	(8.7-14.8)
	No	4,529	21.3	(19.1-23.6)	36.4	(34.0-38.9)	4.3	(3.6-5.2)	26.8	(24.6-29.1)	11.2	(9.9-12.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Virginia, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

			Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics		Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	551	20.7	(16.9-25.0)	36.3	(29.9-43.3)	2.7	(1.5-4.8)	26.3	(21.3-32.0)	14.0	(10.6-18.3)
	No	4,544	21.5	(19.4-23.9)	36.0	(33.6-38.4)	4.3	(3.5-5.2)	27.3	(25.1-29.6)	10.9	(9.6-12.3)
Asthma	Yes	659	16.3	(12.7-20.7)	36.7	(29.9-44.1)	4.4	(2.3-8.1)*	29.2	(22.2-37.3)	13.4	(10.4-17.1)
	No	4,461	22.3	(20.1-24.6)	35.8	(33.4-38.3)	4.1	(3.4-5.0)	26.9	(24.9-29.1)	10.8	(9.6-12.3)
Arthritis	Yes	1,894	20.5	(18.3-22.9)	35.3	(32.0-38.8)	3.3	(2.5-4.3)	28.3	(25.2-31.6)	12.6	(10.6-14.9)
	No	3,210	21.8	(19.2-24.8)	36.0	(33.2-38.9)	4.5	(3.6-5.7)	26.9	(24.3-29.6)	10.7	(9.3-12.3)
Frequent Mental Distress	Yes	443	15.7	(12.0-20.3)	31.3	(24.1-39.5)	4.0	(2.1-7.6)*	30.4	(23.9-37.8)	18.6	(14.1-24.0)
	No	4,625	22.1	(20.0-24.5)	36.4	(34.0-38.9)	4.2	(3.4-5.0)	26.9	(24.7-29.2)	10.4	(9.2-11.8)
Serious Psychological Distress	Yes	180	18.1	(12.4-25.5)	25.4	(17.3-35.7)	4.4	(1.3-14.5)*	32.2	(23.3-42.6)	19.9	(13.7-28.0)
	No	4,876	21.5	(19.4-23.7)	36.3	(33.9-38.7)	4.2	(3.5-5.1)	27.3	(25.1-29.5)	10.7	(9.5-12.1)
Receiving Treatment for Mental Health Condition	Yes	689	17.5	(13.9-21.6)	35.4	(30.0-41.1)	3.5	(2.0-6.1)	28.2	(23.6-33.3)	15.4	(12.3-19.2)
	No	4,433	22.0	(19.8-24.3)	36.0	(33.6-38.5)	4.3	(3.5-5.2)	27.1	(24.9-29.5)	10.6	(9.4-12.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,883	70.6	(69.0-72.1)	22.8	(21.4-24.3)	1.3	(0.9-1.8)	3.5	(2.8-4.4)	1.8	(1.3-2.4)
Age (years)	18-24	207	52.4	(44.5-60.2)	40.1	(32.7-47.9)	2.0	(0.7-5.4)*	4.1	(1.6-10.2)*	1.4	(0.4-5.2)*
	25-34	543	71.2	(66.3-75.6)	22.6	(18.6-27.3)	0.4	(0.1-1.3)*	3.4	(1.8-6.4)*	2.4	(1.2-4.6)*
	35-54	2,433	72.3	(70.0-74.5)	20.5	(18.5-22.6)	1.6	(1.0-2.4)	3.7	(2.8-5.0)	1.9	(1.3-2.9)
	55+	3,651	74.2	(72.4-75.9)	20.1	(18.6-21.8)	1.1	(0.7-1.6)	3.0	(2.3-3.8)	1.6	(1.1-2.3)
Sex	Men	2,753	63.9	(61.4-66.3)	27.5	(25.3-29.9)	1.5	(1.0-2.3)	4.8	(3.7-6.3)	2.2	(1.5-3.4)
	Women	4,130	76.8	(75.0-78.6)	18.3	(16.7-20.0)	1.1	(0.7-1.7)	2.3	(1.6-3.3)	1.4	(1.0-2.1)
Race/ Ethnicity	Hispanic or Latino	309	61.5	(54.7-67.9)	19.7	(14.2-26.5)	4.7	(2.6-8.3)	2.9	(1.5-5.8)*	11.2	(8.7-14.3)
	Not Hispanic or Latino:	5,991	70.5	(68.9-72.1)	23.2	(21.7-24.7)	1.3	(0.9-1.7)	3.5	(2.8-4.3)	1.6	(1.2-2.2)
	White											
	Black or African American	92	75.3	(63.9-84.0)	18.4	(11.0-29.1)	0.5	(0.4-0.7)	5.0	(1.5-15.7)*	0.8	(0.3-2.2)*
	All Other Races	429	63.8	(57.8-69.5)	26.5	(21.4-32.3)	0.8	(0.4-1.6)*	6.8	(4.0-11.3)	2.1	(1.0-4.7)*
Education	<High School	340	63.5	(56.1-70.2)	24.6	(18.5-32.0)	3.5	(1.6-7.5)*	3.8	(2.1-6.8)*	4.6	(2.2-9.5)*
	High School Graduate (or GED)	1,422	63.0	(59.4-66.5)	30.3	(27.0-33.8)	1.5	(0.8-2.5)	3.2	(2.2-4.6)	2.0	(1.2-3.4)
	Some College	2,257	72.0	(69.2-74.6)	21.1	(18.8-23.7)	1.2	(0.7-2.0)	4.3	(2.9-6.4)	1.4	(0.8-2.4)
	College Graduate	2,856	76.8	(74.5-78.9)	18.7	(16.8-20.9)	0.9	(0.4-1.9)*	2.7	(1.8-3.9)	0.9	(0.5-1.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	671	66.9 (61.4-71.9)	23.9 (19.6-28.9)	2.0 (1.1-3.9)*	4.5 (2.6-7.9)	2.6 (1.4-4.8)*				
	\$20,000-\$34,999	1,313	71.4 (67.8-74.8)	22.3 (19.2-25.8)	0.9 (0.4-1.9)*	3.9 (2.6-6.0)	1.4 (0.9-2.3)				
	\$35,000-\$49,999	1,001	68.8 (64.4-72.8)	24.5 (20.8-28.7)	0.8 (0.3-1.9)*	3.7 (1.9-7.3)*	2.2 (1.1-4.6)*				
	\$50,000-\$74,999	1,183	73.8 (70.3-77.1)	20.8 (17.8-24.2)	1.4 (0.6-3.2)*	2.9 (1.9-4.5)	1.0 (0.4-2.5)*				
	\$75,000 and more	1,989	74.7 (72.0-77.2)	21.3 (18.9-23.9)	0.8 (0.4-1.4)*	2.4 (1.6-3.6)	0.9 (0.4-1.8)*				
Employment Status	Employed	3,490	70.0 (67.8-72.0)	22.7 (20.8-24.7)	1.4 (0.9-2.1)	3.7 (2.7-5.0)	2.3 (1.5-3.4)				
	Out of work	412	73.3 (67.2-78.5)	20.9 (16.1-26.6)	1.5 (0.6-3.5)*	3.4 (1.5-7.3)*	1.1 (0.4-2.8)*				
	Homemaker	492	71.2 (65.8-76.1)	21.7 (17.4-26.8)	1.3 (0.3-4.4)*	3.7 (1.8-7.6)*	2.0 (0.8-5.1)*				
	Student	127	72.2 (62.2-80.4)	24.7 (17.1-34.4)	1.8 (0.4-7.1)*	1.3 (0.2-8.6)*	0.0 (0.0-0.0)				
	Retired	1,945	70.4 (67.6-73.1)	23.7 (21.1-26.4)	1.1 (0.7-1.8)	3.8 (2.7-5.3)	1.0 (0.6-1.7)				
	Unable to work	395	70.5 (64.3-76.0)	24.4 (19.1-30.6)	0.5 (0.2-1.5)*	2.9 (1.5-5.5)*	1.7 (0.8-3.6)*				
Veteran Status	Yes	1,033	69.2 (65.5-72.6)	25.2 (21.9-28.8)	1.1 (0.5-2.5)*	2.7 (1.8-4.0)	1.8 (1.0-3.2)*				
	No	5,835	70.7 (68.9-72.4)	22.5 (21.0-24.1)	1.3 (1.0-1.9)	3.6 (2.9-4.6)	1.8 (1.3-2.5)				
Diabetes	Yes	741	72.8 (68.5-76.8)	22.4 (18.6-26.8)	0.7 (0.3-1.7)*	3.0 (1.7-5.3)	1.0 (0.4-2.6)*				
	No	6,139	70.3 (68.6-71.9)	22.9 (21.4-24.5)	1.4 (1.0-1.9)	3.6 (2.8-4.5)	1.9 (1.4-2.6)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	645	69.4	(64.9-73.5)	26.1	(22.1-30.6)	0.3	(0.1-1.4)*	3.0	(1.7-5.2)	1.2	(0.6-2.5)*
	No	6,169	70.8	(69.2-72.4)	22.6	(21.1-24.1)	1.4	(1.0-1.9)	3.4	(2.7-4.3)	1.8	(1.3-2.5)
Asthma	Yes	1,060	72.7	(68.7-76.3)	20.7	(17.4-24.4)	1.9	(0.9-3.9)*	2.6	(1.5-4.5)	2.1	(0.8-5.2)*
	No	5,807	70.3	(68.6-71.9)	23.1	(21.6-24.7)	1.2	(0.9-1.7)	3.7	(2.9-4.7)	1.7	(1.3-2.3)
Arthritis	Yes	2,516	72.5	(69.8-74.9)	21.4	(19.3-23.8)	1.3	(0.7-2.3)*	3.4	(2.1-5.5)	1.4	(0.9-2.2)
	No	4,339	69.9	(68.0-71.8)	23.2	(21.5-25.0)	1.3	(0.9-1.9)	3.6	(2.8-4.6)	1.9	(1.3-2.8)
Frequent Mental Distress	Yes	655	68.2	(63.1-72.8)	21.5	(17.6-26.1)	1.4	(0.6-3.2)*	5.8	(3.4-9.6)	3.1	(1.2-7.5)*
	No	6,157	70.8	(69.1-72.4)	23.0	(21.5-24.6)	1.3	(0.9-1.8)	3.3	(2.6-4.2)	1.6	(1.2-2.2)
Serious Psychological Distress	Yes	214	62.4	(53.4-70.7)	26.2	(19.1-34.8)	0.7	(0.2-2.5)*	8.0	(4.4-14.3)*	2.6	(1.1-6.2)*
	No	6,565	71.3	(69.7-72.9)	22.8	(21.3-24.3)	1.3	(0.9-1.7)	3.2	(2.5-4.1)	1.5	(1.1-2.0)
Receiving Treatment for Mental Health Condition	Yes	1,015	78.5	(74.7-81.8)	17.0	(14.0-20.6)	1.5	(0.6-3.7)*	2.1	(1.2-3.7)	0.9	(0.4-2.0)*
	No	5,850	69.4	(67.6-71.0)	23.7	(22.1-25.3)	1.3	(0.9-1.8)	3.8	(3.0-4.7)	1.9	(1.4-2.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		6,853	18.9	(17.6-20.2)	33.7	(32.2-35.3)	2.3	(1.8-2.9)	28.3	(26.8-29.7)	16.9	(15.7-18.1)
Age (years)	18-24	207	16.4	(11.4-23.0)	44.9	(37.1-53.0)	3.8	(1.6-8.6)*	26.5	(20.1-34.0)	8.4	(4.8-14.3)
	25-34	540	16.9	(13.5-21.1)	35.6	(31.1-40.5)	2.2	(1.1-4.1)*	32.3	(27.9-37.1)	13.0	(10.0-16.6)
	35-54	2,437	16.7	(15.0-18.6)	33.0	(30.8-35.3)	2.0	(1.5-2.8)	28.5	(26.4-30.7)	19.7	(17.8-21.8)
	55+	3,623	24.0	(22.3-25.7)	30.2	(28.4-32.0)	2.2	(1.6-2.9)	26.0	(24.4-27.6)	17.7	(16.3-19.3)
Sex	Men	2,748	21.0	(19.1-23.1)	37.3	(34.9-39.7)	2.2	(1.5-3.2)	26.6	(24.5-28.8)	12.9	(11.3-14.7)
	Women	4,105	17.0	(15.5-18.6)	30.2	(28.3-32.2)	2.3	(1.8-3.0)	29.9	(28.1-31.7)	20.6	(19.1-22.3)
Race/ Ethnicity	Hispanic or Latino	310	32.2	(26.2-38.8)	27.4	(21.5-34.2)	2.9	(1.1-7.0)*	24.5	(18.4-31.7)	13.2	(8.9-19.0)
	Not Hispanic or Latino:	5,958	17.6	(16.3-18.9)	33.3	(31.6-35.0)	2.2	(1.7-2.7)	29.6	(28.0-31.2)	17.4	(16.2-18.7)
	White											
	Black or African American	90	28.5	(18.0-41.9)	22.3	(13.2-35.1)	0.7	(0.5-0.9)	12.0	(6.5-21.2)*	36.5	(25.6-49.1)
	All Other Races	433	21.9	(17.2-27.4)	32.1	(26.7-38.2)	2.1	(0.7-5.8)*	21.6	(17.1-26.8)	22.3	(18.0-27.3)
Education	<High School	349	33.9	(27.3-41.2)	26.4	(20.6-33.3)	1.9	(1.0-3.9)*	21.8	(16.5-28.3)	15.9	(11.0-22.4)
	High School Graduate (or GED)	1,424	22.0	(19.3-24.9)	35.2	(31.7-38.8)	2.2	(1.2-3.8)	25.1	(22.1-28.4)	15.5	(13.2-18.2)
	Some College	2,238	16.7	(14.7-18.8)	33.5	(30.8-36.3)	1.8	(1.1-3.1)	28.7	(26.1-31.4)	19.3	(17.3-21.6)
	College Graduate	2,835	14.1	(12.4-16.0)	34.2	(31.9-36.6)	2.9	(2.1-3.8)	32.3	(30.0-34.7)	16.5	(14.8-18.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	675	20.7	(16.6-25.6)	32.8	(27.6-38.4)	2.9	(1.5-5.6)*	23.1	(18.9-28.0)	20.5	(16.6-25.0)
	\$20,000-\$34,999	1,306	20.6	(17.9-23.6)	32.6	(29.0-36.5)	1.2	(0.6-2.5)*	28.8	(25.3-32.6)	16.7	(14.2-19.5)
	\$35,000-\$49,999	992	18.6	(15.8-21.9)	33.4	(29.5-37.7)	2.4	(1.4-3.9)	29.8	(26.0-34.0)	15.7	(13.2-18.7)
	\$50,000-\$74,999	1,181	16.5	(13.9-19.4)	35.1	(31.7-38.7)	1.9	(1.2-3.0)	29.5	(26.4-32.7)	17.0	(14.4-20.1)
	\$75,000 and more	1,980	17.7	(15.4-20.2)	35.9	(33.2-38.7)	1.6	(1.1-2.4)	28.9	(26.5-31.5)	15.9	(14.0-18.0)
Employment Status	Employed	3,468	18.7	(17.0-20.6)	33.9	(31.9-36.0)	2.2	(1.7-3.0)	28.2	(26.3-30.2)	16.9	(15.3-18.7)
	Out of work	411	16.9	(12.8-22.1)	38.6	(32.3-45.3)	1.2	(0.5-2.6)*	27.1	(22.0-32.9)	16.2	(11.9-21.7)
	Homemaker	495	20.0	(16.0-24.6)	32.1	(27.1-37.5)	2.8	(1.4-5.8)*	29.9	(25.2-35.0)	15.2	(12.1-19.1)
	Student	128	14.7	(8.3-24.9)	35.5	(26.2-46.2)	5.5	(1.7-16.1)*	29.0	(20.3-39.6)	15.2	(9.3-24.0)
	Retired	1,932	22.8	(20.5-25.3)	33.9	(31.2-36.8)	2.1	(1.4-3.2)	26.6	(24.2-29.2)	14.6	(12.7-16.7)
	Unable to work	397	14.1	(10.2-19.2)	24.6	(19.1-31.2)	1.8	(0.7-4.3)*	31.4	(25.3-38.2)	28.1	(22.7-34.2)
Veteran Status	Yes	1,043	18.4	(15.7-21.3)	33.3	(29.8-37.0)	1.8	(1.0-3.0)	28.1	(24.5-32.0)	18.5	(15.4-22.0)
	No	5,796	19.0	(17.6-20.4)	33.7	(32.0-35.4)	2.4	(1.8-3.0)	28.3	(26.8-29.9)	16.7	(15.5-18.1)
Diabetes	Yes	738	18.8	(15.6-22.4)	31.5	(27.2-36.2)	1.4	(0.7-2.5)*	26.8	(22.9-31.1)	21.6	(18.1-25.5)
	No	6,112	18.9	(17.6-20.3)	33.9	(32.3-35.5)	2.3	(1.8-3.0)	28.4	(26.9-29.9)	16.4	(15.2-17.7)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Washington, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	642	19.5	(16.1-23.3)	31.6	(27.3-36.3)	2.0	(1.1-3.5)*	25.5	(21.6-29.9)	21.5	(17.7-25.7)
	No	6,143	18.8	(17.5-20.2)	33.8	(32.2-35.5)	2.3	(1.8-3.0)	28.5	(27.0-30.1)	16.5	(15.3-17.8)
Asthma	Yes	1,056	17.1	(13.9-20.7)	32.7	(28.8-36.9)	2.7	(1.6-4.5)	27.7	(24.3-31.4)	19.8	(16.9-23.2)
	No	5,780	19.2	(17.9-20.7)	33.7	(32.1-35.4)	2.2	(1.7-2.8)	28.5	(27.0-30.1)	16.4	(15.1-17.7)
Arthritis	Yes	2,510	17.9	(16.1-19.9)	32.7	(30.2-35.3)	1.6	(1.1-2.3)	27.9	(25.6-30.3)	19.8	(17.9-22.0)
	No	4,316	19.3	(17.7-20.9)	33.9	(32.1-35.9)	2.5	(1.9-3.2)	28.5	(26.8-30.3)	15.8	(14.4-17.3)
Frequent Mental Distress	Yes	652	13.5	(10.5-17.1)	29.2	(24.5-34.4)	1.2	(0.6-2.3)*	28.7	(24.2-33.7)	27.4	(22.8-32.7)
	No	6,133	19.7	(18.4-21.2)	34.2	(32.6-35.8)	2.3	(1.8-3.0)	28.2	(26.7-29.7)	15.6	(14.4-16.8)
Serious Psychological Distress	Yes	214	10.1	(6.3-15.7)	28.1	(20.4-37.4)	1.8	(0.4-7.9)*	24.4	(17.2-33.5)	35.6	(27.7-44.4)
	No	6,540	19.2	(17.9-20.6)	34.2	(32.6-35.8)	2.3	(1.8-2.9)	28.4	(27.0-29.9)	15.9	(14.7-17.1)
Receiving Treatment for Mental Health Condition	Yes	1,006	16.1	(13.5-19.1)	29.6	(25.7-33.8)	2.0	(1.3-3.3)	28.2	(24.5-32.2)	24.1	(21.0-27.5)
	No	5,833	19.8	(18.4-21.2)	33.5	(31.9-35.2)	2.2	(1.7-2.9)	27.9	(26.4-29.5)	16.6	(15.3-17.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,241	62.9	(60.7-65.1)	30.3	(28.2-32.4)	0.7	(0.4-1.0)	4.4	(3.4-5.7)	1.7	(1.2-2.5)
Age (years)												
	18-24	199	46.6	(37.0-56.5)	43.4	(34.2-53.0)	0.3	(0.1-1.6)*	7.9	(3.8-15.6)*	1.8	(0.4-7.2)*
	25-34	488	58.2	(52.5-63.7)	35.2	(29.9-41.0)	0.5	(0.1-1.8)*	5.5	(3.1-9.6)	0.5	(0.2-1.6)*
	35-54	1,691	64.4	(61.4-67.3)	29.2	(26.4-32.2)	0.6	(0.3-1.1)*	3.7	(2.6-5.2)	2.2	(1.3-3.4)
	55+	1,840	70.4	(67.6-72.9)	23.3	(20.9-25.9)	1.1	(0.6-2.1)*	3.3	(2.4-4.6)	1.9	(1.2-3.0)
Sex												
	Men	1,617	55.9	(52.5-59.2)	34.6	(31.4-37.9)	1.0	(0.6-1.7)	6.3	(4.5-8.7)	2.3	(1.5-3.5)
	Women	2,624	69.4	(66.6-72.1)	26.2	(23.7-29.0)	0.4	(0.2-0.7)*	2.7	(1.7-4.1)	1.2	(0.6-2.4)*
Race/Ethnicity												
	Hispanic or Latino	77	56.9	(42.3-70.4)	29.7	(17.7-45.4)	0.1	(0.0-0.1)	5.7	(1.8-16.6)*	7.6	(1.7-27.7)*
	Not Hispanic or Latino:	3,542	64.4	(62.1-66.6)	30.0	(27.8-32.2)	0.7	(0.4-1.1)	3.7	(2.7-5.1)	1.2	(0.8-1.7)
	White											
	Black or African American	450	60.7	(51.7-69.0)	29.2	(21.2-38.7)	0.7	(0.3-1.6)*	7.3	(3.5-14.5)*	2.2	(1.1-4.3)*
	All Other Races	147	52.7	(38.3-66.7)	34.1	(22.3-48.2)	1.2	(0.2-5.6)*	9.7	(4.7-19.0)*	2.3	(0.8-6.1)*
Education												
	<High School	295	62.1	(53.9-69.7)	29.9	(22.9-37.9)	0.8	(0.3-2.0)*	5.7	(3.2-9.8)	1.5	(0.7-3.1)*
	High School Graduate (or GED)	1,440	59.0	(55.0-63.0)	32.9	(29.2-36.8)	0.8	(0.4-1.4)*	4.9	(3.2-7.4)	2.4	(1.3-4.2)
	Some College	1,206	58.4	(54.2-62.4)	34.8	(30.8-39.0)	0.6	(0.3-1.2)*	4.1	(2.8-6.1)	2.1	(1.1-3.7)
	College Graduate	1,297	70.5	(66.8-74.0)	23.9	(20.7-27.5)	0.8	(0.3-1.8)*	4.0	(2.0-7.8)*	0.7	(0.3-1.9)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	496	55.1	(47.4-62.6)	34.2	(27.1-42.0)	0.6	(0.2-1.4)*	5.2	(3.1-8.6)	4.9	(2.0-11.6)*
	\$20,000-\$34,999	1,196	57.0	(52.6-61.3)	34.8	(30.7-39.2)	0.6	(0.3-1.2)*	5.4	(3.4-8.4)	2.2	(1.2-3.9)
	\$35,000-\$49,999	741	65.2	(59.9-70.2)	26.6	(22.2-31.6)	0.9	(0.3-3.0)*	6.1	(3.1-11.8)*	1.1	(0.5-2.4)*
	\$50,000-\$74,999	688	67.2	(62.4-71.6)	29.1	(24.8-33.8)	0.7	(0.3-1.9)*	2.3	(1.3-4.3)*	0.7	(0.2-2.3)*
	\$75,000 and more	778	67.7	(63.0-72.0)	27.4	(23.3-32.1)	0.6	(0.2-1.6)*	3.4	(2.0-5.9)	0.8	(0.3-2.6)*
Employment Status	Employed	2,581	62.1	(59.3-64.7)	31.0	(28.5-33.7)	0.7	(0.4-1.1)	4.8	(3.4-6.8)	1.4	(0.8-2.5)
	Out of work	162	65.3	(51.4-77.0)	28.4	(18.2-41.5)	0.2	(0.0-0.8)*	4.8	(1.9-11.7)*	1.3	(0.4-4.4)*
	Homemaker	245	68.7	(61.4-75.3)	26.2	(20.0-33.6)	0.9	(0.2-3.8)*	1.7	(0.6-4.3)*	2.5	(1.0-6.1)*
	Student	91	62.7	(47.4-75.8)	33.1	(20.7-48.5)	0.1	(0.0-1.0)*	2.4	(0.8-6.8)*	1.7	(0.3-10.7)*
	Retired	948	64.3	(60.3-68.1)	28.0	(24.3-32.0)	1.1	(0.4-2.7)*	4.6	(3.1-6.9)	2.0	(1.1-3.7)*
	Unable to work	210	62.4	(53.0-70.9)	29.5	(21.4-39.3)	0.2	(0.0-0.9)*	2.7	(1.1-6.8)*	5.2	(2.3-11.5)*
Veteran Status	Yes	422	62.0	(56.3-67.4)	27.4	(22.5-33.0)	1.6	(0.5-4.6)*	7.0	(4.2-11.4)	2.0	(1.0-3.9)*
	No	3,818	62.9	(60.6-65.2)	30.6	(28.4-32.9)	0.6	(0.4-0.9)	4.2	(3.1-5.6)	1.7	(1.1-2.6)
Diabetes	Yes	381	67.3	(60.9-73.1)	26.0	(20.6-32.3)	0.3	(0.1-0.8)*	3.7	(1.7-7.9)*	2.7	(1.3-5.3)*
	No	3,860	62.5	(60.2-64.8)	30.5	(28.4-32.8)	0.8	(0.5-1.1)	4.5	(3.4-5.9)	1.7	(1.1-2.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	369	64.4	(57.9-70.3)	29.6	(23.8-36.1)	0.5	(0.1-1.9)*	3.6	(2.0-6.6)*	1.8	(0.6-5.4)*
	No	3,844	62.6	(60.3-64.8)	30.5	(28.3-32.7)	0.7	(0.4-1.1)	4.5	(3.4-5.9)	1.7	(1.1-2.6)
Asthma	Yes	581	67.5	(61.0-73.4)	27.9	(22.2-34.3)	0.5	(0.2-1.5)*	2.9	(1.5-5.5)*	1.2	(0.5-3.2)*
	No	3,657	62.0	(59.7-64.3)	30.6	(28.4-32.9)	0.8	(0.5-1.2)	4.7	(3.6-6.2)	1.9	(1.3-2.8)
Arthritis	Yes	1,516	65.9	(62.5-69.2)	28.2	(25.1-31.5)	0.8	(0.5-1.4)	3.3	(2.2-4.8)	1.8	(1.0-3.0)
	No	2,723	61.7	(59.0-64.4)	31.1	(28.5-33.7)	0.6	(0.3-1.1)*	4.9	(3.6-6.6)	1.7	(1.1-2.8)
Frequent Mental Distress	Yes	387	63.6	(55.6-71.0)	29.0	(22.3-36.9)	0.1	(0.0-0.4)*	3.0	(1.4-6.5)*	4.2	(2.3-7.4)
	No	3,829	62.9	(60.6-65.2)	30.4	(28.2-32.6)	0.7	(0.4-1.1)	4.5	(3.4-6.0)	1.4	(0.9-2.3)
Serious Psychological Distress	Yes	141	56.2	(43.1-68.5)	35.4	(23.8-48.9)	0.4	(0.1-1.0)*	4.5	(2.0-9.6)*	3.6	(1.7-7.6)*
	No	4,068	63.2	(60.9-65.4)	30.1	(28.0-32.3)	0.7	(0.5-1.0)	4.4	(3.3-5.7)	1.7	(1.1-2.5)
Receiving Treatment for Mental Health Condition	Yes	441	72.7	(65.8-78.7)	21.6	(16.2-28.2)	1.0	(0.2-5.7)*	2.2	(1.0-4.6)*	2.5	(1.3-4.6)*
	No	3,797	62.0	(59.6-64.2)	31.1	(28.9-33.4)	0.6	(0.4-1.0)	4.7	(3.5-6.1)	1.7	(1.1-2.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		4,266	22.3	(20.5-24.2)	39.5	(37.3-41.6)	0.5	(0.4-0.8)	28.0	(26.0-29.9)	9.7	(8.5-11.1)
Age (years)	18-24	200	11.9	(6.9-19.9)	49.0	(39.6-58.5)	0.0	(0.0-0.0)	30.5	(22.6-39.8)	8.6	(4.6-15.2)*
	25-34	494	21.9	(17.3-27.3)	42.1	(36.6-47.8)	0.1	(0.0-0.9)*	28.2	(23.4-33.6)	7.6	(5.1-11.4)
	35-54	1,702	21.9	(19.3-24.8)	38.0	(35.0-41.0)	0.7	(0.4-1.3)*	28.6	(26.0-31.5)	10.7	(9.0-12.8)
	55+	1,845	28.4	(25.8-31.1)	35.8	(33.0-38.7)	0.8	(0.4-1.6)*	25.4	(23.0-28.0)	9.6	(8.0-11.5)
Sex	Men	1,628	23.3	(20.5-26.4)	43.7	(40.4-47.1)	0.9	(0.5-1.4)	25.0	(22.1-28.0)	7.1	(5.5-9.2)
	Women	2,638	21.4	(19.3-23.8)	35.4	(32.7-38.2)	0.3	(0.1-0.6)*	30.8	(28.3-33.5)	12.1	(10.4-14.0)
Race/ Ethnicity	Hispanic or Latino	77	22.5	(12.1-37.8)	39.1	(25.5-54.7)	1.0	(0.2-6.2)*	19.1	(10.7-31.7)	18.3	(7.6-38.1)*
	Not Hispanic or Latino:	3,561	20.2	(18.4-22.1)	40.4	(38.2-42.7)	0.7	(0.4-1.0)	29.6	(27.5-31.7)	9.2	(8.0-10.5)
	White											
	Black or African American	454	15.8	(11.2-21.7)	39.3	(29.6-49.9)	0.4	(0.2-1.0)*	34.1	(24.3-45.6)	10.3	(7.3-14.4)
	All Other Races	150	21.4	(12.7-33.7)	36.3	(24.8-49.7)	0.4	(0.2-0.9)*	23.2	(15.5-33.4)	18.6	(8.8-35.2)*
Education	<High School	300	33.0	(25.7-41.1)	31.9	(24.7-40.2)	0.1	(0.0-0.5)*	23.5	(17.2-31.1)	11.6	(7.5-17.3)
	High School Graduate (or GED)	1,446	23.5	(20.7-26.6)	40.2	(36.3-44.3)	0.7	(0.3-1.4)*	28.3	(24.7-32.1)	7.3	(5.7-9.3)
	Some College	1,218	21.9	(18.5-25.8)	38.9	(35.0-43.1)	0.8	(0.4-1.5)*	27.3	(23.8-31.1)	11.1	(8.6-14.1)
	College Graduate	1,299	18.0	(15.0-21.5)	40.9	(37.3-44.7)	0.4	(0.1-0.9)*	29.7	(26.6-33.1)	11.0	(8.8-13.6)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	506	12.9	(10.0-16.5)	34.1	(27.3-41.5)	1.3	(0.4-3.6)*	32.6	(25.7-40.3)	19.2	(13.5-26.5)
	\$20,000-\$34,999	1,207	24.7	(21.3-28.5)	37.5	(33.3-42.0)	0.5	(0.2-1.2)*	28.5	(24.7-32.7)	8.8	(6.9-11.1)
	\$35,000-\$49,999	741	23.3	(18.5-28.8)	37.9	(32.9-43.3)	0.9	(0.3-2.1)*	28.9	(24.5-33.7)	9.1	(6.6-12.4)
	\$50,000-\$74,999	688	22.9	(19.1-27.2)	44.5	(39.7-49.3)	0.4	(0.1-1.0)*	24.0	(20.2-28.3)	8.2	(6.0-11.1)
	\$75,000 and more	782	22.5	(18.8-26.9)	39.8	(35.6-44.3)	0.3	(0.1-0.9)*	28.6	(24.9-32.7)	8.6	(6.3-11.8)
Employment Status	Employed	2,598	21.9	(19.6-24.4)	41.2	(38.6-44.0)	0.6	(0.5-0.9)	27.3	(25.0-29.8)	8.9	(7.5-10.5)
	Out of work	165	16.7	(9.9-26.9)	36.8	(24.9-50.6)	0.9	(0.3-2.9)*	38.7	(26.5-52.5)	6.9	(3.5-13.1)*
	Homemaker	244	24.8	(19.1-31.5)	40.1	(32.6-48.1)	0.4	(0.3-0.5)	26.4	(20.0-33.9)	8.3	(5.1-13.4)
	Student	91	20.3	(10.1-36.6)*	31.9	(20.1-46.6)	0.5	(0.3-0.6)	29.4	(18.6-43.1)	18.0	(8.2-35.0)*
	Retired	954	25.1	(21.8-28.6)	36.9	(32.9-41.0)	1.1	(0.5-2.4)*	27.4	(23.8-31.3)	9.6	(7.3-12.4)
	Unable to work	210	21.0	(13.6-31.0)	27.6	(19.5-37.4)	2.6	(0.7-9.7)*	29.5	(21.4-39.2)	19.3	(12.6-28.4)
Veteran Status	Yes	428	22.5	(18.1-27.8)	37.0	(31.4-43.0)	1.3	(0.5-3.4)*	27.9	(22.6-33.9)	11.3	(7.6-16.3)
	No	3,837	22.2	(20.3-24.3)	39.6	(37.3-42.0)	0.7	(0.5-1.0)	27.9	(25.9-30.0)	9.6	(8.3-11.0)
Diabetes	Yes	387	25.9	(20.7-31.8)	35.1	(28.9-41.8)	0.5	(0.3-0.7)	25.6	(20.0-32.0)	13.0	(9.3-17.9)
	No	3,879	22.0	(20.1-24.0)	39.7	(37.4-42.0)	0.8	(0.6-1.1)	28.0	(26.1-30.1)	9.5	(8.2-10.9)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wisconsin, 2007 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	372	25.2	(19.7-31.7)	36.1	(29.8-43.0)	0.6	(0.1-4.2)*	26.6	(21.0-33.1)	11.4	(8.0-16.0)
	No	3,867	22.3	(20.4-24.3)	39.7	(37.5-42.0)	0.5	(0.3-0.8)	27.9	(25.9-30.0)	9.5	(8.3-11.0)
Asthma	Yes	581	21.4	(17.2-26.4)	35.7	(29.9-42.0)	0.2	(0.0-0.8)*	31.7	(26.2-37.7)	11.0	(8.4-14.3)
	No	3,681	22.5	(20.5-24.5)	40.0	(37.8-42.4)	0.6	(0.4-1.0)	27.4	(25.4-29.5)	9.5	(8.2-11.0)
Arthritis	Yes	1,522	22.9	(20.2-25.8)	34.9	(31.6-38.4)	0.8	(0.4-1.7)*	29.5	(26.5-32.8)	11.8	(9.7-14.3)
	No	2,742	22.0	(19.7-24.4)	41.1	(38.5-43.9)	0.4	(0.2-0.7)	27.5	(25.2-30.0)	9.0	(7.5-10.6)
Frequent Mental Distress	Yes	388	14.4	(9.8-20.6)	32.0	(25.1-39.9)	0.9	(0.3-3.3)*	32.6	(25.6-40.4)	20.0	(15.4-25.6)
	No	3,851	23.0	(21.1-25.1)	40.1	(37.8-42.4)	0.7	(0.5-1.0)	27.5	(25.6-29.6)	8.6	(7.4-10.0)
Serious Psychological Distress	Yes	141	22.7	(13.3-35.9)	21.9	(13.7-33.1)	1.9	(0.3-12.2)*	30.2	(19.5-43.6)	23.3	(14.7-35.0)
	No	4,095	22.5	(20.7-24.4)	40.0	(37.8-42.2)	0.5	(0.3-0.8)	27.8	(25.9-29.9)	9.2	(8.0-10.6)
Receiving Treatment for Mental Health Condition	Yes	444	21.4	(16.3-27.6)	29.2	(23.2-36.1)	0.9	(0.2-3.4)*	32.4	(26.3-39.1)	16.1	(12.4-20.7)
	No	3,819	22.4	(20.5-24.4)	40.5	(38.2-42.8)	0.5	(0.3-0.8)	27.5	(25.5-29.6)	9.1	(7.8-10.5)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,482	65.1	(63.2-66.9)	27.1	(25.3-28.9)	0.9	(0.6-1.2)	4.9	(3.9-6.1)	2.1	(1.3-3.4)
Age (years)	18-24	125	52.1	(43.2-60.8)	37.6	(29.1-46.8)	0.1	(0.1-0.1)	8.1	(4.2-15.1)*	2.1	(0.3-12.4)*
	25-34	503	60.2	(55.3-65.0)	30.5	(26.2-35.3)	1.3	(0.6-2.8)*	6.5	(4.2-9.9)	1.4	(0.4-4.7)*
	35-54	1,801	67.5	(65.0-70.0)	25.3	(23.0-27.7)	0.7	(0.4-1.3)	3.9	(2.9-5.1)	2.6	(1.6-4.1)
	55+	3,025	69.3	(67.4-71.2)	23.8	(22.1-25.6)	1.3	(0.9-1.9)	3.5	(2.8-4.5)	2.0	(1.5-2.8)
Sex	Men	2,167	57.8	(54.9-60.6)	32.0	(29.1-34.9)	0.5	(0.3-0.9)	6.1	(4.6-8.0)	3.6	(2.0-6.4)
	Women	3,315	72.0	(69.6-74.2)	22.3	(20.3-24.6)	1.2	(0.8-1.8)	3.7	(2.5-5.5)	0.8	(0.5-1.2)
Race/ Ethnicity	Hispanic or Latino	215	54.0	(45.2-62.7)	25.2	(18.0-34.0)	1.8	(0.7-4.5)*	16.4	(10.5-24.8)	2.6	(1.1-5.7)*
	Not Hispanic or Latino:	5,074	65.9	(64.0-67.7)	27.1	(25.3-28.9)	0.9	(0.7-1.2)	4.5	(3.6-5.6)	1.5	(1.1-2.0)
	White											
	Black or African American	20	--	--	--	--	--	--	--	--	--	--
	All Other Races	124	9.0	(1.6-37.4)*	4.4	(0.1-68.0)*	0.2	(0.0-97.9)*	0.4	(0.0-91.3)*	86.1	(53.2-97.1)
Education	<High School	262	62.9	(53.4-71.4)	20.5	(14.5-28.2)	0.8	(0.2-2.6)*	11.5	(6.0-21.1)*	4.3	(1.9-9.7)*
	High School Graduate (or GED)	1,640	57.9	(54.2-61.4)	32.1	(28.6-35.8)	0.8	(0.5-1.4)	6.0	(4.2-8.5)	3.2	(1.4-7.1)*
	Some College	1,763	64.8	(61.5-67.9)	28.7	(25.7-31.9)	1.0	(0.6-1.8)	4.1	(2.6-6.3)	1.4	(0.7-2.7)*
	College Graduate	1,813	72.8	(69.9-75.6)	22.3	(19.7-25.1)	1.0	(0.5-1.9)*	2.9	(1.9-4.3)	1.0	(0.4-2.3)*

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	717	59.0	(52.5-65.1)	26.9	(20.9-34.0)	0.5	(0.2-1.3)*	6.8	(3.7-12.1)*	6.8	(2.7-16.0)*
	\$20,000-\$34,999	968	62.1	(57.4-66.5)	30.3	(26.1-35.0)	1.1	(0.5-2.4)*	5.3	(3.1-9.0)	1.1	(0.6-2.0)
	\$35,000-\$49,999	780	64.3	(59.6-68.7)	30.0	(25.7-34.6)	1.0	(0.5-2.1)*	3.6	(2.0-6.2)	1.2	(0.6-2.5)*
	\$50,000-\$74,999	975	69.8	(65.8-73.5)	25.4	(21.9-29.2)	0.5	(0.2-1.2)*	3.4	(2.1-5.5)	0.9	(0.3-3.2)*
	\$75,000 and more	1,481	70.6	(67.5-73.6)	23.8	(21.0-26.8)	0.8	(0.4-1.4)*	3.1	(2.0-4.7)	1.7	(0.9-3.3)*
Employment Status	Employed	3,073	65.7	(63.4-68.0)	26.9	(24.7-29.1)	0.8	(0.5-1.2)	5.1	(3.8-6.8)	1.5	(1.0-2.3)
	Out of work	181	62.2	(51.9-71.5)	29.5	(20.7-40.1)	0.2	(0.0-1.2)*	5.1	(2.2-11.5)*	3.1	(0.9-10.0)*
	Homemaker	438	60.6	(54.2-66.6)	30.1	(24.3-36.6)	2.6	(1.2-5.7)*	5.9	(2.2-14.7)*	0.9	(0.3-2.6)*
	Student	70	65.1	(49.0-78.3)	21.0	(11.8-34.6)	0.0	(0.0-0.0)	3.0	(0.6-13.3)*	10.9	(2.1-40.7)*
	Retired	1,470	65.2	(62.2-68.2)	28.3	(25.4-31.3)	1.6	(1.0-2.8)	3.4	(2.3-5.0)	1.4	(0.9-2.3)
	Unable to work	237	64.9	(57.0-72.1)	25.6	(19.2-33.2)	0.0	(0.0-0.0)	5.4	(3.1-9.5)	4.1	(1.5-10.5)*
Veteran Status	Yes	811	64.7	(60.7-68.5)	27.3	(23.8-31.2)	1.4	(0.7-2.5)*	4.3	(2.9-6.2)	2.3	(1.4-3.9)
	No	4,658	65.1	(63.0-67.2)	27.1	(25.1-29.2)	0.8	(0.6-1.2)	4.9	(3.7-6.3)	2.1	(1.2-3.7)
Diabetes	Yes	520	66.5	(61.6-71.2)	25.6	(21.3-30.3)	1.3	(0.5-3.1)*	3.9	(2.2-6.9)	2.7	(1.0-7.1)*
	No	4,958	64.9	(62.9-66.9)	27.2	(25.3-29.2)	0.9	(0.6-1.2)	4.9	(3.9-6.3)	2.0	(1.2-3.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, Treatment Can Help People with Mental Illness Lead Normal Lives†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	570	63.7	(58.6-68.6)	21.9	(17.0-27.7)	2.3	(1.2-4.3)*	6.7	(3.4-12.9)*	5.4	(1.4-18.8)*
	No	4,862	65.4	(63.4-67.4)	27.4	(25.6-29.4)	0.8	(0.5-1.1)	4.7	(3.7-6.0)	1.7	(1.2-2.4)
Asthma	Yes	723	66.5	(61.4-71.3)	25.3	(20.8-30.3)	0.7	(0.2-2.2)*	6.7	(3.9-11.2)	0.8	(0.4-1.7)*
	No	4,746	64.9	(62.8-66.8)	27.3	(25.4-29.3)	0.9	(0.7-1.3)	4.6	(3.6-5.9)	2.3	(1.4-3.8)
Arthritis	Yes	2,014	66.9	(64.2-69.5)	23.9	(21.5-26.6)	1.3	(0.7-2.1)	5.5	(3.9-7.8)	2.4	(1.4-3.9)
	No	3,447	64.5	(62.1-66.8)	28.1	(25.9-30.5)	0.8	(0.5-1.1)	4.6	(3.5-6.1)	2.0	(1.0-4.0)*
Frequent Mental Distress	Yes	479	61.4	(54.8-67.6)	23.5	(17.9-30.3)	1.2	(0.5-2.9)*	6.1	(3.2-11.3)*	7.7	(2.8-19.8)*
	No	4,926	65.4	(63.4-67.3)	27.6	(25.7-29.5)	0.9	(0.6-1.2)	4.7	(3.7-6.0)	1.5	(1.0-2.1)
Serious Psychological Distress	Yes	135	60.3	(48.3-71.2)	27.6	(16.8-41.9)	1.6	(0.3-7.7)*	4.9	(2.0-11.2)*	5.6	(1.5-18.4)*
	No	5,279	65.4	(63.4-67.2)	27.1	(25.3-29.0)	0.8	(0.6-1.2)	4.7	(3.7-6.0)	2.0	(1.2-3.3)
Receiving Treatment for Mental Health Condition	Yes	728	73.8	(68.1-78.7)	17.4	(13.3-22.4)	0.6	(0.2-1.9)*	2.9	(1.6-5.0)	5.4	(1.5-17.3)*
	No	4,746	64.0	(62.0-66.0)	28.3	(26.4-30.2)	0.9	(0.7-1.3)	5.2	(4.0-6.5)	1.7	(1.2-2.4)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
TOTAL		5,476	23.5	(21.8-25.2)	39.3	(37.4-41.2)	1.6	(1.3-2.1)	26.5	(24.9-28.1)	9.2	(8.1-10.4)
Age (years)	18-24	127	17.6	(11.6-25.7)	50.7	(41.2-60.0)	0.8	(0.1-5.4)*	24.8	(17.9-33.3)	6.2	(2.4-14.9)*
	25-34	507	19.8	(16.1-24.0)	41.9	(37.3-46.7)	0.7	(0.3-1.8)*	29.9	(25.9-34.2)	7.7	(5.4-11.0)
	35-54	1,801	22.2	(20.0-24.5)	36.8	(34.4-39.4)	2.6	(1.9-3.5)	28.4	(26.2-30.7)	10.0	(8.6-11.7)
	55+	3,010	29.4	(27.6-31.3)	36.1	(34.2-38.1)	1.5	(1.1-2.1)	23.0	(21.5-24.7)	9.9	(8.7-11.2)
Sex	Men	2,172	25.5	(23.0-28.1)	43.7	(40.8-46.6)	1.4	(0.9-2.2)	21.4	(19.3-23.8)	8.0	(6.2-10.2)
	Women	3,304	21.6	(19.6-23.7)	34.8	(32.5-37.3)	1.9	(1.4-2.5)	31.5	(29.3-33.7)	10.2	(9.1-11.4)
Race/ Ethnicity	Hispanic or Latino	220	30.3	(23.5-38.0)	33.4	(25.1-42.8)	1.9	(0.7-4.9)*	22.8	(16.0-31.3)	11.6	(7.5-17.6)
	Not Hispanic or Latino:	5,063	22.2	(20.6-23.8)	39.9	(38.0-41.9)	1.7	(1.4-2.1)	27.7	(26.0-29.4)	8.5	(7.6-9.6)
	White											
	Black or African American	21	--	--	--	--	--	--	--	--	--	--
	All Other Races	125	16.3	(8.6-28.8)*	23.7	(12.0-41.4)*	3.3	(0.5-19.9)*	27.5	(17.2-41.0)	29.2	(14.9-49.1)*
Education	<High School	261	32.2	(23.0-43.0)	34.0	(25.4-43.9)	0.6	(0.1-3.1)*	19.4	(13.2-27.5)	13.8	(7.8-23.3)
	High School Graduate (or GED)	1,641	28.4	(25.3-31.7)	36.5	(33.0-40.3)	1.7	(1.0-2.8)	23.6	(20.8-26.8)	9.8	(7.4-12.8)
	Some College	1,774	21.5	(19.0-24.2)	41.4	(38.2-44.7)	1.9	(1.3-2.6)	26.9	(24.2-29.8)	8.3	(7.0-10.0)
	College Graduate	1,794	18.3	(15.9-20.9)	40.3	(37.4-43.4)	1.7	(1.1-2.6)	30.4	(27.8-33.1)	9.3	(7.8-11.1)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

Characteristics		Sample Size	Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Annual Household Income	< \$20,000	713	20.1	(15.7-25.3)	36.6	(29.8-43.9)	0.9	(0.4-2.0)*	25.3	(20.3-31.1)	17.1	(11.4-24.9)
	\$20,000-\$34,999	962	30.1	(26.3-34.1)	35.6	(31.1-40.4)	1.7	(1.1-2.8)	24.8	(21.2-28.8)	7.8	(5.6-10.7)
	\$35,000-\$49,999	781	23.3	(19.5-27.6)	36.4	(32.0-41.0)	1.3	(0.6-2.7)*	29.8	(25.7-34.3)	9.2	(7.2-11.8)
	\$50,000-\$74,999	973	19.2	(16.2-22.6)	42.6	(38.6-46.7)	1.7	(1.0-2.7)	28.5	(25.1-32.1)	8.1	(6.2-10.5)
	\$75,000 and more	1,486	22.2	(19.6-25.1)	42.7	(39.5-45.9)	2.0	(1.2-3.2)	26.2	(23.6-28.9)	7.0	(5.7-8.5)
Employment Status	Employed	3,070	23.4	(21.3-25.7)	40.2	(37.9-42.6)	1.9	(1.4-2.7)	26.1	(24.2-28.1)	8.3	(7.1-9.7)
	Out of work	180	21.3	(13.1-32.8)	30.2	(20.5-42.2)	0.8	(0.3-2.8)*	32.5	(23.9-42.5)	15.1	(8.9-24.4)
	Homemaker	441	25.6	(20.2-31.7)	39.5	(33.4-46.0)	1.9	(1.0-3.5)*	26.5	(21.7-31.9)	6.6	(4.6-9.3)
	Student	70	10.7	(4.6-23.1)*	45.5	(30.7-61.2)	0.0	(0.0-0.0)	30.1	(19.1-44.1)	13.6	(3.8-38.5)*
	Retired	1,466	27.6	(24.9-30.4)	38.5	(35.5-41.7)	1.4	(0.8-2.3)	24.6	(22.1-27.4)	7.9	(6.5-9.6)
	Unable to work	235	20.0	(14.5-26.9)	29.6	(22.4-38.0)	1.5	(0.6-4.0)*	27.5	(20.2-36.2)	21.4	(16.1-27.9)
Veteran Status	Yes	819	22.9	(19.8-26.3)	38.9	(34.9-43.0)	1.6	(0.9-2.8)	25.2	(21.6-29.2)	11.4	(8.6-15.1)
	No	4,645	23.6	(21.8-25.5)	39.3	(37.2-41.4)	1.7	(1.3-2.2)	26.6	(24.9-28.4)	8.9	(7.7-10.2)
Diabetes	Yes	522	20.8	(17.3-24.8)	35.3	(30.4-40.5)	1.4	(0.6-3.2)*	30.5	(25.6-36.0)	12.0	(9.3-15.2)
	No	4,950	23.7	(22.0-25.5)	39.5	(37.5-41.5)	1.7	(1.3-2.1)	26.2	(24.6-27.9)	8.9	(7.8-10.2)

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

Wyoming, 2009 BRFSS, People are Caring and Sympathetic to People with Mental Illness†

		Agree strongly		Agree slightly		Neither agree nor disagree		Disagree slightly		Disagree strongly	
Characteristics	Sample Size	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Cardiovascular Disease	Yes	563	23.6 (19.0-29.0)	26.9 (22.0-32.6)	1.9 (0.9-3.9)*	27.6 (23.2-32.6)	19.9 (12.4-30.4)				
	No	4,867	23.5 (21.8-25.3)	40.2 (38.2-42.2)	1.6 (1.2-2.1)	26.4 (24.8-28.2)	8.3 (7.4-9.3)				
Asthma	Yes	722	25.3 (20.6-30.7)	30.2 (25.6-35.2)	1.2 (0.3-4.6)*	28.7 (24.7-33.0)	14.7 (11.1-19.2)				
	No	4,741	23.1 (21.5-24.9)	40.7 (38.6-42.7)	1.7 (1.3-2.1)	26.2 (24.5-28.0)	8.3 (7.2-9.5)				
Arthritis	Yes	2,011	20.8 (18.8-22.9)	36.1 (33.4-38.9)	1.9 (1.3-2.9)	29.0 (26.6-31.6)	12.2 (10.3-14.3)				
	No	3,444	24.6 (22.5-26.8)	40.3 (38.0-42.8)	1.5 (1.1-2.1)	25.6 (23.7-27.6)	7.9 (6.6-9.4)				
Frequent Mental Distress	Yes	479	17.5 (12.8-23.6)	30.4 (24.5-37.0)	1.1 (0.5-2.2)*	31.3 (26.0-37.1)	19.7 (13.0-28.6)				
	No	4,924	23.9 (22.3-25.7)	40.3 (38.3-42.4)	1.7 (1.3-2.2)	26.1 (24.4-27.8)	7.9 (7.1-8.9)				
Serious Psychological Distress	Yes	136	15.0 (7.0-29.2)*	23.6 (11.4-42.8)*	1.1 (0.3-4.5)*	40.2 (28.7-52.9)	20.1 (13.3-29.1)				
	No	5,276	23.8 (22.2-25.5)	39.6 (37.7-41.6)	1.6 (1.3-2.1)	26.2 (24.6-27.8)	8.8 (7.7-10.0)				
Receiving Treatment for Mental Health Condition	Yes	719	17.9 (14.5-21.8)	30.0 (25.2-35.4)	1.9 (1.1-3.3)	31.5 (27.3-36.1)	18.6 (13.8-24.6)				
	No	4,748	24.4 (22.7-26.3)	39.9 (37.8-41.9)	1.7 (1.3-2.2)	25.8 (24.1-27.5)	8.3 (7.3-9.4)				

† Adjusted for sex, age group, racial/ethnic group, education, and annual household income level.

* Because the relative standard error exceeds 30%, the estimate is unreliable.

-- Estimate suppressed due to small sample size.

